

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

PROTOCOL CODE: BRLATWAC

(Page 1 of 2)

DOCTOR'S ORDERS Htcm Wtkg BSA	m²	
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form		
DATE: To be given: Cycle #:		
Date of Previous Cycle:		
 Delay treatment week(s) CBC & Diff, platelets day of treatment May proceed with doses as written for weekly paclitaxel portion if labs done within 24 h: ANC greater than or equal to 1.0 x 10⁹/L, <u>Platelets greater than or equal to</u> 90 x 10⁹/L May proceed with doses as written for AC portion if labs done within 96 h: ANC greater than or equal to 1.5 x 10⁹/L, <u>Platelets greater than or equal to</u> 90 x 10⁹/L Dose modification for: Hematology Other Toxicity 		
Proceed with treatment based on blood work from		
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm 45 Minutes Prior to PACLitaxel: dexamethasone 10 mg IV in 50 mL NS over 15 minutes 30 Minutes Prior to PACLitaxel: diphenhydrAMINE 25 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) No pre-medication required (see protocol for guidelines) OR dexamethasone [] 8 mg or [] 12 mg (select one) PO 30 to 60 minutes prior to AC treatment and select ONE of the following: Ondansetron 8 mg PO 30 to 60 minutes prior to AC treatment aprepitant 125 mg PO 30 to 60 minutes prior to AC treatment netupitant-palonosetron 300 mg-0.5 mg PO 30 to 60 minutes prior to AC treatment		
Other: **Have Hypersensitivity Reaction Tray and Protocol Available for Cycles 1 to 4**		
CHEMOTHERAPY: (Note - continued over 2 pages) CYCLE #(Cycle 1-4) PACLitaxel I 80 mg/m² OR Img/m² (select one) x BSA =mg Dose Modification:% =mg/m² x BSA =mg IV in 100 to 500 mL (non-DEHP bag) NS over 1 hour once weekly x 3 weeks (use non-DEHP tubing with 0.2 micron in-line filter)		
*** SEE PAGE 2 FOR CHEMOTHERAPY CYCLES 5 TO 8 ***		
DOCTOR'S SIGNATURE:	SIGNATURE: UC:	



Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

PROTOCOL CODE: BRLATWAC

(Page 2 of 2)

DOCTOR'S ORDERS		
DATE: To be given: Cycle #:		
CHEMOTHERAPY continued		
CYCLE # (Cycle 5-8)		
DOXOrubicin 60 mg/m ² x BSA =mg Dose Modification:% =mg/m ² x BSA =mg IV push		
cyclophosphamide 600 mg/m² x BSA =mg ☐ Dose Modification:% =mg/m² x BSA =mg IV in 100 to 250 mL NS over 20 minutes to 1 hour		
RETURN APPOINTMENT ORDERS		
 Return in <u>three</u> weeks for Doctor and Cycle (Book chemo room weekly x 3 for cycles 1-4, book chemo room every 3 weeks for AC cycles 5-8, cycle 5 to start week 13) Last Cycle. Return in week(s) after last treatment. 		
CBC & Diff, Platelets prior to each treatment		
If clinically indicated: Creatinine ALT Bilirubin Muga Echocardiogram		
☐ Other tests:		
Consults:		
See general orders sheet for additional requests.		
DOCTOR'S SIGNATURE:	SIGNATURE:	
	UC:	