

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <a href="https://www.bccancer.bc.ca">www.bccancer.bc.ca</a> and according to acceptable standards of care

PROTOCOL CODE: GICPART

Page 1 of 1

DOCTOR'S ORDERS Htcm Wt	kg BSAm²
REMINDER: Please ensure drug allergies and previous bleomycin are documen	ted on the Allergy & Alert Form
DATE: To be given:	Cycle #:
Date of Previous Cycle:	
□ Delay treatment week(s) □ CBC & Diff, creatinine day of treatment  May proceed with doses as written if within 24 hours ANC greater than or equal to 1.5 x 10°/L, platelets greater than 100 x 10°/L, creatinine clearance greater than or equal to 60 mL/minute.  Dose modification for: □ Hematology □ Other Toxicity  Proceed with treatment based on blood work from	
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm	·
dexamethasone   8 mg or   12 mg (select one) PO 30 to 60 minutes prior to CISp	platin
AND select ONE of the following:  aprepitant 125 mg PO 30 to 60 minutes prior to CISplatin and ondansetron 8 mg PO 30 to 60 minutes prior to CISplatin	
netupitant-palonosetron 300 mg-0.5 mg PO 30 to 60 minutes prior to CISplatin	
If additional antiemetic required: ☐ OLANZapine ☐ 2.5 mg or ☐ 5 mg or ☐ 10 mg (select one) PO 30 to 60 minutes prior to CISplatin☐ Other:	
PREHYDRATION: 1000 mL NS over 1 hour prior to CISplatin	
TREATMENT: Begins on Day 1 of each radiotherapy course  CISplatin 60 mg/m² x BSA =mg  Dose Modification:% =mg/m² x BSA =mg  IV in 500 mL NS with 20 mEq potassium chloride, 1 g magnesium sulfate, 30 g mannitol over 1 hour on Day 1, Weeks 1 and 5.  capecitabine 825 mg/m² x BSA x (%) =mg PO BID (refer to Capecitabine Suggested Tablet Combination Table for dose rounding). The second dose should be taken 10-12 hours after the first dose. To be dispensed in appropriate weekly intervals Monday to Friday, with Saturday, Sunday and statutory holidays off, beginning on the first day of Radiation Therapy and ending on the last day of RT	
Radiation Therapy to start Week 1 x 5½ weeks Return in <u>four</u> weeks for Doctor and book chemo <b>Day 1, Week 5</b> Return in weeks for Doctor assessment during RT Return in weeks for Doctor and week for PO capecitabine Last Cycle. Return in week(s)	
CBC & Diff, creatinine weekly	
If clinically indicated:	
See general orders sheet for additional requests.	CICNATURE.
DOCTOR'S SIGNATURE:	SIGNATURE: