Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: GIEFFOXRT

DOCTOR'S ORDERS

Ht_____cm Wt_____kg BSA_____m^2

REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form

DATE: **To be given:** **Cycle(s) #:**

Date of Previous Cycle:

☐ Delay treatment _____ week(s)

☐ **CBC & Diff** day of treatment

May proceed with doses as written if within 72 hours **ANC greater than or equal to 1.5 x 10^9/L, platelets greater than or equal to 75 x 10^9/L**

Dose modification for: ☐ **Hematology** ☐ **Other Toxicity** _____

Proceed with treatment based on blood work from _____

PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm _____.

ondansetron 8 mg PO prior to treatment

dexamethasone ☐ **8 mg** or ☐ **12 mg** *(select one)* PO prior to treatment (omit if below dexamethasone IV premedication ordered)

☐ For prior oxaliplatin hypersensitivity reactions (Grade 1 or 2):
45 minutes prior to oxaliplatin: **dexamethasone 20 mg** IV in 50 mL NS over 15 minutes
30 minutes prior to oxaliplatin: **diphenhydrAMINE 50 mg** IV in NS 50 mL over 15 minutes and **famotidine 20 mg** IV in NS 100 mL over 15 minutes (Y-site compatible)

NO ice chips

☐ **Other:**

Every 2 weeks ☐ **During Radiation (Dual Modality) (3 cycles)** **or** ☐ **Post Radiation (3 cycles)**

**** Have Hypersensitivity Reaction Tray & Protocol Available****

TREATMENT: (Note – continued over 2 pages)

☐ **Repeat in two weeks**

All lines to be primed with D5W

oxaliplatin 85 mg/m^2 x BSA = _____mg
☐ Dose Modification: _____mg/m^2 x BSA = _____mg
IV in 250 to 500 mL D5W over 2 hours*

leucovorin 200 mg/m^2 x BSA = _____ mg IV in 250 mL D5W over 2 hours*

*oxaliplatin and leucovorin may be infused over same two hour period by using a Y-site connector placed immediately before the injection site

*** SEE PAGE 2 FOR FLUOROURACIL CHEMOTHERAPY ***

DOCTOR'S SIGNATURE:

SIGNATURE:

UC:

DATE:

TREATMENT: (Continued)

fluorouracil 400 mg/m² x BSA = _____ mg

☐ Dose Modification: _____mg/m² x BSA = _____mg
IV push **THEN**

fluorouracil 1600 mg/m² x BSA = _____ mg**

☐ Dose Modification: _____mg/m² x BSA = _____mg**
IV over 46 hours in D5W to a total volume of 230 mL by continuous infusion at 5 mL/h via Baxter LV5 INFUSOR

** For 3000 to 4600 mg dose, **select INFUSOR per dose range below (doses outside** dose banding range are **prepared as ordered):**

Dose Banding Range	Dose Band INFUSOR (mg)	Pharmacist Initial and Date
Less than 3000 mg	Pharmacy to mix specific dose	
3000 to 3400 mg	3200 mg	
3401 to 3800 mg	3600 mg	
3801 to 4200 mg	4000 mg	
4201 to 4600 mg	4400 mg	

RETURN APPOINTMENT ORDERS

During Radiation (Dual Modality)
Book 3 cycles of chemotherapy every 2 weeks starting Week 1 radiation.
☐ Return in **two** weeks for Doctor and Cycle _____
☐ Last Cycle with Radiation. Return in 2 weeks for Doctor and Cycle _____ .

Post Radiation
☐ Return in **two** weeks for Doctor and Cycle _____
☐ Return in **four** weeks for Doctor and Cycles _____&_____. Book treatment x 2 cycles
☐ Last Cycle Post Radiation (3 cycles completed). Return in _____ week(s)

CBC & Diff, creatinine, total bilirubin, ALT prior to each cycle

If clinically indicated:
☐ **CEA** ☐ **CA 19-9** ☐ **SCC** ☐ **ECG**
☐ **alkaline phosphatase** ☐ **albumin** ☐ **GGT** ☐ **sodium** ☐ **potassium**
☐ **INR** weekly ☐ **INR** prior to each cycle
☐ **Other tests:**
☐ **Book for PICC assessment / insertion per Centre process**
☐ **Book for IVAD insertion per Centre process**
☐ **Weekly nursing assessment for (specify concern): _____**
☐ **Radiation consult before Cycle _____ or in _____ weeks.**
☐ **See general orders sheet for additional requests.**

DOCTOR'S SIGNATURE:

SIGNATURE:

UC: