

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca/terms-of-use and according to acceptable standards of care.

PROTOCOL CODE: GIGAJCPRT

Page 1 of 2

DOCTOR'S ORDERS Htcm Wtkg BSAm²		
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form		
DATE: To be given: Cycle #:		
Date of Previous Cycle:		
 □ Delay treatment week(s) □ CBC & diff day of treatment May proceed with doses as written if within 72 hours ANC greater than or equal to 1.5 x 109/L, Platelets greater than or equal to 75 x 109/L and Creatinine Clearance greater than or equal to 60 mL/minute 		
Dose modification for: Hematology Renal Dysfunction Other Toxicity: Proceed with treatment based on blood work from		
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm		
Cycles 1 and 2, 4 and 5: dexamethasone 8 mg or 12 mg (select one) PO 30 to 60 minutes prior to CISplatin and select ONE of the following: aprepitant 125 mg PO 30 to 60 minutes prior to CISplatin on Day 1, then 80 mg PO daily on Day 2 and 3 ondansetron 8 mg PO 30 to 60 minutes prior to CISplatin		
netupitant-palonosetron 300 mg-0.5 mg PO 30 to 60 minutes prior to CISplatin		
Cycle 3: ☐ ondansetron 8 mg PO prior to chemotherapy ☐ dexamethasone 8 mg PO prior to chemotherapy ☐ prochlorperazine 10 mg PO prior to chemotherapy OR ☐ metoclopramide 10 mg PO prior to chemotherapy		
HYDRATION: 1000 mL NS over 1 hr prior to CISplatin.		
CHEMOTHERAPY: ☐ Cycle 1: 3 week cycle (weeks 1-3) pre-RT ☐ Cycle 4: 3 week cycle starting 2-4 weeks post-RT ☐ Cycle 2: 3 week cycle (weeks 4-6) pre-RT ☐ Cycle 5: 3 week cycle post-RT		
CISplatin 60 mg/m² x BSA = mg Dose Modification:% = mg/m² x BSA = mg IV in NS 500 mL with potassium chloride 20 mEq, magnesium sulfate1 g, and mannitol 30 g over 1 hour Day 1 only.		
capecitabine 1000 mg/m² x BSA x (%) = mg PO BID x 14 days (refer to <u>Capecitabine Suggested Tablet Combination Table</u> for dose rounding)		
☐ Cycle 3: Over 5 weeks (weeks 7-11)		
capecitabine 825 mg/m² x BSA x (%) = mg PO BID each RT day only. (refer to <u>Capecitabine Suggested Tablet Combination Table</u> for dose rounding) The second dose should be taken 10-12 hours after the first dose. To be dispensed in appropriate weekly intervals, Monday-Friday with Saturday, Sunday and Statutory holidays off.		
DOCTOR'S SIGNATURE: UC:		



Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca/terms-of-use and according to acceptable standards of care.

PROTOCOL CODE: GIGAJCPRT

Page 2 of 2

RETURN APPOINTMENT ORDERS	
DATE:	
 □ Return in three weeks for Doctor and chemo Cycle □ 2 or □ 5 (select one). □ Book chemo for Day 1. □ Return in three weeks for Doctor and Cycle 3 oral chemo. □ Return in weeks for Doctor assessment during RT. □ Return in weeks (2-4 weeks) after RT completed for Doctor and Cycle 4. □ Book chemo for Day 1. □ Last Cycle. Return in week(s). 	
CBC & diff, platelets, creatinine prior to each treatment	
CBC & diff, platelets, creatinine weekly during radiation therapy	
If clinically indicated: Tot. Prot Albumin Bilirubin GGT Alk Phos. LDH ALT BUN sodium potassium	
☐ INR weekly ☐ INR prior to each cycle	
☐ Other tests:	
☐ Weekly Nursing Assessment for (specify concern):	
☐ Consults:	
☐ See general orders sheet for additional requests.	
DOCTOR'S SIGNATURE:	SIGNATURE:
	UC: