

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

## PROTOCOL CODE: GUAVIPNI

(Induction)

Page 1 of 1

DOCTOR'S ORDERS Wtkg	
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form	
DATE: To be given: Cycle #:	
Date of Previous Cycle:	
<ul> <li>Delay treatment week(s)</li> <li>Delay for toxicity Type of toxicity</li> <li>May proceed with doses as written if within 96 hours ALT less than or equal to 3 times the upper limit of normal, bilirubin less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, bilirubin less than or equal to 1.5 X baseline.</li> <li>Proceed with treatment based on blood work from</li> </ul>	
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm         For prior infusion reaction:         diphenhydrAMINE 50 mg PO 30 minutes prior to treatment         acetaminophen 325 to 975 mg PO 30 minutes prior to treatment         hydrocortisone 25 mg IV 30 minutes prior to treatment	
IMMUNOTHERAPY:	
nivolumab 3 mg/kg xkg =mg every 3 weeks IV in 50 to 100 mL NS over 30 minutes using a 0.2 micron in-line filter*	
<b>ipilimumab 1 mg/kg</b> x kg =mg every 3 weeks IV in 25 to 50 mL NS over <u>30 minutes</u> using a 0.2 micron in-line filter*	
* Use separate infusion line and filter for each drug	
RETURN APPOINTMENT ORDERS	
<ul> <li>Return in <u>three weeks</u> for Doctor and Cycle (Cycle 5 maint. phase nivolumab every 2 weeks)</li> <li>Return in <u>six weeks</u> for Doctor and Cycle 5. (maint. phase nivolumab for every 4 week dose only)</li> </ul>	
CBC and diff, platelets, creatinine, alkaline phosphatase, ALT, total bilirubin, LDH, sodium, potassium, creatinine kinase, TSH, glucose prior to each treatment Weekly nursing assessment If clinically indicated: Serum HCG or urine HCG – required for woman of child bearing potential Free T3 and free T4 lipase morning serum cortisol serum ACTH levels testosterone estradiol FSH LH Other consults: See general orders sheet for additional requests.	
DOCTOR'S SIGNATURE:	SIGNATURE:
	UC: