

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

PROTOCOL CODE: GUAVNIV

(Page 1 of 1)

DOCTOR'S ORDERS Ht cm Wt kg BSA	m²
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form	
DATE: To be given: Cycle #:	
Date of Previous Cycle:	
Delay treatment week(s)	
May proceed with doses as written if within 96 hours ALT less than or equal to 3 times the upper limit of normal, <u>bilirubin less than or equal to</u> 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal and less than or equal to 1.5 X baseline.	
Proceed with treatment based on blood work from	
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm For prior infusion reaction: Image:	
TREATMENT:	
nivolumab 3 mg/kg xkg = mg (max. 240 mg) every 2 weeks	
IV in 50 to 100 mL NS over 30 minutes using a 0.2 micron in-line filter.	
RETURN APPOINTMENT ORDERS	
 Return in <u>two weeks</u> for Doctor and Cycle # Return in <u>four weeks</u> for Doctor and Cycles # and Book chemo x 2 cycles. Last cycle. Return in week(s). 	
CBC and diff, platelets, creatinine, alkaline phosphatase, ALT, total bilirubin, LDH, sodium, potassium, TSH prior to each treatment	
If clinically indicated: ECG Chest X-ray serum HCG or urine HCG (select one) – required for woman of child bearing potential Free T3 and free T4 lipase morning serum cortisol serum ACTH levels testosterone estradiol FSH LH Glucose Weekly nursing assessment	
Other consults:	
☐ See general orders sheet for additional requests.	
DOCTOR'S SIGNATURE:	SIGNATURE:
	UC: