

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: GUAVPEML

Page 1 of 2

DOCTOR'S ORDERS Htcm Wtkg BS	6Am²	
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form		
DATE: To be given:	Cycle #:	
Date of Previous Cycle:		
Delay treatment week(s)		
May proceed with pembrolizumab as written if within 96 hours ALT <u>less than or equal to</u> 3 times the upper limit of normal, <u>total bilirubin less than or equal to</u> 1.5 times the upper limit of normal, creatinine <u>less than</u> <u>or equal to</u> 1.5 times the upper limit of normal <u>and</u> <u>less than or equal to</u> 1.5 X baseline.		
May proceed with lenvatinib as written if within 96 hours ANC <u>greater than or equal to</u> 1.0 x 10 ⁹ /L, platelets <u>greater than or equal to</u> 75 x 10 ⁹ /L, BP <u>less than</u> 160/100 mmHg, creatinine clearance <u>greater than or equal to</u> 30 mL/min, alkaline phosphatase or ALT <u>less than or equal to</u> 5 X ULN, total bilirubin <u>less than or equal to</u> 3 X ULN, urine protein <u>less than</u> 1 g/24 h		
Dose modification for: Hematology Hypertension Diarrhea QTc prolongation Other Toxicity		
Proceed with treatment based on blood work from		
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm		
Antiemetics per protocol		
For prior infusion reaction to pembrolizumab: diphenhydrAMINE 50 mg PO 30 minutes prior to pembrolizumab acetaminophen 325 to 975 mg PO 30 minutes prior to pembrolizumab hydrocortisone 25 mg IV 30 minutes prior to pembrolizumab		
TREATMENT:		
pembrolizumab 2 mg/kg x kg = mg (max. 200 mg) IV in 50 mL NS over 30 minutes using a 0.2 micron in-line filter		
lenvatinib		
(select one) 14 mg PO once daily		
☐ 10 mg PO once daily		
☐ 8 mg PO once daily		
☐ 4 mg PO once daily		
Mitte: 25 days or days. Order in increments of 5 days (only available as 5-day supply unit)		
DOCTOR'S SIGNATURE:	SIGNATURE:	
	UC:	



Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: GUAVPEML

Page 2 of 2

Date:	
RETURN APPOINTMENT ORDERS	
 □ Return in three weeks for Doctor and Cycle □ Please book Nurse for BP monitoring q 2 weeks x □ Last Cycle. Return in week(s) 	
CBC & Diff, platelets, creatinine, ALT, alkaline phosphatase, total bilirubin, sodium, potassium, magnesium, calcium, albumin, TSH, dipstick or laboratory urinalysis for protein, blood pressure measurement prior to each cycle	
Every two weeks for first 2 months: ALT, alkaline phosphatase, total bilirubin, albumin	
During cycle 1 and 2: weekly telephone nursing assessment Cycle 3 onward: every weeks telephone nursing assessment for weeks	
If clinically indicated:	
☐ 24 hour urine protein within 3 days prior to next cycle for laboratory urinalysis for protein greater than or equal to 1g/L or dipstick proteinuria 2+ or 3+	
☐ ECG ☐ Chest X-ray ☐ MUGA scan or ☐ echocardiogram ☐ serum HCG or ☐ urine HCG – required for woman of child bearing potential	
☐ Free T3 and free T4 ☐ lipase ☐ morning serum cortisol	
☐ random glucose ☐ creatine kinase ☐ serum ACTH levels	
☐ testosterone ☐ estradiol ☐ FSH ☐ LH ☐ GGT ☐ total protein ☐ phosphorus ☐ C-reactive protein ☐ troponin ☐ INR	
☐ Other consults:	
☐ See general orders sheet for additional requests.	
DOCTOR'S SIGNATURE:	SIGNATURE:
	UC: