

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

PROTOCOL CODE: GUTEM

Page 1 of 1

DOCTOR'S ORDERS	
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form One cycle consists of <u>4 weeks</u> of once weekly temsirolimus	
DATE: To be given: Cycle #:	weeks
Date of Previous Cycle:	
Delay treatment week(s)	
CBC & Diff, Platelets day of treatment	
May proceed with doses as written if within 24 hours ANC greater than or equal to 1.0 x 10 ⁹ / or equal to 75 x 10 ⁹ /L	L, Platelets <u>greater than</u>
Dose modification for: Hematology Other Toxicity	
Proceed with treatment based on blood work from	
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm	
diphenhydrAMINE 25 mg or 50 mg (circle one) IV in 50 mL NS over 20 minutes prior to temsirolimus	
Other:	
** Have Hypersensitivity Reaction Tray and Protocol Available **	
CHEMOTHERAPY:	
temsirolimus 25 mg Dose Modification: 20 mg or 15 mg (circle one) IV in 250 mL NS (non-DEHP bag) over 30 minutes weekly xweeks. (use non-DEHP tubing with 0.2 micron in-line filter)	
RETURN APPOINTMENT ORDERS	
Return in week(s) for Doctor and Cycle week	
Book weekly treatment xweeks	
Last Cycle. Return in week(s).	
CBC & Diff, Platelets prior to each treatment	
Sodium, Potassium, Creatinine, BUN, Glucose, Calcium, Phosphorus, ALT, LDH, Total Bili, Alk. Phos., Total Cholesterol, Triglycerides prior to each cycle (every 4 weeks)	
If clinically indicated: 🗌 Tot. Prot 🔄 Albumin 🗌 GGT 🔛 INR/PTT	
☐ Other tests:	
Consults:	
See general orders sheet for additional requests.	
DOCTOR'S SIGNATURE:	SIGNATURE:
	UC: