

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: GOOVBEVP

Page 1 of 2

D	OCTOR'S	ORDERS	Ht	cm	Wt	kg	BSA	m²
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form								
	TE:		be given:		С	ycle #:		
Da	te of Previous Cyc	de:						PAGE 1 of 2
	☐ Delay treatment week(s) and repeat CBC & Diff, Platelets on day of treatment							
	May proceed with doses as written if within 96 hours ANC greater than or equal to 1.0 x 10 ⁹ /L, Platelets greater than or equal to 100 x 10 ⁹ /L, BP less than or equal to 150/100, and urine dipstick for protein negative or 1+.							
Dose modification for:								
PF	REMEDICATIO	ONS: Patient to take	own supply. RN/Pha	armacist to	confirm	n		
45 minutes prior to PACLitaxel: dexamethasone 20 mg IV in 50 mL NS over 15 minutes 30 minutes prior to PACLitaxel: diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible)								
	Other:							
Have Hypersensitivity Reaction Tray and Protocol Available								
CHEMOTHERAPY:								
PACLitaxel ☐ 175 mg/m² or ☐ 155 mg/m² or ☐ 135 mg/m² (select one) x BSA = mg ☐ Dose Modification: % = mg/m² x BSA = mg IV in 250 to 500 mL (non-DEHP bag) NS over 3 hours. (Use non-DEHP tubing with 0.2 micron in-line filter)								
Flush line with 10 mL NS pre-bevacizumab. Blood pressure measurement pre-bevacizumab dose.								
bevacizumab								
		•			by Policy	III-190		
		ect bevacizumab brand		emic Therap		III-190 macist Initia	al and Date	e
	Pharmacy to sele	ect bevacizumab brand	as per Provincial Syste	emic Therap			al and Date	9
	Pharmacy to sele	ect bevacizumab brand	as per Provincial Syste	emic Therap			al and Date	
	Pharmacy to sele	ect bevacizumab brand Brand (Pharmacist to	as per Provincial Syste	emic Therap			al and Date	SIGNATURE:
	Pharmacy to sele Drug bevacizumab	ect bevacizumab brand Brand (Pharmacist to	as per Provincial Syste	emic Therap			al and Date	



Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: GOOVBEVP

Page 2 of 2

DATE:	
RETURN APPOINTMENT ORDERS	
Return in three weeks for Doctor and Cycle	
Last Treatment. Return in week(s).	
CBC & Diff, Platelets, Laboratory urinalysis or Urine dipstick for protein (select one) prior to next cycle (within 96 hours OK).	
☐ 24 h urine for total protein within 3 days prior to next bevacizumab dose if 2+ or 3+ dipstick or greater than or equal to 1 g/L laboratory urinalysis for protein	
☐ INR weekly ☐ INR prior to next cycle	
Prior to next cycle, if clinically indicated: Bilirubin Alk Phos GGT ALT	
□LDH □Tot Prot □Albumin □Creatinine □CA 15-3 □CA 125 □CA 19-9 □SCC	
□CEA	
☐ Other tests:	
☐ Consults:	
☐ See general orders sheet for additional requests.	
DOCTOR'S SIGNATURE:	SIGNATURE:
	uc∙