

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: GOOVPLDC

Page 1 of 2

REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form DATE: To be given: Cycle #: Date of Previous Cycle: Delay treatment week(s) CBC & Diff, Platelets day of treatment May proceed with doses as written if within 96 hours ANC greater than or equal to 1.0 x 10°/L, Platelets greater than or equal to 100 x 10°/L Dose modification for: Hematology Other Toxicity Proceed with treatment based on blood work from PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm If prior infusion reaction: 45 minutes prior to DOXOrubicin pegylated liposomal: dexamethasone 20 mg IV in 50 mL NS over 15 minutes 30 minutes prior to DOXOrubicin pegylated liposomal: diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) dexamethasone 8 mg PO 30 to 60 minutes prior to CARBOplatin AND select ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin, and ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin, and ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin		
Date of Previous Cycle: Delay treatment week(s) CBC & Diff, Platelets day of treatment May proceed with doses as written if within 96 hours ANC greater than or equal to 1.0 x 10°/L, Platelets greater than or equal to 100 x 10°/L Dose modification for:		
Delay treatment week(s) CBC & Diff, Platelets day of treatment May proceed with doses as written if within 96 hours ANC greater than or equal to 1.0 x 10°/L, Platelets greater than or equal to 100 x 10°/L Dose modification for:		
□ CBC & Diff, Platelets day of treatment May proceed with doses as written if within 96 hours ANC greater than or equal to 1.0 x 10 ⁹ /L, Platelets greater than or equal to 100 x 10 ⁹ /L Dose modification for: □ Hematology □ Other Toxicity □ Proceed with treatment based on blood work from □ PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm □ If prior infusion reaction: 45 minutes prior to DOXOrubicin pegylated liposomal: □ dexamethasone 20 mg IV in 50 mL NS over 15 minutes 30 minutes prior to DOXOrubicin pegylated liposomal: □ diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) dexamethasone 8 mg PO 30 to 60 minutes prior to CARBOplatin AND select □ ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and		
May proceed with doses as written if within 96 hours ANC greater than or equal to 1.0 x 10°/L, Platelets greater than or equal to 100 x 10°/L Dose modification for:		
or equal to 100 x 10 ⁹ /L Dose modification for: Hematology Other Toxicity Proceed with treatment based on blood work from PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm If prior infusion reaction: 45 minutes prior to DOXOrubicin pegylated liposomal: dexamethasone 20 mg IV in 50 mL NS over 15 minutes 30 minutes prior to DOXOrubicin pegylated liposomal: diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) dexamethasone 8 mg PO 30 to 60 minutes prior to CARBOplatin AND select ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin AND select aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and		
Dose modification for:		
Proceed with treatment based on blood work from PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm If prior infusion reaction: 45 minutes prior to DOXOrubicin pegylated liposomal: dexamethasone 20 mg IV in 50 mL NS over 15 minutes 30 minutes prior to DOXOrubicin pegylated liposomal: diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) dexamethasone 8 mg PO 30 to 60 minutes prior to CARBOplatin AND select ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin ONE of the following: aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and		
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm If prior infusion reaction: 45 minutes prior to DOXOrubicin pegylated liposomal: dexamethasone 20 mg IV in 50 mL NS over 15 minutes 30 minutes prior to DOXOrubicin pegylated liposomal: diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) dexamethasone 8 mg PO 30 to 60 minutes prior to CARBOplatin AND select		
If prior infusion reaction: 45 minutes prior to DOXOrubicin pegylated liposomal: dexamethasone 20 mg IV in 50 mL NS over 15 minutes 30 minutes prior to DOXOrubicin pegylated liposomal: diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) dexamethasone 8 mg PO 30 to 60 minutes prior to CARBOplatin AND select ONE of the following: aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and		
dexamethasone 20 mg IV in 50 mL NS over 15 minutes 30 minutes prior to DOXOrubicin pegylated liposomal: diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) dexamethasone 8 mg PO 30 to 60 minutes prior to CARBOplatin AND select ONE of the following: aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and		
30 minutes prior to DOXOrubicin pegylated liposomal: diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) dexamethasone 8 mg PO 30 to 60 minutes prior to CARBOplatin AND select ONE of the following: aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and		
diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) dexamethasone 8 mg PO 30 to 60 minutes prior to CARBOplatin AND select ONE of the following: aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and		
dexamethasone 8 mg PO 30 to 60 minutes prior to CARBOplatin AND select ONE of the following: aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and		
dexamethasone 8 mg PO 30 to 60 minutes prior to CARBOplatin AND select ONE of the following: aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and		
AND select ONE of the following:		
AND select ONE of the following:		
ONE of the following: aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and		
I I following: · · ·		
following: ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin		
netupitant-palonosetron 300 mg-0.5 mg PO 30 to 60 minutes prior to CARBOplatin		
If additional antiemetic required:		
□ OLANZapine □ 2.5 mg or □ 5 mg or □ 10 mg (select one) PO 30 to 60 minutes prior to CARBOplatin		
☐ Other:		
Have Hypersensitivity Reaction Medications and Protocol Available		
CHEMOTHERAPY:		
All lines to be primed with D5W (CARBOplatin is compatible with both NS and D5W)		
DOXOrubicin pegylated liposomal 30 mg/m² or 25 mg/m² (select one) x BSA = mg		
☐ Dose Modification:mg/m² x BSA =mg		
IV in 250 mL D5W over 1 h*		
*In Cycle 1, infuse over at least 1 h (maximum 1mg/min). For subsequent doses and no prior reaction, infuse over 1 h.		
CARBOplatin AUC 5 or 4 (select one) x (GFR + 25) = mg		
☐ Dose Modification:% = mg		
IV in 100 to 250 mL NS over 30 minutes.		
DOCTOR'S SIGNATURE: SIGNATURE:		
UC:		



Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: GOOVPLDC

Page 2 of 2

DATE:		
RETURN APPOINTMENT ORDERS		
Return in four weeks for Doctor and Cycle		
Last Treatment. Return in week(s).		
Cycle 1: CBC & Diff, Platelets, Creatinine prior to Day 1, and CBC & Diff, Platelets on Days 14, and 21.		
Subsequent cycles: CBC & Diff, Platelets, Creatinine prior to Day 1;		
if indicated, also on ☐ Day 14 and/or ☐ Day 21.		
Prior to next cycle, if clinically indicated:		
☐ Bilirubin ☐ Alk Phos ☐ GGT ☐ ALT ☐ LDH		
☐ Tot Prot ☐ Albumin		
☐ CA 15-3 ☐ CA 125 ☐ CA 19-9 ☐ CEA		
☐ Other tests:		
Consults:		
☐ See general orders sheet for additional requests.		
DOCTOR'S SIGNATURE:	SIGNATURE:	
	UC:	