

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

PROTOCOL CODE: UGOCXCATBP

(Page 1 of 3) A BC Cancer "Compassionate Access Program" request form must be completed and approved prior to treatment

REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form DATE: Cycle #: Date of Previous Cycle: □ Delay treatment week(s)	DOCTOR'S	ORDERS	Ht	cm	Wt	kg	BSA	m²
Date of Previous Cycle: Delay treatment	REMINDER: Please	ensure drug allergie	s and previous blee	omycin are d	ocumente	ed on the All	ergy & Al	ert Form
□ Delay treatment	DATE:		To be give	n:			Cycle	ə #:
□ CBC & Diff on day of treatment May proceed with doses as written if within 96 hours ANC greater than or equal to 1.0 x 10%L, Platelets greater than or equal to 1.5 times the baseline, ALT less than or equal to 3 times the upper limit of normal, and less than or equal to 1.5 times the baseline, ALT less than or equal to 3 times the upper limit of normal, total bilirubin less than or equal to 1.5 times the upper limit of normal, by less than or equal to 1.5 times the upper limit of norma	Date of Previous	Cycle:						
greater than or equal to 10 x 10 ⁹ /L, creatinine less than or equal to 1.5 times the upper limit of normal, local billinut biness than or equal to 1.5 times the baseline, ALT less than or equal to 1.5 times the upper limit of normal, local billinut biness than or equal to 1.5 times the upper limit of normal, BP less than or equal to 1.5 times the upper limit of normal, local billinut biness than or equal to 1.5 times the upper limit of normal, BP less than or	•	()			_		_	
Proceed with treatment based on blood work from	May proceed with doses as written if within 96 hours ANC <u>greater than or equal to</u> 1.0 x 10 ⁹ /L, Platelets <u>greater than or equal to</u> 100 x 10 ⁹ /L, creatinine <u>less than or equal to</u> 1.5 times the upper limit of normal <i>and</i> <u>less than or equal to</u> 1.5 times the baseline, ALT <u>less than or equal to</u> 3 times the upper limit of normal, <u>total bilirubin less than or equal to</u> 1.5 times the upper limit of normal,							
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm ¬No prior infusion reaction to pembrolizumab: administer premedications as sequenced below <u>45 minutes prior to PACLitaxel:</u> dexamethasone 20 mg IV in 50 mL NS over 15 minutes <u>30 minutes prior to PACLitaxel:</u> diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) Prior infusion reaction to pembrolizumab: administer PACLitaxel premedications prior to pembrolizumab: dexamethasone 20 mg IV in 50 mL NS over 15 minutes <u>45 minutes prior to pembrolizumab:</u> dexamethasone 20 mg IV in 50 mL NS over 15 minutes <u>30 minutes prior to pembrolizumab:</u> dexamethasone 20 mg IV in 50 mL NS over 15 minutes <u>30 minutes prior to pembrolizumab:</u> diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over <u>15 minutes of the pembrolizumab:</u> diphenhydrAMINE 50 mg PO 30 minutes prior to pembrolizumab <u>AND</u> select <u>0 ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin</u> <u>aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin</u> <u>0LANZapine 0 2.5 mg or 0 5 mg or 0 10 mg (select one) PO 30 to 60 minutes prior to CARBOplatin If additional antiemetic required: <u>0 LANZapine 2.5 mg or 5 mg or 10 mg (select one) PO 30 to 60 minutes prior to CARBOplatin <u>0 CARBOplatin </u> </u></u>	Dose modification	for: 🗌 Hema	itology	Other To	cicity			
 No prior infusion reaction to pembrolizumab: administer premedications as sequenced below	Proceed with trea	atment based on	blood work from	m				
45 minutes prior to PACLitaxel: dexamethasone 20 mg IV in 50 mL NS over 15 minutes 30 minutes prior to PACLitaxel: diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) Prior infusion reaction to pembrolizumab: administer PACLitaxel premedications prior to pembrolizumab: dexamethasone 20 mg IV in 50 mL NS over 15 minutes 30 minutes prior to pembrolizumab: dexamethasone 20 mg IV in 50 mL NS over 15 minutes 30 minutes prior to pembrolizumab: dexamethasone 20 mg IV in S0 mL NS over 15 minutes 30 minutes prior to pembrolizumab: dexamethasone 20 mg IV in NS 50 mL over 15 minutes 30 minutes prior to pembrolizumab: dexamethasone 20 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) acetaminophen 325 to 975 mg PO 30 minutes prior to pembrolizumab AND select ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin If additional antiemetic required: OLANZapine 2.5 mg or 5 mg or 10 mg (select one) PO 30 to 60 minutes prior to CARBOplatin Other: Continued on page 2	PREMEDICATI	DNS: Patient to	take own supply.	RN/Pharm	acist to	confirm		
AND ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin select aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and ONE of the ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin following: netupitant-palonosetron 300 mg-0.5 mg PO 30 to 60 minutes prior to CARBOplatin If additional antiemetic required: OLANZapine 2.5 mg or 5 mg or 10 mg (select one) PO 30 to 60 minutes prior to CARBOplatin Other: Continued on page 2	 <u>45 minutes prior to PACLitaxel:</u> dexamethasone 20 mg IV in 50 mL NS over 15 minutes <u>30 minutes prior to PACLitaxel:</u> diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) Prior infusion reaction to pembrolizumab: administer PACLitaxel premedications prior to pembrolizumab <u>45 minutes prior to pembrolizumab:</u> dexamethasone 20 mg IV in 50 mL NS over 15 minutes <u>30 minutes prior to pembrolizumab:</u> dexamethasone 20 mg IV in 50 mL NS over 15 minutes <u>30 minutes prior to pembrolizumab:</u> diphenhydrAMINE 50 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 							
select aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin Image: Ima	acetaminophen 325 to 975 mg PO 30 minutes prior to pembrolizumab							
ONE of the following: aprepriant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin If additional antiemetic required: netupitant-palonosetron 300 mg-0.5 mg PO 30 to 60 minutes prior to CARBOplatin If additional antiemetic required: 2.5 mg or 5 mg or 10 mg (select one) PO 30 to 60 minutes prior to CARBOplatin Other: Continued on page 2		ondansetron 8	mg PO 30 to 60	minutes pri	or to CA	RBOplatin		
Image:	ONE of the	• • •						
 OLANZapine 2.5 mg or 5 mg or 10 mg (select one) PO 30 to 60 minutes prior to CARBOplatin Other: Continued on page 2 						CARBOplatin		
	□ OLANZapine □ 2.5 mg or □ 5 mg or □ 10 mg (select one) PO 30 to 60 minutes prior to CARBOplatin □ Other:							
UC:	DOCTOR'S SIG	NATURE:						



PROTOCOL CODE: UGOCXCATBP

(Page 2 of 3) A BC Cancer "Compassionate Access Program" request form must be completed and approved prior to treatment

DOCTOR'S	DRDERS Page 2 of 3					
DATE:	To be given:					
Have Hypersensitivity Reaction Tray and Protocol Available						
TREATMENT:						
pembrolizumab 2 mg/kg x kg = mg (max. 200 mg) IV in 50 mL NS over 30 minutes using a 0.2 micron in-line filter*						
PACLitaxel ☐ 175 mg/m ² x BSA = mg ☐ Dose Modification:% = mg/m ² x BSA = mg IV in 250 to 500 mL (non-DEHP bag) NS over 3 hours. (Use non-DEHP tubing with 0.2 micron in-line filter*)						
CARBOplatin AUC 5 x (GFR + 25) x = mg ☐ Dose Modification:% = mg IV in 100 to 250 mL NS over 30 minutes.						
Blood pressure me	easurement pre-bevacizumab dose.					
bevacizumab 15 mg/kg or mg/kg (select one) x kg = mg IV in 100 to 250 mL NS over 1 hour (If no infusion reaction observed in Cycle 1, may administer subsequent cycles over 30 minutes). (Blood pressure measurement post-bevacizumab infusion for first 3 cycles)						
Pharmacy to select bevacizumab brand as per Provincial Systemic Therapy Policy III-190						
Drug	Brand (Pharmacist to complete. Please print.)	Pharmacist Initial and Da	ite			
bevacizumab						
* use separate infusion line and filter for each drug						
DOCTOR'S SIG	NATURE:		SIGNATURE: UC:			



Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

PROTOCOL CODE: UGOCXCATBP

(Page 3 of 3)

A BC Cancer "Compassionate Access Program" request form must be completed and approved prior to treatment

DATE:	Page 3 of 3	
RETURN APPOI	NTMENT ORDERS	
Return in <u>three</u> weeks for Doctor and Cyc Last Cycle. Return in three weeks for G pembrolizumab with or without bevacizuma	GOCXBP or GOCXBP6 (to continue	
CBC & Diff creatinine, ALT, alkaline pho potassium, TSH, dipstick or laboratory u measurement prior to each cycle.	• • • •	
24 hr urine for total protein within 3 d 3+ dipstick or greater than or equal to 1 g/L	ays prior to next bevacizumab dose if 2+ or . laboratory urinalysis for protein	
INR weekly INR prior to next cycl	e	
If clinically indicated: 🗌 ECG 🗌 Chest X-	ray	
serum HCG or urine HCG – require	•	
☐ Free T3 and free T4 ☐ lipase ☐ ☐ GGT ☐ total protein ☐ albumin	morning serum cortisol 🛛 🗌 Glucose	
serum ACTH levels testosterone		
Weekly nursing assessment		
Other consults		
See general orders sheet for additio	nal requests.	
DOCTOD'S SIGNATURE.		
DOCTOR'S SIGNATURE:		SIGNATURE:
		UC: