

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <a href="https://www.bccancer.bc.ca/terms-of-use">www.bccancer.bc.ca/terms-of-use</a> and according to acceptable standards of care.

PROTOCOL CODE: HNNAVPG

Page 1 of 1

DOCTOR'S ORDERS	Ht	cm	Wt	kg	BSA	m²
REMINDER: Please ensure drug allergies	and previous bleomy	cin are	documen	ted on the	Allergy	& Alert Form
DATE: To	be given:		C	ycle #:		
Date of Previous Cycle:						
□ Delay treatment week(s) □ CBC & Diff, Platelets day of treatment  May proceed with doses as written if within 24 hours ANC greater than or equal to 1.0 x 10°/L, Platelets greater than or equal to 100 x 10°/L, and if ordered, Creatinine Clearance greater than or equal to 50 mL/minute (for CISplatin only)  Dose modification for: □ Hematology □ Other Toxicity  Proceed with treatment based on blood work from						
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm						
dexamethasone    8 mg or    12 mg (select one) PO 30 to 60 minutes prior to CISplatin or CARBOplatin on Day 1 and Day 8 and select ONE of the following:  aprepitant 125 mg PO 30 to 60 minutes prior to CISplatin or CARBOplatin on Day 1 and Day 8 ondansetron 8 mg PO 30 to 60 minutes prior to CISplatin or CARBOplatin on Day 1 and Day 8  netupitant-palonosetron 300 mg-0.5 mg PO 30 to 60 minutes prior to CISplatin or CARBOplatin on Day 1 and Day 8  Other:						
**Have Hypersensitivity Reaction Tray and Protocol Available**						
PREHYDRATION: 1000 mL NS IV over 60 minutes prior to CISplatin on Day 1 and Day 8						
CHEMOTHERAPY:  gemcitabine 1000 mg/m² x BSA = mg  Dose Modification: % = mg/m² x BSA = mg  IV in 250 mL NS over 30 minutes on Days 1, 8 and 15						
CISplatin 50 mg/m² x BSA = mg  Dose Modification: % = mg/m² x BSA = mg  IV in 500 mL NS, with potassium chloride 10 mEq and magnesium sulfate 1 g, mannitol 30 g over 1 hour on Days 1 and 8.  OR						
CARBOplatin AUC  5 or 6 (select one) x (GFR + 25) = mg IV in 100 to 250 mL NS over 30 minutes Day 1 (if using AUC 6, may dose gemcitabine at 800 mg/m²)						
DOSE MODIFICATION FOR DAY 8, DAY gemcitabine 1000 mg/m² x BSA x (%)	= mg IV in 250					
Dose Modification: (%) =	mg/m² x BS	A =	mg	J IV in 250 n	nL NS ov	er 30 minutes
RETURN APPOINTMENT ORDERS						
☐ Return in <u>four</u> weeks for Doctor and Cycle _ ☐ Last Cycle. Return in week(s).	Book chemo D	ау 1, 8 а	nd 15.			
CBC & Diff, creatinine, ALT, total bilirubin process.  CBC & Diff, creatinine prior to Day 8  CBC & Diff prior to Day 15  Other tests:  Consults:  See general orders sheet for additional recognitions.	·					
DOCTOR'S SIGNATURE:				SIGNAT	URE:	
				UC:		