

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

## PROTOCOL CODE: LUAVNIV4

Page 1 of 1

DOCTOR'S ORDERS Htcm Wtkg BSA_	m²
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the	e Allergy & Alert Form
DATE: To be given: Cycle #:	
Date of Previous Cycle:	
Delay treatment week(s)	
May proceed with doses as written if within 96 hours ALT less than or equal to 3 times the upper limit of normal, bilirubin less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal and less than or equal to 1.5 X baseline.	
Proceed with treatment based on blood work from	
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm         For prior infusion reaction:         Image: Display the section of the section	·
hydrocortisone 25 mg IV 30 minutes prior to treatment	
TREATMENT:         nivolumab 6 mg/kg xkg = mg (max. 480 mg) every 4 weeks         IV in 50 to 100 mL NS over 30 minutes using a 0.2 micron in-line filter.	
RETURN APPOINTMENT ORDERS	
<ul> <li>Return in <u>four weeks</u> for Doctor and Cycle #</li> <li>Last cycle: Return in week(s).</li> </ul>	
CBC and diff, platelets, creatinine, alkaline phosphatase, ALT, total bilirubin, LDH, sodium, potassium, TSH prior to each treatment	
If clinically indicated: ECG Chest X-ray serum HCG or urine HCG – required for woman of child bearing potential Free T3 and free T4 IIpase morning serum cortisol serum ACTH levels testosterone estradiol FSH LH Glucose Weekly nursing assessment Other consults:	
See general orders sheet for additional requests.	
DOCTOR'S SIGNATURE:	SIGNATURE:
	UC: