

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

PROTOCOL CODE: LUMMIPNI

Page 1 of 1

DOCTOR'S ORDERS	Ht	cm	Wt	kg	BSA	m²
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form						
DATE:	To be given:			Сус	:le #:	
Date of Previous Cycle:						
Delay treatment week(s)						
May proceed with doses as written if within 96 hours ALT less than or equal to 3 times the upper limit of normal, bilirubin less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal, creatinine less the upper limit of normal,						
Proceed with treatment based on bloc	od work from					
PREMEDICATIONS: Patient to take of For prior infusion reaction:	inutes prior to trea 30 minutes prior to	tment o treatmer		firm		
TREATMENT:						
nivolumab 3 mg/kg xkg = mg (max. 240 mg) IV in 50 to 100 mL NS over 30 minutes using a 0.2 micron in-line filter* on Days 1, 15, and 29						
<pre>ipilimumab 1 mg/kg xkg = mg IV in 25 to 100 mL NS over 30 minutes using a 0.2 micron in-line filter* on Day 1 only</pre>						
* Use separate infusion line and filter for each drug						
RETURN APPOINTMENT ORDERS						
Return in <u>six weeks</u> for Doctor and C	ycle Bool	k chemo D	ay 1, 1	5, and 29.		
Last cycle. Return in week(s)).					
CBC and differential, platelets, creatin LDH, sodium, potassium, TSH, creatin					٦,	
☐ serum HCG or ☐ urine HC ☐ Free T3 and free T4 ☐ lipa	•	erum cor	isol	aring potential		
Weekly nursing assessment						
☐ Other consults:						
☐ See general orders sheet for addit	ional requests.					
DOCTOR'S SIGNATURE:						SIGNATURE:
						UC: