

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca/terms-of-use</u> and according to acceptable standards of care.

PROTOCOL CODE: LYPEM6

Page 1 of 1

DOCTOR'S ORDERS	Ht	cm	Wt	kg	BSA_	m²
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form						
DATE:	To be given:			Сус	:le #:	
Date of Previous Cycle:						
Delay treatment week(s)						
May proceed with doses as written if within 96 hours ALT <u>less than or equal to</u> 3 times the upper limit of normal, <u>bilirubin less than or equal to</u> 1.5 times the upper limit of normal, creatinine <u>less than or equal to</u> 1.5 times the upper limit of normal and <u>less than or equal to</u> 1.5 times the baseline. Proceed with treatment based on blood work from						
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm For prior infusion reaction: Image:						
TREATMENT:						
pembrolizumab 4 mg/kg x kg =mg (max. 400 mg) IV in 50 mL NS over 30 minutes using a 0.2 micron in-line filter						
RETURN APPOINTMENT ORDERS						
Return in <u>six weeks</u> for Doctor and C	Cycle					
Last cycle. Return in week(s)					
CBC & Diff, creatinine, alkaline phosp potassium, TSH prior to each treatment		ıl bilirubir	n, LDH, s	odium,		
If clinically indicated: ECG che serum HCG or urine HCG - r free T3 and free T4 lipase serum ACTH levels testoste C-reactive protein creatinin HBV viral load	equired for woman morning serue more estradio	m cortiso	I		cose	
Weekly nursing assessment						
Other consults:						
☐ See general orders sheet for addit	tional requests.					
DOCTOR'S SIGNATURE:						SIGNATURE:
						UC: