

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: SMAVALIPNI (Induction)

Page 1 of 1

DOCTOR'S ORDERS Wtkg	
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form	
DATE: To be given: Cycle #:	
Date of Previous Cycle:	
□ Delay treatment week(s) □ Delay for toxicity Type of toxicity May proceed with doses as written if within 96 hours ALT less than or equal to 3 times the upper limit of normal, bilirubin less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal and less than or equal to 1.5 X baseline. Proceed with treatment based on blood work from	
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm	
For prior infusion reaction: diphenhydrAMINE 50 mg PO 30 minutes prior to treatment acetaminophen 325 to 975 mg PO 30 minutes prior to treatment hydrocortisone 25 mg IV 30 minutes prior to treatment	
IMMUNOTHERAPY:	
nivolumab 3 mg/kg xkg =mg every 3 weeks IV in 50 to 100 mL NS over 30 minutes using a 0.2 micron in-line filter*	
<pre>ipilimumab 1 mg/kg x kg = mg every 3 weeks IV in 25 to 100 mL NS over 30 minutes using a 0.2 micron in-line filter* * Use separate infusion line and filter for each drug</pre>	
RETURN APPOINTMENT ORDERS	
Return in three weeks for Doctor and Cycle (Cycle 5 maint. phase nivolumab q2wk) Return in six weeks for Doctor and Cycle 5. (maint. phase nivolumab for q4wk dose only)	
CBC & Diff, creatinine, alkaline phosphatase, ALT, total bilirubin, LDH, sodium, potassium, creatine kinase (CK), TSH, glucose prior to each treatment	
Weekly nursing assessment	
If clinically indicated: ECG	
DOCTOR'S SIGNATURE:	SIGNATURE:
	UC: