

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

PROTOCOL CODE: SMAVIPI

Page 1 of 1

DOCTOR'S ORDERS	Ht	cm	Wt	kg
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form				
DATE: To be given: Cycle #		of 4		
Date of Previous Cycle:				
 Delay treatment week(s) for: Hepatotoxicity Other Toxicity: May proceed with doses as written if within 96 hours AST or ALT less than or equal to 2.5 times the upper limit of normal, total bilirubin less than or equal to 1.5 times the upper limit of normal Proceed with treatment based on blood work from 				
TREATMENT:				
ipilimumab 3 mg/kg x kg = mg IV in 50 to 250 mL NS over 30 minutes using a 0.2 micron in-line filter.				
RETURN APPOINTMENT ORDERS				
☐ Return in <u>three</u> weeks for Doctor and Cycle #				
Last Treatment. Return in week((s)			
CBC & Diff, creatinine, alkaline phosphatase, ALT, total bilirubin, LDH, sodium, potassium, TSH, creatine kinase prior to each treatment.				
During treatment: weekly telephone nursing assessment After treatment: every weekly telephone nursing assessment for weeks				
If clinically indicated:				
serum HCG or urine HCG – required for woman of child bearing potential				
☐ free T3 and free T4 ☐ lipase ☐ morning serum cortisol ☐ glucose				
☐ serum ACTH levels ☐ testosterone ☐ es ☐ troponin	stradiol _	FSH 🗌 L	H 🗌 ECG	
Other Tests:				
Consults:				
──				
DOCTOR'S SIGNATURE:			SIGNATURE:	
				UC: