

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

PROTOCOL CODE: SMAVNIV4

(Page 1 of 1)

DOCTOR'S ORDERS	Ht	cm Wt	kg BSA_	m²
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form				
DATE:	To be given:		Cycle #:	
Date of Previous Cycle:				
Delay treatment week(s)				
May proceed with doses as written if w <u>bilirubin less than or equal to</u> 1.5 til upper limit of normal <i>and</i> <u>less than or</u>	mes the upper limi	t of normal, creatini		
Proceed with treatment based on b	lood work from			
PREMEDICATIONS: Patient to take of For prior infusion reaction: diphenhydrAMINE 50 mg PO 30 acetaminophen 325 to 975 mg P	minutes prior to tre	atment		·
hydrocortisone 25 mg IV 30 min	utes prior to treatme	ent		
nivolumab 6 mg/kg xkg = mg (max. 480 mg) every 4 weeks IV in 50 to 100 mL NS over 30 minutes using a 0.2 micron in-line filter. RETURN APPOINTMENT ORDERS				
			-	
Return in <u>four weeks</u> for Doctor an	ıd Cycle #			
Last cycle. Return in week	α(s).			
CBC & Diff, creatinine, alkaline pho potassium, TSH, creatine kinase pri	•		odium,	
If clinically indicated: ECG C serum HCG or urine HCG - re Free T3 and free T4 lipase serum ACTH levels testoste troponin Weekly nursing assessment Other consults: See general orders sheet for ad	equired for woman o morning serum o erone estradio	ortisol		
DOCTOR'S SIGNATURE:				SIGNATURE:
				UC: