

PROTOCOL CODE: SCDRUGRX

**Preprinted Order A: Immediate management of infusion-related reaction and resumption of infusion**

**DOCTOR'S ORDERS**

**DATE:**

**TIME:**

**Cycle #:**

Infusion-Related Reaction Management:

**Stop infusion** and administer the following:

- |   |  |
|---|--|
| <input type="checkbox"/> diphenhydrAMINE 50 mg IV   | <input type="checkbox"/> epinephrine 0.5 mg* IM STAT                       |
| <input type="checkbox"/> hydrocortisone sodium succinate 100 mg IV  | Repeat at 5 minute intervals twice more as needed                          |
| <input type="checkbox"/> salbutamol 100 mcg inhaler, 2 puffs via aerosol chamber  | *Note: epinephrine 1:1,000 = 1 mg/mL and epinephrine 1:10,000 = 1 mg/10 mL |
| <input type="checkbox"/> ipratropium 20 mcg/salbutamol 100 mcg (COMBIVENT RESPIMAT) inhaler, 1 puff via aerosol chamber | <input type="checkbox"/> salbutamol 5 mg via nebulizer**                   |
| <input type="checkbox"/> ipratropium 20 mcg inhaler, 2 puffs via aerosol chamber  | <input type="checkbox"/> ipratropium 0.5 mg via nebulizer**                |

\*\* See Procedure: [Application of COVID-19 Personal Protective Equipment \(PPE\) Framework for BC Cancer](#)

Others:

SCDRUGRX: Use for drugs that have no specific direction in protocol

- Resume \_\_\_\_\_ (indicate drug name) infusion at 25% OR \_\_\_\_% of the rate at which the reaction occurred and titrate as per SCDRUGRX or cap rate at \_\_\_\_\_ once symptoms of reaction have subsided
- 25% of the rate at time of reaction for 5 minutes
  - 50% of the rate at time of reaction for 5 minutes
  - 75% of the rate at time of reaction for 5 minutes
  - 100% of the rate at time of reaction
- DO NOT resume infusion

SPECIFIC PROTOCOL: Refer to patient specific protocol for infusion reaction guidelines

- Resume \_\_\_\_\_ (indicate drug name) as per patient's treatment protocol.
- Resume PACLitaxel infusion at 20 mL/h for 5 minutes, 30 mL/h for 5 minutes, 40 mL/h for 5 minutes, then 60 mL/h for 5 minutes. If no reaction, increase to full rate.
- Resume DOCEtaxel infusion at 30 mL/h x 5 minutes, then 60 mL/h x 5 minutes, then 120 mL/h x 5 minutes, then complete infusion at 250 mL/h (for 500 mL bag, continue 250 mL/h for 5 minutes, then complete infusion at 500 mL/h).
- Resume riTUXimab at \_\_\_\_\_ (one infusion rate below the rate at which the reaction occurred) and continue with escalation of infusion rates as per protocol.
- Resume daratumumab at \_\_\_\_\_ (half the infusion rate at which the reaction occurred) and continue with escalation of infusion rates as per protocol.
- DO NOT resume infusion

SPECIFIC GUIDELINES: Physician to provide specific orders for resuming infusion

DOCTOR'S SIGNATURE:

SIGNATURE:

UC: