

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Aldesleukin 22 million units (1.3 mg) (Novartis) (F)(PFL) no preservative ¹	1.2 mL SWI ^{1,2} direct diluent against side of vial during reconstitution ¹ do not shake ¹	18 million unit/mL (1.1 mg/mL) ^{1,2}	48 h F ¹	50 mL D5W ¹ 30 – 70 mcg/mL ¹ Less than 30 mcg/mL: dilute in D5W containing human albumin 0.1% ²	48 h F ¹	- do not use in-line filter ^{1,2} - avoid bacteriostatic water for injection or NS due to increased aggregation ¹
				SC syringe ^{3,4}	14 d F ⁴ **(PFL)	
Alemtuzumab 30 mg/mL (Genzyme previously Bayer) ⁵ (F)(PFL) do not shake no preservative ⁶	N/A	filter NOT required ⁶ 30 mg/mL ⁶	discard unused portion ⁶	SC syringe ⁷	discard at the end of the day F or RT	- do not shake ⁸
				100 mL NS or D5W ⁶	8 h F or RT ⁶ **(PFL) ⁸	
				100 mL NS or D5W ⁹	8 h F or RT ⁸ **(PFL) ⁸	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Amifostine 500 mg (MedImmune) (RT) no preservative ¹⁰	9.7 mL NS only ¹⁰	50 mg/mL ¹⁰	24 h F, 5 h RT ¹⁰	25–50 mL * NS only ¹⁰	5–40 mg/mL: 24 h F, 5 h RT ¹⁰	- discard cloudy solution ¹¹
Amsacrine 75 mg/1.5 mL (Erfa Canada) (RT) no preservative ¹²	glass syringes preferred during reconstitution; max. time in plastic syringe ¹² : 15 min 13.5 mL supplied diluent (L-lactic acid) ¹ transfer 1.5mL from ampoule into the diluent vial ¹²	5 mg/mL ¹²	24 h RT ¹² PFL ¹²	500 mL D5W ¹² (plastic or glass container) ¹²	7 d F, 48 h RT ¹²⁻¹⁴	- contains DMA***
Arsenic 10 mg/10 mL (Lundbeck/Teva) (RT) no preservative ¹⁵	N/A	1 mg/mL ¹⁵ (use filter needle to withdraw from ampoule)	discard unused portion ¹⁵	100-250 mL NS , D5W ¹⁵	24 h RT, 48 h F ¹⁵	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Asparaginase (asparaginase <i>E. coli</i>) 10,000 units (CGF/EUSA) (F) no preservative{[12004]}	4 mL SWI ¹⁶ do not shake; rotate gently ¹⁶	2500 units/mL	3 h RT or 72 h F ¹⁶	syringe	complete administration within 3 h RT or 72 h F ¹⁶	
				50-250 mL NS or D5W ¹⁷	complete administration within 3 h RT ^{16,18}	
Erwinia asparaginase (asparaginase <i>Erwinia</i> <i>chrysanthemi</i>) 10,000 units (CGF/EUSA) (F) no preservative ¹⁹	1-2 mL NS ¹⁹ do not shake; mix gently to minimize bubbles and contact with stopper ¹⁹	10 000-5000 units/mL (use 5 micron filter needle to withdraw from vial) ²⁰	15 min RT ¹⁹	glass or polypropylene syringe ¹⁹	4 h RT ¹⁹	- contact with the rubber stopper may denature the reconstituted drug, creating filaments of insoluble material ¹⁹ - discard if particulate matter is present ²⁰ - do not use sterile water for reconstitution as the resulting product is not isotonic ¹⁹
PEG-asparaginase - see pegaspargase in L-Z chart (pegylated asparaginase <i>E. coli</i>)						

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Atezolizumab 1200 mg/20 mL (Hoffman-La Roche) (F)(PFL) do not shake no preservative ²¹	N/A	60 mg/mL ²¹	discard unused portion ²¹	250 mL NS only ²²	complete administration within 8 h RT, 24 h F ²¹	- discard vial if cloudy, discoloured (should be clear to pale yellow), or visible particles ²² - do NOT shake; mix by slow inversion ²²
Avelumab 200 mg (Merck Serono) (F)(PFL) do not shake no preservative ²³	N/A	20 mg/mL ²³	discard unused portion ¹⁸	250 mL NS , 0.45% sodium chloride ²⁴ dilute to final volume by withdrawing volume from bag equal to volume of drug to be added ²⁴ mix by gentle inversion ²⁴	complete administration within 8 h RT, 24 h F ²⁴	- if refrigerated, allow vials/product to come to RT prior to use ²⁴ - do not shake ²⁴ - use 0.2 micron in- line filter ²⁴

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
azaCITIDine 100 mg (Celgene) (RT) no preservative ²⁵	4 mL SWI ²⁵ shake vigorously ²⁵ record time of reconstitution	25 mg/mL ²⁵	45 min RT, 8 h F ²⁵	SC syringe ²⁵	45 min RT (including preparation time), 8 h F ²⁵ refrigerate syringe immediately after preparation if not to be used within 45 minutes of reconstitution ²⁶	- discard if contains large particles ²⁵ - re-suspend syringe contents before injection by vigorously rolling syringe between palms ²⁵ -if cold diluent reconstitution is used to extend stability, minimize exposure to RT; ensure proper refrigeration of diluent, reconstituted vial, and final product
	cold diluent reconstitution: 4 mL SWI at 2- 8°C ^{27,28}	25 mg/mL ²⁵	22 h F ^{27,28}		22 h F ^{27,28}	
					Refrigerated syringes²⁵: <ul style="list-style-type: none"> • allow up to 30 min prior to administration to reach a temperature of ~20- 25°C • discard syringe if time elapsed at RT is greater than 30 min 	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p>azaCITIDine 100 mg (Dr. Reddy's) (RT) no preservative²⁹</p>	<p>4 mL SWI²⁹ shake vigorously²⁹</p>	<p>25 mg/mL²⁹</p>	<p>45 min RT, 8 h F²⁹</p>	<p>SC syringe²⁹</p>	<p>45 min RT (including preparation time), 8 h F²⁹</p> <p>refrigerate syringe immediately after preparation if not to be used within 45 minutes of reconstitution²⁹</p> <p>Refrigerated syringes²⁹:</p> <ul style="list-style-type: none"> • allow up to 30 min prior to administration to reach a temperature of approximately 20- 25°C • discard syringe if time elapsed at RT is greater than 30 min 	<ul style="list-style-type: none"> - do not filter²⁹ - discard if contains large particles²⁹ - re-suspend syringe contents before injection by vigorously rolling syringe between palms²⁹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p>BCG intravesical 81 mg (Sanofi Pasteur) (F)(PFL) preservative³⁰</p>	<p>do not shake; roll to reconstitute³⁰</p> <p>3 mL supplied diluent³⁰</p> <p>record time of reconstitution</p>	<p>10.5 ± 8.7×10⁸ CFU/vial (Connaught strain)³⁰</p>	<p>2 h F, RT³⁰</p>	<p>50 mL NS³⁰</p>	<p>2 h F or RT after reconstitution³⁰</p> <p>** (PFL)³⁰</p>	<p>- auxiliary label: biohazard¹⁸</p>
<p>BCG (Tice substrain) intravesical 50 mg = 1 to 8 x 10⁸ CFU (Hospira/Organon) (F)(PFL) no preservative³¹</p>	<p>1 mL preservative free NS for injection³¹</p> <p>use reconstitution device provided</p> <p>allow to stand for a few minutes, then gently swirl to suspend³¹</p>	<p>1 to 8×10⁸ CFU/vial³¹</p>	<p>2 h F (PFL)³¹</p>	<p>transfer from vial to 60 mL syringe, rinse vial with another 1 mL NS. Add rinse to same 60 mL syringe. qs to 50 mL with NS³¹</p>	<p>2 h F³¹</p>	<p>- auxiliary label: biohazard¹⁸ - overfill unknown - protect from light³¹ - do not filter³¹</p>

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Bendamustine 25 mg 100 mg (Lundbeck/Teva) (RT)(PFL) no preservative ³²	25 mg vial: add 5 mL SWI ³² 100 mg vial: add 20 mL SWI ³² shake well; dissolves completely in 5 minutes ³²	5 mg/mL ³²	30 minutes ³²	500 mL NS ³² 0.2-0.6 mg/mL ³²	complete administration within 24 h F, 3 h RT ³²	
Bevacizumab 100 mg/4 mL 400 mg/16 mL (Roche) (F)(PFL) do not shake no preservative ³³	N/A	25 mg/mL ³³	discard unused portion ³³	1.4-16.5 mg/mL ³⁴ 100-250 mL NS <u>only</u> ^{33,34}	48 h F, RT ³³⁻³⁵	- do not shake ³³
Bleomycin 15 units (NB: dose in units only) (Bristol) (F) no preservative ³⁶	6 mL* NS ³⁶	2.5 units/mL	48 h F ³⁶	50 mL* NS ³⁶	24 h RT ³⁶	- no overfill ³⁷

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Bleomycin 15 units (NB: dose in units only) (Hospira) (F)(PFL) no preservative ³⁸	6 mL * NS or SWI ³⁸	2.5 units/mL ³⁸	48 h F, 24 h RT ³⁸	50 mL * NS , SWI ³⁸	24 h RT ³⁹	- no overfill ⁴⁰
Bleomycin 15 units (NB: dose in units only) (Fresenius Kabi) (F)(PFL) no preservative ⁴¹	6 mL NS ⁴¹	2.5 units/mL ⁴¹	48 h F ⁴¹	50 mL NS ⁴¹	24 h RT ⁴¹	
Blinatumomab 38.5 mcg (Amgen) (F)(PFL) do not shake no preservative ⁴²	3 mL SWI ⁴² do NOT use supplied IV solution stabilizer to reconstitute vials ⁴² direct diluent against side of vial during reconstitution ⁴² gently swirl to avoid excess foaming ⁴²	12.5 mcg/mL ⁴²	4 h RT, 24 h F ⁴²	250 mL NS ⁴² add supplied IV solution stabilizer to NS bag and gently mix to avoid foaming ⁴² add reconstituted drug to bag following addition of IV solution stabilizer ⁴²	complete administration within 96 h RT, 10 d F ⁴²	- use non-DEHP bag and IV administration set ⁴² - use 0.2 or 0.22 micron low protein binding in-line filter ⁴² - prime lines with blinatumomab solution; do NOT use NS

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Bortezomib SC injection 3.5 mg (Actavis) (RT)(PFL) no preservative ⁴³	1.4 mL NS ⁴³	2.5 mg/mL ⁴³	8 h RT ⁴³	SC syringe ⁴³	8 h RT ⁴³	- auxiliary info: WARNING: SUBCUTANEOUS use only. Fatal if given by other routes.
Bortezomib 3.5 mg (Actavis) (RT)(PFL) no preservative ⁴³	3.5 mL NS ⁴³	1 mg/mL ⁴³	8 h RT ⁴³	IV syringe ⁴³	8 h RT ⁴³	- auxiliary info: WARNING: INTRAVENOUS use only. Fatal if given by other routes.
Bortezomib SC injection 3.5 mg (Janssen) (RT)(PFL) no preservative ⁴⁴	1.4 mL NS ⁴⁴	2.5 mg/mL ⁴⁴	2 d RT, F ^{18,45}	SC syringe ⁴⁴	48 h RT, 14 d F ^{18,45}	- auxiliary info: WARNING: SUBCUTANEOUS use only. Fatal if given by other routes.
Bortezomib 3.5 mg (Janssen) (RT)(PFL) no preservative ⁴⁴	3.5 mL NS ⁴⁴	1 mg/mL ⁴⁴	2 d RT, F ^{18,46}	IV syringe ⁴⁴	8 h RT ⁴⁴	- auxiliary info: WARNING: INTRAVENOUS use only. Fatal if given by other routes.

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Bortezomib SC injection 3.5 mg (Teva) (RT)(PFL) no preservative ⁴⁷	1.4 mL NS ⁴⁷	2.5 mg/mL ⁴⁷	2 d RT, F ^{18,48}	SC syringe ⁴⁷	48 h RT, 14 d F ^{18,48}	- auxiliary info: WARNING: SUBCUTANEOUS use only. Fatal if given by other routes.
Bortezomib 3.5 mg (Teva) (RT)(PFL) no preservative ⁴⁷	3.5 mL NS ⁴⁷	1 mg/mL ⁴⁷	2 d RT, F ⁴⁷	IV syringe ⁴⁷	48 h RT, 14 d F ^{18,48}	- auxiliary info: WARNING: INTRAVENOUS use only. Fatal if given by other routes.
Brentuximab vedotin 50 mg (GMD Distribution for Seattle Genetics) (F)(PFL) no preservative ⁴⁹	10.5 mL SWI ⁴⁹ Direct diluent against side of vial during reconstitution ⁴⁹ Do NOT shake ⁴⁹	5 mg/mL ⁴⁹	24 h F ⁴⁹	0.4-1.8 mg/mL in NS, D5W, Lactated Ringer's (i.e. 100-250 mL) ⁴⁹	24 h F ⁴⁹ Do NOT freeze ⁴⁹	- solution should be clear to slightly opalescent, colorless, and free of visible particulates ⁴⁹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p>Busulfan 60 mg/10 mL (Orphan Medical) (F) no preservative⁵⁰</p>	<p>N/A</p>	<p>use 5-micron nylon filter provided with ampoule to withdraw drug⁵⁰ 6 mg/mL⁵⁰</p>	<p>discard unused portion⁵⁰</p>	<p>NS or D5W (dilute in volume 10 times the busulfan volume to ~0.5 mg/mL)⁵⁰</p>	<p>complete administration within 12 h F: NS⁵⁰ 8 h RT: NS, D5W</p>	<p>- contains DMA***</p>
<p>Cabazitaxel 60 mg/1.5 mL (sanofi-aventis) (RT) no preservative⁵¹</p>	<p>supplied diluent: withdraw entire contents of diluent vial and inject into the concentrate vial⁵¹ slowly direct diluent against inside of vial to limit foaming⁵¹ mix by repeated inversions for 45 sec⁵¹ do NOT shake⁵¹ let sit for 5 min⁵¹</p>	<p>10 mg/mL⁵¹</p>	<p>1 h RT⁵¹</p>	<p>0.10 – 0.26 mg/mL NS, D5W⁵¹ (e.g., 250 mL*)</p>	<p>complete administration within 8 h RT, 48 h F⁵¹</p>	<p>- concentrate and diluent vials contain overflow⁵¹ - use non-DEHP bag and tubing⁵¹ - use 0.22 micron in- line filter⁵¹ - diluent contains 13% (w/w) ethanol in water⁵¹ - discard if crystallization occurs⁵¹</p>

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
CARBOplatin 50 mg/5 mL 150 mg/15 mL 450 mg/45 mL (Accord) (RT)(PFL) no preservative ⁵²	N/A	10 mg/mL ⁵²	discard unused portion ⁵²	0.5-10 mg/mL ⁵² NS, D5W⁵²	8 h RT, 24 h F ⁵²	- do NOT use aluminum-containing needle, syringe, or tubing ⁵²
CARBOplatin 50 mg/5 mL 150 mg/15 mL 450 mg/45 mL 600 mg/60 mL (Hospira) (RT)(PFL) no preservative ⁵³	N/A	10 mg/mL ⁵³	discard unused portion ⁵³	0.3-10 mg/mL ⁵⁴ NS, D5W^{11,53}	24 h RT, ⁵⁵ 48 h F ⁵³	- do NOT use aluminum-containing needle, syringe, or tubing ⁵⁴
CARBOplatin 50 mg/5 mL 150 mg/15 mL 450 mg/45 mL 600 mg/60 mL (Omega) (RT)(PFL) no preservative ⁵⁶	N/A	10 mg/mL ⁵⁶	discard unused portion ⁵⁶	0.3-10 mg/mL ⁵⁶ NS, D5W⁵⁶ do NOT use aluminum-containing needle or syringe ⁵⁶	24 h RT ⁵⁷ , 48 h F ⁵⁶	- do NOT use aluminum-containing needle, syringe or tubing ⁵⁶

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
CARBOplatin 50 mg/5 mL 150 mg/15 mL 450 mg/45 mL (Teva/Novopharm) (RT)(PFL) no preservative ⁵⁸	N/A	10 mg/mL ⁵⁸	discard unused portion RT ⁵⁸	0.5-10 mg/mL ⁵⁹ NS, D5W ^{11,58,60}	8 h RT ⁵⁸	- do NOT use aluminum-containing needle, syringe, or tubing ⁵⁸
Carfilzomib 10 mg 30 mg 60 mg (Amgen) (F)(PFL) no preservative ⁶¹	10 mg: 5 mL SWI ⁶¹ 30 mg: 15 mL SWI ⁶¹ 60 mg: 29 mL SWI ⁶¹ direct diluent against side of vial during reconstitution ⁶¹ swirl gently; do NOT shake ⁶¹ if foaming occurs, allow to settle until clear (about 5 minutes) ⁶¹ record time of reconstitution	2 mg/mL ⁶¹	24 h F, 4 h RT ⁶¹	50 -100 mL D5W <u>only</u> ⁶¹ do NOT dilute in NS ⁶¹	complete administration within 24 h F, 4 h RT after reconstitution ⁶¹	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Carmustine 100 mg (Bristol Labs) (F) no preservative ⁶²	3 mL diluent (supplied) ⁶² diluent to reach RT, then dissolve drug with 3 mL diluent; add 27 mL SWI ⁶² record time of reconstitution	3.3 mg/mL in 10% ethanol ⁶²	24 h F, 8 h RT ⁶²	glass ⁶² or polyolefin container ¹¹ 500 mL NS or D5W ⁶²	24 h F: in glass ⁶² or polyolefin container ¹¹ use within 4 h of reconstitution RT ⁶²	- do not use if product has oily droplets ⁶²
Cetuximab 100 mg/50 mL 200 mg/100 mL (ImClone/BMS) (F) do not dilute do not shake no preservative ⁶³	N/A	2 mg/mL ⁶³	discard unused portion after 12 h F, 8 h RT ⁶³	syringe ⁶³ sterile evacuated container or bag e.g. polyolefin, polyethylene, ethylene vinyl acetate, DEHP plasticized PVC, PVC bag, or glass ⁶³	12 h F, 8 h RT ⁶³ 12 h F, 8 h RT ⁶³	- administer with a 0.2 or 0.22 micron low protein binding in-line filter ⁶³ - normal saline may be used to flush the line ⁶³ - solution may contain white particulates which do not affect product quality ⁶³

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
CISplatin 10 mg/10 mL 50 mg/50 mL 100 mg/100mL (Hospira) (RT)(PFL) no preservative ⁶⁴	N/A	1 mg/mL ⁶⁴	48 h RT ⁶⁵	Less than or equal to 60 mg: 100 mL* NS Greater than 60 mg: 250 mL* NS 500 or 1000 mL* NS , D5-NS, D5-1/2S, D5- NS with mannitol, D5- 1/2S with mannitol ^{64,66} , D5W- 1/3S with mannitol ⁶⁴	48 h RT ⁶⁵	- do NOT use aluminum-containing needle, syringe or tubing ⁶⁴
CISplatin 10 mg/10 mL 50 mg/50 mL 100 mg/100mL (Sandoz) (RT)(PFL) no preservative ⁶⁷	N/A	1 mg/mL ⁶⁷	48 h RT ^{67,68}	Less than or equal to 60 mg: 100 mL NS* Greater than 60 mg: 250 mL NS* NS; 0.45% Sodium Chloride with or without mannitol ⁶⁹ 2 L of D5 in one-half or one-third NS containing 37.5 g of mannitol ⁶⁷	24 h RT ⁶⁷	- do NOT use aluminum-containing needle, syringe or tubing ⁶⁷

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
CISplatin 10 mg/10 mL 50 mg/50 mL 100 mg/100mL (Teva) (RT)(PFL) no preservative ⁷⁰	N/A	1 mg/mL ⁷⁰	discard unused portion ¹⁸	Less than or equal to 60 mg: 100 mL* NS Greater than 60 mg: 250 mL* NS 2 L of D5 in one-half or one-third NS containing 37.5 g of mannitol ⁷⁰	24 h RT ⁷⁰	- do NOT use aluminum-containing needle, syringe or tubing ⁷⁰

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Cladribine 10 mg/10 mL (Fresenius Kabi) (F)(PFL) no preservative ⁷¹	N/A	1 mg/mL ⁷¹	discard unused portion ⁷¹	SC syringe ⁷²	discard end of day ^{13,71,73}	
				500 mL NS only Do NOT use D5W	24 h RT	
				Cassette: qs to 100 mL with bacteriostatic NS only via SIMS DELTEC INC. MEDICATION CASSETTES® ⁷¹ filter drug and diluent through 0.22 micron filter as each solution is being introduced into the cassette	at least 7 days ⁷¹	
Clodronate 300 mg/10 mL (Oryx) (RT) no preservative ⁷⁴	N/A	30 mg/mL ⁷⁴	discard unused portion ⁷⁴	500 mL NS or D5W ⁷⁴	12 h RT ⁷⁴	- do NOT mix with calcium containing solution (e.g., Ringer's) ⁷⁵

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Cyclophosphamide 200 mg 500 mg 1000 mg 2000 mg (Baxter) (RT)(PFL) no preservative ⁷⁶	NS⁷⁷ 200 mg: 10 mL 500 mg: 25 mL 1000 mg: 50 mL 2000 mg: 100 mL ⁷⁶	20 mg/mL ⁷⁶	48 h F, ^{68,76,78} 24 h RT ⁷⁶	Less than or equal to 1 g: 100 mL NS* Greater than 1 g: 250 mL NS* high dose in BMT: may need 500 NS* NS, D5W, D5NS⁷⁶	72 h F, ^{76,78} 24 h RT ⁷⁶	
Cytarabine 100 mg/1 mL 1000 mg/10mL 2000 mg/20mL (Hospira) (RT)(PFL) no preservative ⁷⁹	N/A record time of puncture	100 mg/mL ⁷⁹	24 h RT ⁷⁹	100 mL* NS , Water for Injection, D5W, Lactated Ringer's ⁷⁹	72 h F , 24 h RT from initial vial puncture ⁷⁹	- do not use for IT injection
Cytarabine IT injection 100 mg/1 mL 1000 mg/10mL 2000 mg/20mL (Hospira) (RT)(PFL) no preservative ⁷⁹	N/A record time of puncture	100 mg/mL ⁷⁹	24 h RT ⁷⁹	diluents containing preservatives should NOT be used for intrathecal administration ⁷⁹ qs to 6 mL with preservative free NS ⁸⁰	use within 4 h of initial vial puncture ¹⁸	- auxiliary label: IT injection ¹⁸ - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag ¹⁸

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Cytarabine SC injection 100 mg/1 mL 1000 mg/10mL 2000 mg/20mL (Hospira) (RT)(PFL) no preservative ⁷⁹	N/A record time of puncture	100 mg/mL ⁷⁹	24 h RT ⁷⁹	syringe	14 d F, 48 h RT ^{18,81}	- do not use for IT injection
Dacarbazine 100 mg 200 mg (Abraxis) (F)(PFL) no preservative ⁸²	100 mg: 9.9 mL SWI ⁸² 200 mg: 19.7 mL SWI ⁸²	10 mg/mL ⁸²	72 h F, 8 h RT ⁸²	250-1000 mL* NS or D5W	24 h F, 8 h RT ⁸² **(PFL) ^{11,82} see Special Precautions/Notes Column	- protect container from light during storage and administration ⁸³ - overfill unknown
Dacarbazine 200 mg 600 mg (Hospira) (F)(PFL) no preservative ⁸⁴	200 mg: 19.7 mL SWI ⁸⁴ 600 mg: 59.1 mL SWI ⁸⁴	10 mg/mL ⁸⁴	48 h F, 8 h RT ⁸⁴ (PFL) ⁸⁵	0.19–3.0 mg/mL ^{13,84} 250-1000 mL* NS or D5W	24 h F ⁸⁴ **(PFL) ⁸³ see Special Precautions/Notes Column	- protect container from light during storage and administration ⁸³ - no overfill ^{40,85}

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
DACTINomycin 0.5 mg (GMD Pharma for Recordati) (RT)(PFL) no preservative ⁸⁶	1.1 mL SWI (preservative-free) ⁸⁶ Do NOT use SWI with preservative (may form precipitate) ⁸⁶	0.5 mg/mL (500 mcg/mL) ⁸⁶	discard unused portion ⁸⁸	syringe ⁸⁶	use within 4 h of initial vial puncture ⁶⁸	- drug loss reported with some cellulose ester membrane in- line filters ⁸⁶
				10 mcg/mL or greater ⁸⁶ NS, D5W ^{86,87}		
Daratumumab 100 mg/5mL 400 mg/20mL (Janssen) (F)(PFL) do not shake no preservative ⁸⁸	N/A	20 mg/mL ⁸⁸	discard unused portion ⁸⁸	500-1000 mL NS dilute to final volume by withdrawing volume from bag equal to volume of drug to be added ⁸⁸ mix by gentle inversion ⁸⁸	24 h F , followed by 15 h infusion (total 39 h) ⁸⁸ allow bag to come to room temperature, then use immediately ⁸⁸ **(PFL)	- administer with a 0.22 or 0.2 micron low protein binding in-line filter ⁸⁸ - discard if visible particles are observed ⁸⁸ - complete infusion within 15 hours ⁸⁸
DAUNOrubicin 20 mg (Erfa Canada Inc.) ⁸⁹ (RT)(PFL) ⁹⁰ no preservative ⁹¹	4 mL SWI ⁸⁹	5 mg/mL ^{89,92}	48 h F , 24 h RT ⁹¹	100-250 mL in isotonic solution e.g., NS ⁸⁹ no data for D5W ⁹¹	24 h RT , 48 h F ⁸⁹	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
DAUNOrubicin 20 mg (Teva/Novopharm) (RT)(PFL) no preservative ⁹³	4 mL SWI ⁹³	5 mg/mL ⁹³	24 h RT, 48 h F ⁹³ (PFL) ⁹³	100-250 mL NS or D5W ¹¹	48 h F, 24 h RT ⁹³ **(PFL) ⁹³	
Degarelix 80 mg 120 mg (Ferring) (RT) do not shake ⁹⁴ no preservative ⁹⁵	80 mg: 4.2 mL SWI (supplied diluent) ⁹⁴	20 mg/mL ⁹⁴	2 h RT ⁹⁴	SC syringe ⁹⁴	2 h RT ⁹⁴	
	120 mg: 3 mL SWI (supplied diluent) ⁹⁴	40 mg/mL ⁹⁴				
	swirl gently; avoid shaking to prevent foam formation ⁹⁴ reconstitution may take up to 15 min ⁹⁴					

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Denosumab (XGEVA) 120 mg/1.7 mL (Amgen) (F)(PFL) do not shake no preservative ⁹⁶	N/A	71 mg/mL ⁹⁶	discard unused portion ^{68,96}	SC syringe ⁹⁶	use within 4 h of initial puncture ⁶⁸	- not interchangeable with PROLIA ⁹⁶ - do not use if solution is cloudy; trace amounts of translucent to white proteinaceous particles are acceptable ⁹⁶ - avoid vigorous shaking ⁹⁶ - bring to room temperature 15-30 minutes prior to administration ⁹⁶
Dexrazoxane 250 mg 500 mg (Pfizer) (RT) no preservative ⁹⁷	SWI ⁹⁷ 250 mg: 25 mL 500 mg: 50 mL	10 mg/mL ⁹⁷	30 min RT, 3 h F ⁹⁸	MUST BE FURTHER DILUTED With Lactated Ringers Injection to 1.3 – 3.0 mg/mL ⁹⁷	1 h RT, 4 h F ⁹⁷	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p>DOCEtaxel 20 mg/2 mL 80 mg/8 mL 160 mg/16 mL (Hospira) (F, RT)(PFL) preservative⁹⁹</p>	<p align="center">N/A</p>	<p align="center">10 mg/mL⁹⁹</p>	<p>20mg/2 mL vial: discard unused portion^{18,99}</p>	<p align="center">0.3-0.74 mg/mL⁹⁹ (250 mL NS or D5W)⁹⁹</p>	<p align="center">complete administration within 4 h F,⁹⁹ 48 h RT^{18,100}</p>	<p>- use non-DEHP bag and IV administration set⁹⁹</p>
			<p>80 mg/8 mL or 160 mg/16 mL vial (maximum number of punctures: up to 3 doses can be removed when a venting needle is also inserted, i.e., 6 punctures total)¹⁰¹ 14 d F^{18,99} **(PFL)^{18,99}</p>			

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p>DOCEtaxel 20 mg/0.5 mL 80 mg/2 mL (sanofi-aventis) (F, RT)(PFL) no preservative¹⁰²</p>	<p>supplied diluent : - if vials were refrigerated, allow to warm for 5 min at RT. Withdraw entire contents of the diluent and inject the entire contents of the syringe into the corresponding concentrate vial. Mix by repeated inversions for 45 sec¹⁰²</p> <p>DO NOT SHAKE¹⁰²</p> <p>Let sit for 5 minutes¹⁰²</p>	<p>10 mg/mL¹⁰²</p>	<p>48 h F, RT^{18,102,103}</p>	<p>0.3-0.74 mg/mL¹⁰² (250 mL NS or D5W)¹⁰²</p>	<p>complete administration within 4 h F,¹⁰² 48 h RT^{18,103}</p>	<p>- use non-DEHP bag and IV administration set¹⁰²</p>
<p>DOXOrubicin 10 mg/5 mL 20 mg/10 mL 50 mg/25 mL 200 mg/100 mL (Accord) (F)(PFL) no preservative¹⁰⁴</p>	<p>N/A</p>	<p>2 mg/mL¹⁰⁴</p>	<p>8 h¹⁰⁴</p>	<p>syringe¹⁰⁴</p>	<p>24 h F, RT from initial vial puncture¹⁰⁴</p>	<p>- for ULYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing etoposide, DOXOrubicin, vinCRISStine)</p>

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
DOXOrubicin 10 mg 50 mg 150 mg (Hospira) (RT)(PFL) no preservative ¹⁰⁵	NS, SWI, D5W ¹⁰⁵ (NS reconstitution takes longer) 10 mg: 5 mL 50 mg: 25 mL 150 mg: 75 mL	2 mg/mL ¹⁰⁵	48 h F, 24 h RT ^{13,105}	syringe ¹⁰⁵	48 h F, 24 h RT ^{13,106}	- for ULYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing etoposide, DOXOrubicin, vinCRISStine)
DOXOrubicin 10 mg/5 mL 20 mg/10 mL 50 mg/25 mL 200 mg/100 mL (Teva/Novopharm) (F)(PFL) no preservative ¹⁰⁷	N/A record time of puncture	2 mg/mL ¹⁰⁷	8 h ¹⁰⁷	syringe ¹⁰⁷	48 h F, 24 h RT ¹⁰⁷ from initial vial puncture	- for ULYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing etoposide, DOXOrubicin, vinCRISStine)
DOXOrubicin 10 mg/5 mL 50 mg/25 mL 200 mg/100 mL (Pfizer) (F) no preservative ¹⁰⁸	N/A	2 mg/mL ¹⁰⁸	discard unused portion ^{68,108}	syringe ¹⁰⁸	48 h F, 24 h RT ¹⁰⁸	- for ULYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing etoposide, DOXOrubicin, vinCRISStine)

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
DOXOrubicin Pegylated Liposomal 20 mg/10 mL (Janssen) (F) no preservative ¹⁰⁹	N/A	2 mg/mL ¹⁰⁹	discard unused portion ¹⁰⁹	Less than 90 mg: 250 mL D5W only ¹⁰⁹ Greater than or equal to 90 mg: 500mL D5W only ¹⁰⁹	24 h F ¹⁰⁹	- do not filter ¹⁰⁹
Epirubicin 10 mg/5 mL 20 mg/10 mL 50 mg/25 mL 150 mg/75 mL 200 mg/100 mL (Teva/Novopharm) (F)(PFL) no preservative ¹¹⁰	N/A	2 mg/mL ¹¹⁰	8 h F, RT ¹¹⁰	syringe ¹¹⁰	48 h F, 24 h RT from initial vial puncture ¹¹⁰	
Epirubicin 10 mg/5 mL 50 mg/25 mL 200 mg/100 mL (Fresenius Kabi) (F)(PFL) no preservative ¹¹¹	N/A record time of puncture	2 mg/mL ¹¹¹	8 h ¹¹¹	syringe ¹¹¹	48 h F, 24 h RT from initial vial puncture ¹¹¹	
				100 mL* NS or D5W	2 d F, RT: NS or D5W ^{18,111}	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Epirubicin 10 mg/5 mL 50 mg/25 mL 200 mg/100 mL (Pfizer) (F)(PFL) no preservative ¹¹²	N/A record time of puncture	2 mg/mL ¹¹²	8 h ¹¹²	syringe ¹¹²	48 h F , 24 h RT from initial vial puncture ¹¹²	
				100 mL* NS or D5W ¹¹	2 d F , RT: NS or D5W ⁶⁵	
EPOCHR (ULYEPOCHR protocol) (RT)(PFL) no preservative ^{18,113-116}	see brand specific entries for: DOXOrubicin as applicable	see brand specific entries for: DOXOrubicin, etoposide, vinCRISTine	see brand specific entries for: DOXOrubicin, etoposide, vinCRISTine	etoposide dose ≤125 mg/24 h: in 500 mL NS etoposide dose >125 mg/24 h: in 1000 mL NS	etoposide concentration ≤0.25 mg/mL: complete administration within 72 h RT precipitation occurs at etoposide concentrations >0.25 mg/mL **(PFL) see Special Precautions/Notes column	- final product is a 3-in-1 solution containing etoposide, DOXOrubicin, vinCRISTine (refer to ULYEPOCHR protocol) - use non-DEHP bag and tubing only - use inline filter - protect container from light during administration and storage

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
eriBULin 1 mg/2 mL (Eisai Limited) (RT)(PFL) ¹¹⁷ no preservative ¹⁸	N/A	0.5 mg/mL ¹¹⁷	discard unused portion ^{18,117}	IV syringe ¹¹⁷	24 h F, 6 h RT ¹¹⁷	- do not administer through dextrose containing lines ¹¹⁷ - vials contain dehydrated alcohol USP (5% v/v) ¹¹⁷
Etoposide 100 mg/5 mL 200 mg/10 mL 500 mg/25 mL 1000 mg/50 mL (Sandoz) (RT)(PFL) preservative ¹¹⁸	N/A	20 mg/mL ¹¹⁸	14 d RT ¹¹⁸	0.2-0.4 mg/mL NS, D5W ¹¹⁸ 500 mL * NS or D5W	0.2 mg/mL: 7 d F, RT ¹¹⁸ 0.4 mg/mL: 12 h F, RT ¹¹⁸	- use non-DEHP bag and tubing only - use 0.22 micron in- line filter ¹¹⁹ - for ULYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing etoposide, DOXOrubicin, vinCRISine)

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Etoposide 100 mg/5 mL 200 mg/10 mL 500 mg/25 mL 1000 mg/50 mL (Teva/Novopharm) (RT)(PFL) no preservative ¹²⁰	N/A	20 mg/mL ¹²⁰	discard unused portion ¹²⁰	NS Stability is concentration dependent	0.2-0.3 mg/mL: 7 d F, ¹²¹ 2 d RT ^{121,122} 0.4-0.5 mg/mL: 1 d F, ¹²¹ 1d RT ¹²¹ 0.6-9.0mg/mL: generally unstable 9.5 mg/mL: 2 d F, ¹²¹ 1d RT ¹²¹ 10-12 mg/mL: 7 d F, ¹²¹ 2 d RT ^{121,122}	- use non-DEHP bag and tubing only - use 0.22 micron in-line filter ¹¹⁹ - for ULYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing etoposide, DOXOrubicin, vinCRISTine)
				D5W ¹²⁰	4 h RT ^{120,123}	
Etoposide phosphate (ETOPOPHOS®) 100 mg (BMS) (F)(PFL) no preservative ¹²⁴	5 mL NS, D5W, SWI, BWI ^{124,125}	20 mg/mL ^{124,125}	24 h RT ^{124,125} , 48 h F ^{18,124,125}	500 mL* NS, D5W ^{124,125} (do not dilute to less than 0.1 mg/mL) ^{124,125}	24 h F, RT ^{124,125}	
	10 mL NS, D5W, SWI, BWI ^{124,125}	10 mg/mL ^{124,125}				

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Filgrastim (NEUPOGEN®) 300 mcg/1 mL 480 mcg/1.6 mL (Amgen) (F)(PFL) do not shake no preservative ¹²⁶	N/A	300 mcg/mL ¹²⁶	discard unused portion ¹⁸	SC syringe ¹²⁶	14 d F ^{18,127}	- albumin is added to D5W to prevent filgrastim adsorption to plastic ¹²⁶ - incompatible with saline ^{126,128} - do NOT dilute to less than 5 mcg/mL ¹²⁶
				50-100 mL D5W only ¹²⁸ in PVC, polyolefin, or glass ¹²⁶ (for filgrastim concentrations of 5- 15 mcg/mL in D5W, add albumin 2 mg/mL) ¹²⁶	48 h RT, 7 d F ^{18,127}	
Fludarabine 50 mg (Berlex) (F) no preservative ¹²⁹	2 mL SWI ¹²⁹	25 mg/mL ¹²⁹	48 h F or RT ^{13,65}	dilute to maximum of 1 mg/mL ^{129,130} 50-100 mL* NS or D5W ¹²⁹	48 h F, RT ^{13,65}	
Fludarabine 50 mg (Teva/Novopharm) (F) no preservative ¹³¹	N/A	25 mg/mL ¹³¹	discard unused portion ¹³¹	dilute to maximum of 1 mg/mL ¹³¹ 50-100 mL* NS or D5W	48 h F, 24 h RT ¹³¹	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Fluorouracil 5000 mg/100 mL (Accord) (RT)(PFL) no preservative ¹³²	N/A	50 mg/mL ¹³²	48 h RT ^{18,133}	syringe ¹³²	48 h RT ^{18,133}	
				0.5-10 mg/mL ¹³³ 50-1000 mL* D5W	48 h RT ^{18,133}	
				CIVI: ambulatory pump ¹³⁴	complete within 8 d ¹³³	
Fluorouracil 5000 mg/100 mL (Hospira) (RT)(PFL) no preservative ¹³⁵	N/A	50 mg/mL ¹³⁵	8 h RT ^{134,135}	syringe ¹³	48 h RT ^{13,35,134}	
				2-10 mg/mL ^{134,135} 50-1000 mL* D5W	24 h RT ^{134,135}	
				CIVI: ambulatory pump ¹³⁴	complete within 8 d ^{11,13,136,137}	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Fluorouracil 500 mg/10 mL 5000 mg/100 mL (Sandoz) (RT)(PFL) no preservative ¹³⁸	N/A	50 mg/mL ¹³⁸	4 h RT ¹⁸	syringe	4 h RT ¹⁸	
				300-500 mL* D5W ¹³⁸	24 h RT ¹³⁸	
				CIVI: ambulatory pump ¹³⁴	complete within 8 d ^{11,13,136,137}	
Gemcitabine 200 mg 1000 mg 2000 mg (Accord) (RT) no preservative ¹³⁹	200 mg: 5 mL NS 1000 mg: 25 mL NS 2000 mg: 50 mL NS ¹³⁹	38 mg/mL ¹³⁹	24 h RT ¹³⁹	0.1-10 mg/mL NS ¹³⁹	48 h RT ^{18,140,141}	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Gemcitabine 200 mg 1000 mg (Eli-Lilly) (RT) no preservative ¹⁴²	200 mg: 5 mL NS 1000 mg: 25 mL NS ¹⁴²	38 mg/mL ¹⁴²	48 h RT ^{142,143}	syringe ¹⁴²	48 h RT ^{13,142,143}	
				0.1–10 mg/mL NS ^{142,143}	48 h F, RT ^{13,142,143}	
Gemcitabine 200 mg 1000 mg 2000 mg (Hospira) (RT) ¹⁴⁴ no preservative ¹⁴⁵	200 mg: 5 mL NS 1000 mg: 25 mL NS 2000 mg: 50 mL NS ¹⁴⁴	38 mg/mL ¹⁴⁴	48 RT ^{68,144,146}	syringe ¹⁴⁴	24 h RT ^{144,146}	
				0.1 - 26 mg/mL NS ^{144,146}	48 h RT ^{68,146}	
Gemcitabine 200 mg/5.3 mL 1000 mg/26.3 mL 2000 mg/52.6 mL (Hospira) (F) no preservative ¹⁴⁷	N/A	38 mg/mL ¹⁴⁸	discard unused portion ¹⁸	0.1 – 38 mg/mL NS , D5W ¹⁴⁸	24 h RT ¹⁴⁸	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Gemcitabine 200 mg 1000 mg (Teva/Novopharm) (RT) no preservative ¹⁴⁹	200 mg: 5mL NS 1000 mg: 25 mL NS ¹⁴⁹	38 mg/mL ¹⁴⁹	24 h RT ¹⁴⁹	0.1 - 38 mg/mL NS ¹⁴⁹	24 RT ¹⁴⁹	
Gemcitabine 200 mg 1000 mg (Sandoz Standard) (RT) no preservative ¹⁵⁰	200 mg: 5 mL NS 1000 mg: 25 mL NS ¹⁵⁰	38 mg/mL ¹⁵⁰	48 h RT ^{150,151}	syringe ¹⁵⁰	48 h RT ¹⁵⁰⁻¹⁵²	
				0.1 - 38 mg/mL NS or D5W ^{150,153}	48 h RT ^{13,154}	
IDArubicin 5 mg 10mg (Pfizer) (RT)(PFL) no preservative ¹⁵⁵	5 mg: 5 mL SWI ¹⁵⁵ 10 mg: 10 mL SWI ¹⁵⁵ vial contents under negative pressure ¹⁵⁵ do NOT use BWI to reconstitute ¹⁵⁵	1 mg/mL ¹⁵⁵	48 h F , 24 h RT ¹⁵⁵ **(PFL) ¹⁵⁵	syringe ¹⁵⁵	48 h F , 24 h RT ¹⁵⁵	- avoid alkaline solutions ¹⁵⁵

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
IDArubicin PFS 5 mg/5 mL 10 mg/10 mL 20 mg/20 mL (Pfizer) (F)(PFL) no preservative ¹⁵⁵	N/A	1 mg/mL ¹⁵⁵	24 h RT, 48 h F **(PFL) ¹⁵⁵	syringe ¹⁵⁵	4 h from initial puncture ¹⁸	- avoid alkaline solutions ¹⁵⁵
IDArubicin 5 mg/5 mL 10 mg/10 mL 20 mg/20 mL (Fresenius Kabi) (F)(PFL) no preservative ¹⁵⁶	N/A	1 mg/mL ¹⁵⁶	discard unused solution ¹⁵⁶	syringe ¹⁵⁶	4 h from initial puncture ¹⁸	- avoid alkaline solutions ¹⁵⁶
Ifosfamide 1000 mg 3000 mg (Baxter) (RT) no preservative ¹⁵⁷	1000 mg: 20 mL SWI ¹⁵⁷ 3000 mg: 60 mL SWI ¹⁵⁷ shake well	50 mg/mL ¹⁵⁷	24 h RT, 48 h F ^{18,157}	0.6–20 mg/mL ¹⁵⁷ 500–1000 mL* NS, D5W, Lactated Ringer's ¹⁵⁷	24 h RT, 72 h F ¹⁵⁷ 24 h F, RT when mixed with mesna ¹¹	
Ifosfamide 1000 mg 3000 mg (Fresenius Kabi) (RT) no preservative ¹⁵⁸	1000 mg: 20 mL SWI 3000 mg: 60 mL SWI ¹⁵⁸ shake well	50 mg/mL ¹⁵⁸	24 h RT, 48 h F ^{18,158}	0.6-20 mg/mL ¹⁵⁸ 500-1000 mL* NS D5W, Lactated Ringer's ¹⁵⁸	24 h RT, 72 h F ¹⁵⁸ 24 h F, RT when mixed with mesna ¹¹	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Iniparib 100 mg/10 mL (sanofi-aventis) (F) no preservative ¹⁵⁹	N/A	10 mg/mL ¹⁵⁹	discard unused portion ¹⁵⁹	250 mL NS , D5W dilute to 250 mL final volume by withdrawing volume from bag equal to volume of drug to be added* ¹⁵⁹	24 h RT ¹⁵⁹	- *may also use empty IV bag and qs to final volume of 250 mL with NS , D5W ¹⁵⁹
Interferon Alfa -2b 10 million units/1 mL (Merck) (F) preservative ^{160,161}	N/A	10 million units/mL ¹⁶⁰	7 d F ¹⁶⁰	syringe ¹⁶⁰	7 d F ¹⁸	- vials can be kept at RT for up to 7 days before use; discard if not used within this time ¹⁶⁰
				final concentration ≥ 0.3 million IU/mL ¹⁶⁰ 50 mL NS ¹⁶⁰	24 h F , RT ¹⁶⁰	
Interferon Alfa -2b 18 million units/3 mL (Merck) (F) preservative ^{160,161}	N/A	6 million units/mL ¹⁶⁰	14 d F ^{18,160}	syringe ¹⁶⁰	14 d F ^{18,161}	- vials can be kept at RT for up to 7 days before use; discard if not used within this time ¹⁶⁰
				final concentration ≥ 0.3 million IU/mL ¹⁶⁰ 50 mL NS ¹⁶⁰	24 h F , RT ¹⁶⁰	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Interferon Alfa -2b 25 million units/2.5 mL (Merck) (F) preservative ^{160,161}	N/A	10 million units/mL ¹⁶⁰	14 d F ^{18,160}	syringe ¹⁶⁰	14 d F ^{18,161}	- vials can be kept at RT for up to 7 days before use; discard if not used within this time ¹⁶⁰
				final concentration ≥ 0.3 million IU/mL ¹⁶⁰ 50 mL NS ¹⁶⁰	24 h F, RT ¹⁶⁰	
Interferon Alfa -2b 10 million units (Merck) (F) no preservative (unless reconstituted with BWI) ¹⁶⁰	1 mL supplied diluent (SWI) ¹⁶⁰ do not shake; roll to reconstitute ¹⁶⁰	10 million units/mL ¹⁶⁰	24 h F ¹⁶⁰	syringe ¹⁶⁰	24 h F ^{18,161}	- after reconstitution, provides an isotonic solution which may be used for intralesional injection ¹⁶⁰ - non-reconstituted vials can be kept at RT for up to 4 weeks before use; discard if not reconstituted for use within this time ¹⁶⁰
				final concentration ≥ 0.1 million IU/mL ¹⁶⁰ 100 mL NS ¹⁶⁰	24 h F, RT ¹⁶¹	
	1 mL BWI ¹⁶⁰ do not shake; roll to reconstitute ¹⁶⁰		14 d F ^{18,160}	syringe ¹⁶⁰	14 d F ^{18,160}	
				final concentration ≥ 0.1 million IU/mL ¹⁶⁰ 100 mL NS ¹⁶⁰	24 h F, RT ¹⁶¹	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Interferon Alfa -2b 18 million units (Merck) (F) no preservative (unless reconstituted with BWI) ¹⁶⁰	1 mL supplied diluent ¹⁶⁰	18 million units/mL ¹⁶⁰	24 h F ¹⁶⁰	syringe ¹⁶⁰	24 h F ^{18,161}	- non-reconstituted vials can be kept at RT for up to 4 weeks before use; discard if not reconstituted for use within this time ¹⁶⁰
	do not shake; roll to reconstitute ¹⁶⁰			final concentration ≥ 0.1 million IU/mL ¹⁶⁰ 100 mL NS ¹⁶⁰	24 h F, RT ¹⁶¹	
	1 mL BWI ¹⁶⁰		14 d F ^{18,160}	syringe ¹⁶⁰	14 d F ^{18,160}	
	do not shake; roll to reconstitute ¹⁶⁰			final concentration ≥ 0.1 million IU/mL ¹⁶⁰ 100 mL NS ¹⁶⁰	24 h F, RT ¹⁶¹	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Ipilimumab 50 mg/10 mL 200 mg/40 mL (BMS Canada) (F)(PFL) no preservative ¹⁶²	N/A	5 mg/mL ¹⁶²	24 h F,RT ¹⁶²	1 – 4 mg/mL in NS, D5W 100 mL ¹⁶² OR undiluted in empty viaflex bag or glass bottle (allow vials to stand at RT for ~5 min prior to withdrawal of contents) ¹⁶²	24 h F,RT ¹⁶²	- do NOT shake ¹⁶² - administer with 0.2 or 0.22 low protein binding in-line filter ¹⁶² - vials may contain translucent-to-white amorphous particles ¹⁶² - discard if cloudy or has pronounced colour change (should be clear to pale yellow) ¹⁶² - flush line with NS or D5W after infusion ¹⁶²
Irinotecan 40 mg/2 mL 100 mg/5 mL 500 mg/25 mL (Accord) (RT)(PFL) no preservative ¹⁶³	N/A	20 mg/mL ¹⁶³	discard unused portion ¹⁶³	0.12 – 2.8 mg/mL ¹⁶³ 500 mL* D5W (preferred), NS ¹⁶³	48 h F, 24 h RT **(PFL) ¹⁶³	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Irinotecan 40 mg/2 mL 100 mg/5 mL 500 mg/25 mL (Hospira) (RT)(PFL) no preservative ¹⁶⁴	N/A	20 mg/mL ¹⁶⁴	2 days RT ^{13,165,166}	0.12– 2.8 mg/mL ¹⁶⁴ 500 mL ¹¹ D5W (preferred), NS ¹⁶⁴	24 h RT: D5W, NS ¹⁶⁴ 48 h F: D5W **(PFL) ¹⁶⁴	- do NOT refrigerate if in NS ¹⁶⁷
Irinotecan 40 mg/2 mL 100 mg/5 mL (Pfizer) (RT)(PFL) no preservative ¹⁶⁷	N/A	20 mg/mL ¹⁶⁷	discard unused portion ¹⁶⁷	0.12– 2.8 mg/mL ¹⁶⁷ 500 mL ¹¹ D5W (preferred), NS ¹⁶⁷	24 h RT: D5W, NS ¹⁶⁷ 48 h F: D5W **(PFL) ¹⁶⁷	- do NOT refrigerate if in NS ¹⁶⁷
Irinotecan 40 mg/2 mL 100 mg/5 mL 500 mg/25 mL (Sandoz) (RT)(PFL) no preservative ¹⁶⁸	N/A	20 mg/mL ¹⁶⁸	discard unused portion ^{68,168}	0.12-2.8 mg/mL ¹⁶⁸ D5W (recommended), NS ¹⁶⁸	24 h RT: D5W, NS ¹⁶⁸ 48 h F: D5W ¹⁶⁸ **(PFL) ¹⁶⁸	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Irinotecan Liposome SAP supply 50 mg/10 mL (Baxalta/Baxter) (F)(PFL) no preservative ¹	N/A	5 mg/mL ¹⁶⁹	discard unused portion ¹⁶⁹	dilute to a final volume of 500 mL with NS , D5W ¹⁶⁹	6 h RT, 24 h F ¹⁶⁹ **(PFL) (allow product to come to RT prior to administration if stored in F) ¹⁷⁰	- do not use in-line filter ¹⁷⁰
Irinotecan Liposome commercial supply 43 mg/10 mL (Baxalta) (F)(PFL) no preservative ¹⁷¹	N/A	4.3 mg/mL ¹⁷¹	discard unused portion ¹⁷¹	to a final volume of 500 mL with NS , D5W ¹⁷¹	4 h RT, 24 h F ¹⁷¹ **(PFL) (allow product to come to RT prior to administration if stored in F) ¹⁷¹	- do not use in-line filter ¹⁷¹ - expressed as irinotecan free base
Ixabepilone 15 mg (contains 16 mg) 45 mg (contains 47 mg) (BMS) (F)(PFL) no preservative ¹⁷²	15 mg: 8 mL supplied diluent ¹⁷² 45 mg: 23.5 mL supplied diluent ¹⁷²	2 mg/mL ¹⁷²	1 h RT ¹⁷²	0.2 – 0.6 mg/mL in Lactated Ringer's Injection USP (use non-DEHP infusion container) ¹⁷²	6 h RT ¹⁷²	- use 0.2-1.2 micron in-line filter ¹⁷² - use non-DEHP bag and administration set ¹⁷²

* Suggested volume based on usual dose range and any concentration range of stability data

** Protect from light means minimizing exposure to direct sunlight over a *storage* period. More specific information on protection from light (eg, protecting container and tubing during *administration*) will be indicated in the Special Precautions/Notes column.

*** Contains DMA (N,N dimethylacetamide). Product may be incompatible with closed system transfer devices such as ChemoLock.

Centres are not to change the content locally but should forward suggestions to the Cancer Drug Manual staff.

Explanatory Notes

Stability data assumes products prepared using standard aseptic technique in biological safety cabinet at low risk for contamination according to the classification outlined in USP 797.^{35,173}

Vial stability: Stability of solution after first puncture or reconstituted solution.

Storage temperature: If information states same stability with refrigerator and room temperature storage, then fridge stability is bolded as preferred (ie, to minimize growth of micro-organisms).

Discard unused portion: Unused portion from single use vials should be discarded at the end of the day.

“overfill known” is stated if the manufacturer states overfill that is present is within acceptable limits.

“Complete administration within ___” is stated if the manufacturer specifies that the infusion must be completed in a specific time frame following preparation, usually including entire time required for preparation (from first puncture), storage, and administration of infusion.

Abbreviations

BWI = bacteriostatic water for injection

CIVI: ambulatory pump = Continuous Intravenous Infusion (e.g., elastomeric infusor)

D5W = dextrose 5% in water

DMA = N,N dimethylacetamide

F = refrigerate

Non-DEHP = not containing Di(2-ethylhexyl) phthalate (DEHP)

NS = normal saline

PFL = protect from light

RT = room temperature

SWI = sterile water for injection

References

1. Novartis Pharmaceuticals Canada Inc. PROLEUKIN® product monograph. Dorval, Quebec; 6 July 2006.
2. McEvoy GK, editor. AHFS 2008 Drug Information. Bethesda, Maryland: American Society of Health-System Pharmacists, Inc. p. 917-925.
3. Koreth J, Matsuoka K, Kim HT, et al. Interleukin-2 and regulatory T cells in graft-versus-host disease. *N Engl J Med* 2011;365(22):2055-2066.
4. Koreth J, Alyea EP, Cutler C, Ho VT, et al. Clinical Study Protocol: A phase I study of ultra-low dose subcutaneous interleukin-2 (IL-2) for treatment of refractory chronic graft versus host disease. Boston, MA, USA: Dana Farber Cancer Institute; Harvard Medical Centre; 14 Dec 2010.
5. Rui Paiva. Personal communication. Business Unit Director, Transplant and Oncology; 1 June 2009.
6. Bayer HealthCare Pharmaceuticals. MabCampath® Package Insert. Toronto, Ontario; 1 September 2007.
7. Lundin J, Porwit-MacDonald A, Rossmann ED, et al. Cellular immune reconstitution after subcutaneous alemtuzumab (anti-CD52 monoclonal antibody, CAMPATH-1H) treatment as first-line therapy for B-cell chronic lymphocytic leukaemia. *Leukemia* 2004(18):484-490.
8. Berlex Canada Inc. Campath Drug Information. San Antonio, Texas; undated.
9. Schering AG. MABCAMPATH® Package Leaflet. Berlin, Germany; April 2002.
10. MedImmune Pharma B.V. Etyol Package Insert. The Netherlands; 2003.
11. Trissel LA. Handbook on Injectable Drugs. 13th ed. Bethesda, MD: American Society of Health-System Pharmacists, Inc.; 2005.
12. Erfa Canada Inc. AMSA PD® injection product monograph. Westmount, Quebec; 16 August 2005.

13. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 6 January 2006.
14. Tanya Leduc. Personal communication. Acting editor, BC Cancer Agency Cancer Drug Manual; 2 June 2008.
15. Lundbeck Canada Inc. TRISENOX® product monograph. Montreal, Quebec; 6 June 2013.
16. CGF Pharmatec for EUSA Pharma. KIDROLASE® product monograph. Montreal, Quebec; 17 April 2008.
17. Lexi-Drugs® (database on the Internet). Asparaginase (E. coli). Lexi-Comp Inc., 1 July 2015. Available at: <http://online.lexi.com>. Accessed 25 August 2015.
18. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 19 September 2007.
19. CGF Pharmatec for EUSA Pharma. ERWINASE® for Injection product monograph. Montreal, Quebec; 19 February 2015.
20. Health Canada. MedEffect® e-Notice - Important Safety Information on Shortage of Erwinase for Injection. 12 July 2017. Available at: <http://www.hc-sc.gc.ca/ahc-asc/media/advisories-avis/>. Accessed 12 July 2017.
21. Hoffman-La Roche Limited. TECENTRIQ® product monograph. 7070 Mississauga Road, Ontario; 12 April 2017.
22. Genentech Inc. TECENTRIQ® full prescribing information. South San Francisco, CA, USA; October 2016.
23. Merck KGaA. MSB0010718C Product Label (Compassionate Use Only). Darmstadt, Germany; Undated.
24. Merck KGaA. Pharmacy Manual Protocol Early Access - Single patient use of avelumab (MSB0010718C). 10 April 2017 - version 2.0.
25. Celgene Inc. VIDAZA® product monograph. Mississauga, Ontario; 17 May 2012.
26. Kevin Mejo. Personal communication. Medical Information, Celgene Inc.; 21 June 2017.
27. Tutino A, Lai M. Cold water reconstitution of Vidaza with subsequent refrigerated storage prolongs drug stability. *Eur J Oncol Pharm* 2011;5(3-4):24-25, 34.
28. Celgene Corporation. VIDAZA® full prescribing information. Summit, NJ, USA; December 2012.
29. Dr. Reddy's Laboratories Limited. Azacitidine for injection product monograph. Mississauga, Ontario; 19 April 2017.
30. Repchinsky C editor. ImmuCyst monograph, Compendium of Pharmaceuticals and Specialties. Ottawa, Ontario; 2005.
31. Organon Canada Ltd. OncoTICE® Product Monograph. Scarborough, Ontario; 24 October 2001.
32. Lundbeck Canada Inc. TREANDA® product monograph. Montreal, Quebec; 22 August 2012.
33. Roche. Bevacizumab for injection, Summary product information. Mississauga, Ontario; 2005.
34. Hoffmann-La Roche Limited. Ro 4876647 Avastin Investigator's Brochure. Mississauga, Ontario; 14th edition, November 2006.
35. The United States Pharmacopeia (USP). General Chapter 797: Pharmaceutical compounding - sterile preparations. USP 27-NF 22. Rockville, Maryland: The United States Pharmacopeial Convention, Inc.; 2004.
36. Repchinsky C editor. Blenoxane monograph, Compendium of Pharmaceuticals and Specialties. Ottawa, Ontario; 2005.
37. Catherine Dehaut M.Sc. Personal communication. Medical Information Associate, Bristol Laboratories of Canada; February 2008.
38. Faulding (Canada) Inc. Bleomycin Product Monograph. Montreal, Quebec; 3 June 2002.
39. Sheila Ahmed. Personal communication Bleomycin. Regulatory Affairs Associate, Bleomycin, Mayne Pharma Canada. Ref JUN-128-2004; July 2, 2004.
40. Robert Caunce. Personal communication. Quality System Manager, Hospira Australia; 12 March 2008.
41. Pharmaceutical Partners of Canada Inc. Bleomycin For Injection product monograph. Richmond Hill, Ontario; 14 January 2008.
42. Amgen Canada Inc. BLINCYTO® product monograph. Mississauga, Ontario; 12 July 2016.
43. Actavis Pharma Company. ACT BORTEZOMIB® Bortezomib for injection product monograph. Mississauga, Ontario; 24 September 2015.
44. Janssen Inc. VELCADE® product monograph. Toronto, Ontario; 20 June 2013.
45. Walker SE, Charbonneau LF, Law S. Stability of bortezomib 2.5 mg/mL in vials and syringes stored at 4°C and room temperature (23°C). *Can J Hosp Pharm* 2014;67(2):102-107.
46. Walker SE, Milliken D, Law S. Stability of bortezomib reconstituted with 0.9% sodium chloride at 4°C and room temperature (23°C). *Canadian Journal of Hospital Pharmacy* 2008;61(1):14-20.
47. Teva Canada Limited. Bortezomib for injection® product monograph. Toronto, Ontario; 22 January 2015.
48. Walker SE, Charbonneau LF, Tyono I, et al. Stability of 1.0 and 2.5 mg/mL bortezomib in vials and syringes following reconstitution with sodium chloride at 4 °C and 23 °C. Poster Presentation Canadian Society of Hospital Pharmacists Professional Practice Conference 2016. Jan 30 - Feb 3, 2016.
49. GMD Distribution Inc. for Seattle Genetics Inc. ADCETRIS® product monograph. Oakville, Ontario; 1 February 2013.
50. Orphan Medical Inc. Busulfex Package Insert. Flamborough, Ontario; August 1999.
51. sanofi-aventis Canada Inc. JEV TANA® product monograph. Laval, Quebec; 13 September 2013.
52. Accord Healthcare Inc. Carboplatin injection® product monograph. Markham, Ontario; 15 December 2015.
53. Mayne Pharma (Canada) Inc. Carboplatin Package Insert. Montreal, QC; Undated.

54. Nancy Grigoriadis. Personal communication. Regulatory Affairs Associate, Mayne Pharma (Canada) Inc. 2 October 2006.
55. Nancy Grigoriadis. Personal communication. Regulatory Affairs Associate, Mayne Pharma (Canada) Inc. 2 October 2006.
56. Omega Laboratories Ltd. Carboplatin injection product monograph. Montreal, Quebec; 24 March 2011.
57. Nayla El Zir. Personal communication. Associate, Regulatory Affairs, Omega Laboratories Limited; 12 April 2017.
58. Novopharm Limited. Carboplatin Package Insert. Toronto, Canada; Undated.
59. Manjinder S Kang. Personal communication. Regulatory Affairs Drug Information Pharmacist, Novopharm Canada. 14 March 2005.
60. Repchinsky C editor. Paraplatin-AQ, Compendium of Pharmaceuticals and Specialties. 12th ed. Ottawa, Ontario: Canadian Pharmacists Association; 2004.
61. Amgen Canada Inc. KYPROLIS® product monograph. Mississauga, Ontario; 6 July 2017.
62. Bristol Laboratories of Canada. BiCNU Package Insert. Montreal, Canada; 2002.
63. ImClone Systems Incorporated and Bristol-Myers Squibb Company. ERBITUX® product monograph. Branchburg, NJ, USA; March 2009.
64. Mayne Pharma (Canada) Inc. Cisplatin Package Insert. Montreal, QC; Undated.
65. Trissel LA. Handbook on Injectable Drugs. 12th ed. Bethesda, MD: American Society of Health-System Pharmacists, Inc.; 2003.
66. Repchinsky C editor. Cisplatin CPhA monograph, Compendium of Pharmaceuticals and Specialties. 12th ed. Ottawa, Ontario: Canadian Pharmacists Association; 2004.
67. Sandoz Canada Inc. Cisplatin Injection BP product monograph. Boucherville, Quebec; 13 April 2011.
68. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 19 September 2007.
69. Trissel LA. Handbook on Injectable Drugs. 16th ed. Bethesda, Maryland: American Society of Health-System Pharmacists, Inc; 2011. p. 378.
70. Teva Canada Limited. Cisplatin injection® product monograph. Toronto, Ontario; 6 March 2013.
71. Pharmaceutical Partners of Canada, Inc. Cladribine For Injection product monograph. Richmond Hill, Ontario; 27 November 2008.
72. BC Cancer Agency Lymphoma Tumour Group. (LYCDA) BCCA Protocol Summary for Treatment of Hairy Cell Leukemia with Cladribine. Vancouver, British Columbia: BC Cancer Agency; 1 February 2007.
73. de Lemos ML, Hamata L. Stability issues of parenteral chemotherapy drugs. J Oncol Pharm Pract 2007;13(1):27-31.
74. Oryx Pharmaceuticals Inc. Clasteon Product Monograph. Mississauga, Ontario; 25 November 2004.
75. Bayer Inc. BONEFOS® product monograph. Mississauga, Ontario; 30 March 2017.
76. Baxter Corporation. Procytox Package Insert. Toronto, Ontario; 2004.
77. Baxter Corporation. Procytox Package Insert. Mississauga, Ontario; 1 October 2003.
78. Paul Agro. Personal communication. Medical Information, cyclophosphamide, Baxter. 12 July, 2006.
79. Hospira Healthcare Corporation. Cytarabine Injection® product monograph. Saint-Laurent, Quebec; 25 November 2013.
80. BC Cancer Agency Miscellaneous Origin Tumour Group. (MOIT) BCCA Protocol Summary for Solid Tumours using Intrathecal Methotrexate and/or Thiotepa and/or Cytarabine. Vancouver, British Columbia: BC Cancer Agency; 1 May 2009.
81. My Dang, Regulatory Affairs Associate. Personal communication. Mayne Pharma (Canada) Inc.; 4 May 2007.
82. Abraxis Pharmaceutical Products. Dacarbazine product information package. Schaumburh, IL; December 2006.
83. Trissel L. Handbook on injectable drugs. 13th ed. Bethesda, Maryland: American Society of Health-System Pharmacists; 2005. p. 428-431.
84. Mayne Pharma (Canada) Inc. DACARBAZINE FOR INJECTION product monograph. Montreal, Quebec; 25 July 2003.
85. John Korontzis. Personal communication. Regulatory Affairs Associate, Dacarbazine, Mayne Pharma Canada; #FEB-14-2005 (february 8, 2005).
86. Recordati Rare Diseases Inc. COSMEGEN® product monograph. Lebanon, New Jersey USA; 24 July 2014.
87. Andy Harbrow. Personal communication. Global Medical Services Manager, Primevigilance (for Recordati Rare Diseases Inc.); 15 July 2014.
88. Janssen Inc. DARZALEX® product monograph. Toronto, Ontario; 29 June 2016.
89. Erfa Canada Inc. Daunorubicin injection product monograph. Westmount, Quebec; 6 December 2002.
90. Erfa Canada Inc. Material Safety Data Sheet. Montreal, Quebec; 3 October 2007.
91. Henri Knafo MD. Personal communication. Medical Director, Erfa Canada Inc; 14 July 2008.
92. Henri Knafo MD. Personal communication. Medical director, Erfa Canada Inc; 09 July 2008.
93. Novopharm Limited. Daunorubicin Package Insert. Toronto, Canada; Undated.
94. Ferring Pharmaceuticals. FIRMAGON® product monograph. North York, Ontario; 20 March 2013.
95. Ferring Pharmaceuticals. FIRMAGON® product monograph. North York, Ontario; 06 November 2009.
96. Amgen Canada Inc. XGEVA® product monograph. Mississauga, Ontario; 14 October 2011.

97. Pfizer Canada Inc. ZINECARD® product monograph. Kirkland, Quebec; 12 August 2010.
98. Pfizer Canada Inc. ZINECARD® product monograph. Kirkland, Quebec; 30 March 2015.
99. Hospira Healthcare Corporation. DOCETAXEL FOR INJECTION® product monograph. Saint-Laurent, Quebec; 21 February 2011.
100. Hospira Canada Clinical Support Team. Personal communication. Hospira Canada Healthcare Corporation; 21 March 2011.
101. Josee Lloyd, Senior Clinical Specialist. Subject : Docetaxel Injection 160mg/16mL and 80 mg/8mL multidosing and venting needles. Hospira Clinical Support Team; 13 July 2011.
102. sanofi-aventis Canada Inc. TAXOTERE® product monograph. Laval, Quebec; 15 April 2011.
103. Walker S. Stability of docetaxel solution after Dilution in Ethanol and Storage in vials and after Dilution in Normal Saline and Storage in Bags. Can J Hosp Pharm 2007;60(4):231-237.
104. Accord Healthcare Inc. Doxorubicin injection® product monograph. Montreal, Quebec; 9 April 2014.
105. Mayne Pharma (Canada) Inc. Doxorubicin Package Insert. Montreal, QC; Undated.
106. Mayne Pharma (Canada) Inc. Doxorubicin Product Monograph. Montreal, Quebec; 2002.
107. Novopharm Limited. Doxorubicin Product Monograph. Scarborough, Ontario; 8 November 1996.
108. Pfizer Canada Inc. ADRIAMYCIN® injection product monograph. Kirkland, Quebec; 28 August 2007.
109. Janssen Inc. CAELYX® product monograph. Toronto, Ontario; 10 October 2013.
110. Novopharm. Epirubicin for Injection product monograph. Toronto, Ontario; 16 March 2009.
111. Pharmaceutical Partners of Canada, Inc. Epirubicin Hydrochloride Injection product monograph. Richmond Hill, Ontario; 6 July 2010.
112. Pharmacia Canada Inc. Pharmorubicin PFS Package Insert. Mississauga, Ontario; May 2003.
113. BC Cancer Agency Lymphoma Tumour Group. (ULYEPOCHR) Interim BCCA Protocol Summary for Treatment of Lymphoma with Dose-Adjusted Etoposide, DOXOrubicin, VinCRISTine, Cyclophosphamide, PredniSONE, and riTUXimab (LYEPOCHR) with Intrathecal Methotrexate. Vancouver, British Columbia: BC Cancer Agency; 1 July 2015.
114. Barry Goldspiel. Personal communication. NIH Clinical Centre; 14 April 2015.
115. Wolfe JL, Thoma LA, Du C, et al. Compatibility and stability of vincristine sulfate, doxorubicin hydrochloride, and etoposide in 0.9% sodium chloride injection. Am J Health-Syst Pharm 1999;56:985-989.
116. Dunleavy K, Pittaluga S, Shovlin M, et al. Low-intensity therapy in adults with Burkitt's lymphoma. N Engl J Med 2013;369:1915-1925.
117. Eisai Limited. HALAVEN® product monograph. Mississauga, Ontario; 17 January 2013.
118. Sandoz Canada Inc. Etoposide injection® product monograph. Kirkland, Quebec; 27 February 2012.
119. BC Cancer Agency. Provincial Pharmacy Directive III-50-04: Management of Particulate During Sterile Preparation. Vancouver, British Columbia: BC Cancer Agency; 9 July 2014.
120. Novopharm Limited. Etoposide Product Monograph. Toronto, Ontario; 2000.
121. Lepage R, Walker S, Godin J. Stability and compatibility of etoposide in normal saline. Canadian Journal of Hospital Pharmacy 2000;53(5):338-345.
122. The United States Pharmacopeial Convention, Inc. General Chapter 797: Pharmaceutical compounding - sterile preparations. USP 27-NF 22. Rockville, Maryland: The United States Pharmacopeial Convention, Inc.; 2003.
123. Angie Chan. Personal communication. Drug Information Pharmacist, Novopharm. 29 September 2006.
124. Bristol-Myers Squibb Company. ETOPOPHOS® product monograph. Princeton, New Jersey, USA; March 2011.
125. Bristol-Myers Squibb Company. ETOPOPHOS® product monograph. Princeton, New Jersey, USA; September 2013.
126. Amgen Canada Inc. NEUPOGEN® product monograph. Mississauga, Ontario; 21 March 2014.
127. Amgen Medical Information. Personal communication. Amgen Canada Inc.; 8 July 2014.
128. Trissel LA. Handbook on Injectable Drugs. 13th ed. Bethesda, Maryland: American Society of Health-System Pharmacists, Inc; 2005. p. 648-655.
129. Berlex Canada Inc. Fludara Package Insert. Lachine, Quebec; December 1998.
130. Trissel's™2 Clinical Pharmaceutics Database (Parenteral Compatibility) [database on the internet]. Fludarabine. Thomson MICROMEDEX®, Available at: <http://www.micromedex.com/>. Accessed 14 September, 2007.
131. Novopharm Limited. Fludarabine product information package. Toronto, Ontario; 21 June 2007.
132. Accord Healthcare Inc. Fluorouracil injection® product monograph. Kirkland, Quebec; 30 September 2013.
133. Charles Vachon. Personal communication. Quality and Regulatory Affairs, Accord Healthcare Inc.; 29 September 2016.
134. John Korontzis. Personal communication. Regulatory Affairs Associate, Fluorouracil, Mayne Pharma Canada; reference # FEB-43-2005. February 16, 2005.
135. Mayne Pharma (Canada) Inc. Fluorouracil Package Insert. Montreal, Quebec; Undated.
136. Stiles ML, Allen Jr LV, Tu YH. Stability of fluorouracil administered through four portable infusion pumps. American Journal of Hospital Pharmacy 1989;46(10):2036-2040.

137. BC Cancer Agency Experimental Therapeutics. Physicochemical stability analysis of fluorouracil products in final chemotherapeutic preparations. Vancouver, BC. 19 August 2011; Study number 50009:1-43.
138. Sandoz Canada Inc. Fluorouracil Injection product monograph. Boucherville, Quebec; 3 April 2012.
139. Accord Healthcare Inc. Gemcitabine injection® product monograph. Kirkland, Quebec; 29 September 2014.
140. Astron Research LTD. UK. Gemcitabine for Injection (STBRG/ACGEM/01) Stability Study Report (Dilution Study) 2001.
141. Purvi Agrawal BPharm (Regulatory Affairs Manager). Personal communication. Accord Healthcare Inc.; 07 September 2012.
142. Eli Lilly Canada Inc. Gemzar Package Insert. Toronto, Ontario; 16 January 2002.
143. Marilyn Bain BSc N. Personal communication. Sr Therapeutic Area Specialist Medical Information, Gemzar. September 2004.
144. Hospira Healthcare Corporation. Gemcitabine for injection, USP product monograph. Montreal, Quebec; 3 November 2008.
145. Rose Toussaint. Personal communication. Clinical Specialist, Hospira Canada Healthcare Corporation; 26 May 2009.
146. Hospira Canada Clinical Support Team. Personal communication. Hospira Healthcare Corporation; 13 March 2009.
147. Hospira Healthcare Corporation. Gemcitabine Injection product monograph. Montreal, QC; 27 February 2013.
148. Hospira Healthcare Corporation. Gemcitabine Injection product monograph. Montreal, Quebec; 29 August 2014.
149. Novopharm Limited. Gemcitabine product monograph. Toronto, Ontario; 08 February 2008.
150. Sandoz Standard. Gemcitabine for injection product monograph. Boucherville, Quebec; 13 November 2007.
151. Amelie Fontaine B.Sc (N). Personal communication. Drug Information Specialist, Sandoz (vial); 9 April 2009.
152. Stephane Jean. Personal communication. Drug Information Specialist; Sandoz Canada; 7 May 2008.
153. Stephane Jean. Personal communication. Drug Information Specialist, Sandoz Canada; 25 September 2009.
154. Amelie Fontaine B.Sc (N). Personal communication. Drug Information Specialist, Sandoz (infusion); 9 April 2009.
155. Pfizer Canada Inc. IDAMYCIN® product monograph. Kirkland, Quebec; 19 February 2009.
156. Pharmaceutical Partners of Canada, Inc. IDARUBICIN HYDROCHLORIDE INJECTION® product monograph. Richmond Hill, Ontario; 12 November 2009.
157. Baxter Corporation. IFEX® product monograph. Mississauga, Ontario; 5 April 2012.
158. Pharmaceutical Partners of Canada, Inc. Ifosfamide for Injection product monograph. Richmond Hill, Ontario; 17 January 2008.
159. sanofi-aventis Canada. Iniparib (BSI-201; SAR240550) Special Access Program Guidance for the Physician. Laval, Quebec; 15 December 2010.
160. Merck Canada Inc. INTRON A® product monograph. Kirkland, Quebec; 13 March 2015.
161. Edward Kavalec BSc(Pharm), Medical Services Specialist. Personal communication. Merck Canada Inc. Medical Information; 1 April 2015.
162. Bristol Myers Squibb Canada. YERVOY® product monograph. Montreal, Quebec; 1 February 2012.
163. Accord Healthcare Inc. Irinotecan injection® product monograph. Kirkland, Quebec; 6 May 2014.
164. Mayne Pharma (Canada) Inc. Irinotecan Package Insert. Montreal, Quebec; 28 April 2005.
165. Beryl Chan. Personal communication. Mayne Pharma (Canada) Inc. Scientific Affairs Manager, Irinotecan. 2 February 2006.
166. Walker S, Iazzetta J, Law S. Irinotecan stability in vials following puncture when stored at 23c or 4c. Can J Hosp Pharm 2006;59 (Suppl 2):36.
167. Pharmacia Canada Inc. Camptosar Package Insert. Mississauga, Ontario; May 2002.
168. Sandoz Canada Inc. Irinotecan injection package insert. Boucherville, Quebec; April 2008.
169. Baxalta. ONIVYDE® Summary of product characteristics. Vienna, Austria; undated.
170. Merrimack Pharmaceuticals Inc. ONIVYDE® prescribing information. Cambridge, Massachusetts, USA; Oct 2015.
171. Baxalta Canada Corporation. ONIVYDE® product monograph. Toronto, Ontario; 11 August 2017.
172. Bristol-Myers Squibb. IXEMPRA® product monograph. Princeton, New Jersey; 01 October 2007.
173. Kastango ES. The ASHP discussion guide for compounding sterile preparations. Bethesda (MD): American Society of Health-System Pharmacists, Inc.; 2004. p. 5.