

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Leucovorin 50 mg/5 mL 200 mg/20 mL 1000 mg/100 mL (GMP) (F)(PFL) no preservative ¹	N/A	10 mg/mL ¹	50 mg: discard unused portion ^{1,2} 200 mg, 1000 mg: 8 h F ^{1,2}	syringe	8 h RT ^{1,2}	
				0.05-10 mg/mL NS, D5W, Ringer's, LR, D10W, D5-NS ^{1,2} 50-250 mL†	NS , D5W, LR, Ringer's: 24 h RT ¹ D10W, D5-NS: 8 h RT ¹	
Leucovorin 50 mg/5 mL 500 mg/50 mL (Pfizer/Hospira) (F)(PFL) no preservative ³	N/A	10 mg/mL ³	8 h ³	syringe	8 h RT ³	
				0.05–10 mg/mL NS , D5W, LR, Ringer's, D10W, D5NS ³ 50-250 mL†	NS , D5W, LR, Ringer's: 24 h RT ³ D10W, D5NS: 8 h RT ³	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Leucovorin 50 mg/5 mL 500 mg/50 mL (Teva) (F)(PFL) no preservative ⁴	N/A	10 mg/mL ⁵	discard unused portion ⁵	syringe	8 h ^{6,7}	
				0.4 - 4.8 mg/mL NS , D5W ⁸ 50-250 mL†	72 h F , RT ⁸	
				0.06 - 0.4 mg/mL NS , D5W ⁴ 50-250 mL†	NS : 24 h RT ⁴ D5W: 12 h RT ⁴	
				0.06 - 1 mg/mL Ringer's, Lactated Ringer's, D10W, D10-NS ⁴	Ringer's, LR: 24 h RT ⁴ D10W: 12 h RT ⁴ D10NS: 6 h RT ⁴	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Lurbinectedin 4 mg (Jazz) (F) no preservative ⁹	8 mL SWI ⁹	0.5 mg/mL ⁹	12 h F , RT ^{9,10}	100-250 mL NS , D5W ⁹	complete administration within 24 h F , RT ⁹	- larger infusion volume is recommended for peripheral line ⁹ - do not use nylon membrane filters for administration if diluted in NS ⁹ ; BD Alaris pumps and syringe sets have polyethersulfone membrane in-line filters ¹¹
Lurbinectedin 4 mg (Pharma Mar) (F) no preservative ¹² (SAP)	8 mL SWI ¹²	0.5 mg/mL ¹²	12 h F , RT ^{10,12}	100–250 mL NS , D5W ¹²	30 h F , RT ¹²	- larger infusion volume is recommended for peripheral line ¹²

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Melphalan 50 mg (Marcan) (RT)(PFL) no preservative ¹³	10 mL supplied diluent ¹³ rapidly add diluent and immediately shake vigorously to dissolve ¹³ record time of reconstitution	5 mg/mL ¹³	2 h RT ¹³ do NOT refrigerate ¹³	0.1-0.45 mg/mL NS only ¹³	complete administration within 50 min RT from time of initial reconstitution ¹³	- will precipitate if stored in fridge ¹³
Melphalan 50 mg (Taro) (RT)(PFL) no preservative ¹⁴	10 mL supplied diluent ¹⁴ rapidly add diluent and immediately shake vigorously to dissolve ¹⁴ record time of reconstitution	5 mg/mL ¹⁴	2 h RT ¹⁴ do NOT refrigerate ¹⁴	0.1-0.45 mg/mL NS only ¹⁴	complete administration within 50 min RT from time of initial reconstitution ¹⁴	- will precipitate if stored in fridge ¹⁴

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Mesna 400 mg/4 mL 1000 mg/10 mL (Baxter) (RT) no preservative ¹⁵	N/A	100 mg/mL ¹⁵ (use filter needle to withdraw from ampoule)	discard unused portion ¹⁵	≥1 mg/mL NS , D5W, D5½-NS, LR ¹⁵⁻¹⁷ 100 mL†	24 h RT ¹⁵	
Mesna 1000 mg/10 mL 5000 mg/50 mL (Baxter) (RT) preservative ¹⁵	N/A	100 mg/mL ¹⁵	8 d RT ¹⁵ (vial may be punctured up to 4 times) ¹⁵	≥1 mg/mL NS , D5W, D5½-NS, LR ¹⁵⁻¹⁷ 100 mL†	24 h RT ¹⁵	
Mesna 1000 mg/10 mL (Fresenius Kabi) (RT) preservative ¹⁸	N/A	100 mg/mL ¹⁸	14 d F , RT ^{10,18}	≥1 mg/mL NS , D5W ¹⁹ 100 mL†	48 h F, 24 h RT ¹⁸	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Methotrexate 50 mg/2 mL 500 mg/20 mL 1 g/40 mL (Accord) (RT)(PFL) no preservative ²⁰	N/A	25 mg/mL ²⁰	50mg: discard unused portion ²⁰ 500 mg, 1 g: 8 h RT ²⁰	syringe	use within 8 h RT of initial puncture ²⁰	- for high-dose regimens (e.g., 1-12 g/m ² as a single dose): use preservative-free methotrexate ²⁰ - do not use for IT injection
				0.4–2 mg/mL NS , D5W ²⁰ 50-500 mL†	use within 24 h RT of initial puncture ²⁰ **(PFL)	
				high dose (e.g., 1-12 g/m ² as a single dose): 1000 mL * NS	use within 24 h RT of initial puncture ²⁰ **(PFL)	
<u>Methotrexate intravitreal injection</u> 50 mg/2 mL (Accord) (RT)(PFL) no preservative ²⁰	N/A	25 mg/mL ²⁰	discard unused portion ²⁰	syringe for intravitreal use	use within 4 h of initial puncture ¹⁰	- for intravitreal use preservative-free methotrexate is preferred ²¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Methotrexate IT Injection only preservative free methotrexate may be administered by the intrathecal route ²² 50 mg/2 mL (Accord) (RT)(PFL) no preservative ²⁰	N/A	25 mg/mL ²⁰	discard unused portion ²⁰	IT syringe qs to 6 mL with preservative free NS ^{23,24} diluent containing preservatives should NOT be used for intrathecal administration ²⁵	use within 4 h of initial puncture ¹⁰	- auxiliary info ¹⁰ : IT - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer Ziploc bag ²⁶
Methotrexate 50 mg/2 mL 500 mg/20 mL (Accord) (RT)(PFL) preservative ²⁰	N/A	25 mg/mL ²⁰	28 d F ^{10,20}	syringe	10 d F ^{10,20}	- contains benzyl alcohol ²⁰ - do NOT use for high-dose regimens (e.g., 1-12 g/m ² as a single dose) ²⁰ - do NOT use for IT injection ²⁰
				0.4–2 mg/mL NS, D5W ²⁰ 50-500 mL†	24 h RT ²⁰	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Methotrexate 1 g (Fresenius Kabi USA) (RT)(PFL) no preservative ²⁷	39.4 mL NS, D5W ²⁸	25 mg/mL ²⁸	discard unused portion ²⁷	syringe	use within 4 h of initial puncture ¹⁰	- for high-dose regimens (e.g., 1-12 g/m ² as a single dose): use preservative-free methotrexate
				0.4–2 mg/mL NS , D5W ²⁰ 50-500 mL†	use within 4 h of initial puncture ¹⁰	
				high dose (e.g., 1-12 g/m ² as a single dose): 1000 mL* D5W ²⁷ , NS ²⁰	use within 4 h of initial puncture ¹⁰	
Methotrexate intravitreal injection 1 g (Fresenius Kabi USA) (RT)(PFL) no preservative ²⁷	39.4 mL NS, D5W ²⁸	25 mg/mL ²⁸	discard unused portion ²⁷	syringe for intravitreal use	use within 4 h of initial puncture ¹⁰	- for intravitreal use preservative-free methotrexate is preferred ²¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Methotrexate IT Injection Only preservative free methotrexate may be administered by the intrathecal route ²² 1 g (Fresenius Kabi USA) (RT)(PFL) no preservative ²⁷	39.4 mL NS, D5W ²⁸	25 mg/mL ²⁸	discard unused portion ²⁷	IT syringe qs to 6 mL with preservative free NS ^{23,24} diluents containing preservatives should NOT be used for intrathecal administration ²⁵	use within 4 h of initial puncture ¹⁰	- auxiliary info ¹⁰ : IT - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer Ziploc bag ²⁶
Methotrexate 50 mg/2 mL 500 mg/20 mL 1 g/40 mL 2.5 g/100 mL (Pfizer/Hospira) (RT)(PFL) no preservative ²⁹	N/A	25 mg/mL ²⁹	50mg: discard unused portion ²⁹ 500 mg, 1 g, or 2.5 g: 8 h RT ²⁹	syringe	use within 8 h RT of initial puncture ²⁹	- for high-dose regimens (e.g., 1-12 g/m ² as a single dose): use preservative-free methotrexate ²⁹ - do not use for IT injection
				0.4–2 mg/mL NS , D5W ²⁹ 50-500 mL†	use within 24 h RT of initial puncture ²⁹ **(PFL)	
				high dose (e.g., 1-12 g/m ² as a single dose): 1000 mL * NS	use within 24 h RT of initial puncture ²⁹ **(PFL)	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Methotrexate intravitreal injection 50 mg/2 mL (Pfizer/Hospira) (RT)(PFL) no preservative ²⁹	N/A	25 mg/mL ²⁹	discard unused portion ²⁹	syringe for intravitreal use	use within 4 h of initial puncture ¹⁰	- for intravitreal use preservative-free methotrexate is preferred ²¹
Methotrexate IT Injection only preservative free methotrexate may be administered by the intrathecal route ²² 50 mg/2 mL (Pfizer/Hospira) (RT)(PFL) no preservative ²⁹	N/A	25 mg/mL ²⁹	discard unused portion ²⁹	IT syringe qs to 6 mL with preservative free NS ^{23,24} diluent containing preservatives should NOT be used for intrathecal administration ²⁵	use within 4 h of initial puncture ¹⁰	- auxiliary info ¹⁰ : IT - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer Ziploc bag ²⁶
Methotrexate 50 mg/2 mL 500 mg/20 mL (Pfizer/Hospira) (RT)(PFL) preservative ²⁹	N/A	25 mg/mL ²⁹	28 d F ^{10,29}	syringe	10 d F ^{10,29}	- contains benzyl alcohol ²⁹ - do NOT use for high-dose regimens (e.g., 1-12 g/m ² as a single dose) ²⁹ - do NOT use for IT injection ²⁹
				0.4–2 mg/mL NS, D5W ²⁹ 50-500 mL†	24 h RT ²⁹	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Mirvetuximab soravtansine 100 mg (AbbVie EU) (F)(PFL) do not shake no preservative ³⁰ (SAP)	n/a	5 mg/mL ³⁰ allow vials to reach RT before using ³⁰	discard unused portion ³⁰	dilute with D5W in empty infusion bag to final concentration of 1-2 mg/mL ³⁰ qs to 200-600 mL† do not use NS ³⁰ gently invert to mix; do not shake ³⁰	8 h RT or 24 h F plus an additional 8 h RT, including infusion ³⁰ if refrigerated, bring bag to RT prior to administration ³⁰	- incompatible with saline ³⁰ - do not use if discoloured or cloudy, or has particulates ³⁰ - use non-DEHP bag and tubing ³⁰ - administer using 0.2 micron in-line filter ³⁰
Mitomycin 20 mg (Accord) (RT)(PFL) no preservative ³¹	40 mL SWI ³¹ shake well ³¹	0.5 mg/mL ³¹	12 h F, 6 h RT ^{10,31} **(PFL) ³¹	syringe	72 h F, 6 h RT ³¹ **(PFL) ³¹	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Mitomycin intravesical 20 mg (Accord) (RT)(PFL) no preservative ³¹	40 mL SWI ³¹ shake well ³¹	0.5 mg/mL ³¹	12 h F, 6 h RT ^{10,31} **(PFL) ³¹	syringe	72 h F, 6 h RT ³¹ **(PFL) ³¹	
	10 mL SWI ³² shake well ³¹	2 mg/mL ³²	use immediately after preparation to prevent precipitation ³³	syringe	use immediately after preparation to prevent precipitation ³³	- may precipitate due to low solubility ^{33,34} - do NOT refrigerate ³³
	25 mL SWI shake well	0.8 mg/mL ³⁵	discard unused portion ^{2,35} **(PFL) ^{2,35}	syringe	4 days RT ³⁵ **(PFL) ^{2,35}	- do NOT refrigerate ³⁵
	33.3 mL SWI shake well	0.6 mg/mL ³⁵	discard unused portion ^{2,35} **(PFL) ^{2,35}	syringe	4 days F, RT ³⁵ **(PFL) ^{2,35}	
Mitomycin intraperitoneal 20 mg (Accord) (RT)(PFL) no preservative ³¹	40 mL SWI ³¹ shake well ³¹	0.5 mg/mL ³¹	12 h F, 6 h RT ^{10,31} **(PFL) ³¹	0.02-0.04 mg/mL NS , sodium lactate ³¹	NS ³¹ : 18 h F, 3 h RT sodium lactate ³¹ : 6 h F, 3 h RT	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Mitomycin 20 mg (Hikma) (RT)(PFL) no preservative ³⁶	40 mL SWI ³⁶ shake well ³⁶	0.5 mg/mL ³⁶	12 h F, 6 h RT ^{10,36} **(PFL) ³⁶	syringe	72 h F, 6 h RT ³⁶ **(PFL) ³⁶	
Mitomycin intravesical 20 mg (Hikma) (RT)(PFL) no preservative ³⁶	40 mL SWI ³⁶ shake well ³⁶	0.5 mg/mL ³⁶	12 h F, 6 h RT ^{10,36} **(PFL) ³⁶	syringe	72 h F, 6 h RT ³⁶ **(PFL) ³⁶	
	10 mL SWI ³² shake well	2 mg/mL ³²	use immediately after preparation to prevent precipitation ³³	syringe	use immediately after preparation to prevent precipitation ³³	- may precipitate due to low solubility ^{33,34} - do NOT refrigerate ³³
	25 mL SWI shake well	0.8 mg/mL ³⁵	discard unused portion ^{10,35} **(PFL) ^{10,35}	syringe	4 days RT ³⁵ **(PFL) ^{10,35}	- do NOT refrigerate ³⁵
	33.3 mL SWI shake well	0.6 mg/mL ³⁵	discard unused portion ^{10,35} **(PFL) ^{10,35}	syringe	4 days F, RT ³⁵ **(PFL) ^{10,35}	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Mitomycin <u>intraperitoneal</u> 20 mg (Hikma) (RT)(PFL) no preservative ³⁶	40 mL SWI ³⁶ shake well ³⁶	0.5 mg/mL ³⁶	12 h F, 6 h RT ^{10,36} **(PFL) ³⁶	0.02-0.04 mg/mL NS ³⁶	NS ³⁶ : 18 h F, 3 h RT	
Mitomycin 20 mg (Teva) (RT)(PFL) no preservative ³⁷	40 mL SWI ³⁷ shake well ³⁷	0.5 mg/mL ³⁷	12 h F, 6 h RT ^{10,37} **(PFL) ³⁷	syringe	72 h F, 6 h RT ³⁷ **(PFL) ³⁷	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Mitomycin intravesical 20 mg (Teva) (RT)(PFL) no preservative ³⁷	40 mL SWI ³⁷ shake well ³⁷	0.5 mg/mL ³⁷	12 h F, 6 h RT ^{10,37} **(PFL) ³⁷	syringe	72 h F, 6 h RT ³⁷ **(PFL) ³⁷	
	10 mL SWI ³² shake well ³⁷	2 mg/mL ³²	use immediately after preparation to prevent precipitation ³³	syringe	use immediately after preparation to prevent precipitation ³³	- may precipitate due to low solubility ^{33,34} - do NOT refrigerate ³³
	25 mL SWI shake well	0.8 mg/mL ³⁵	discard unused portion ^{2,35} **(PFL) ^{2,35}	syringe	4 days RT ³⁵ **(PFL) ^{2,35}	- do NOT refrigerate ³⁵
	33.3 mL SWI shake well	0.6 mg/mL ³⁵	discard unused portion ^{2,35} **(PFL) ^{2,35}	syringe	4 days F, RT ³⁵ **(PFL) ^{2,35}	
Mitomycin intraperitoneal 20 mg (Teva) (RT)(PFL) no preservative ³⁷	40 mL SWI ³⁷ shake well ³⁷	0.5 mg/mL ³⁷	12 h F, 6 h RT ^{10,37} **(PFL) ³⁷	0.02-0.04 mg/mL NS , sodium lactate ³⁷	NS ³⁷ : 18 h F, 6 h RT sodium lactate ³⁷ : 6 h F, RT	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
mitoXANTRONE 20 mg/10 mL (Fresenius Kabi) (RT) no preservative ³⁸	N/A	2 mg/mL ³⁸	discard unused portion ³⁸	0.2-0.6 mg/mL NS , D5W ³⁸ 50 mL†	24 h RT ³⁸	
mitoXANTRONE 20 mg/10 mL 25 mg/12.5 mL 30 mg/15 mL (Pfizer/Hospira) (RT)(PFL) no preservative ³⁹	N/A	2 mg/mL ³⁹	discard unused portion ³⁹	0.2-0.6 mg/mL NS , D5W ³⁹ 50 mL†	72 h F, 24 h RT ³⁹ **(PFL) ³⁹	
Mogamulizumab 20 mg/5 mL (Kyowa) (F)(PFL) do not shake no preservative ⁴⁰	N/A	4 mg/mL ⁴⁰	discard unused portion ⁴⁰	0.1-3 mg/mL NS 100 mL* mix by gentle inversion; do not shake ⁴⁰	24 h F ⁴⁰	- discard if cloudy, discoloured, or visible particulates are present ⁴⁰ - administer with 0.2 micron in-line filter ⁴⁰
Nelarabine 250 mg/50 mL (Sandoz) (RT) no preservative ⁴¹	N/A	5 mg/mL ⁴¹	discard unused portion ⁴¹	undiluted in empty PVC infusion bag or glass container ⁴¹	8 h RT ⁴¹	- discard if discoloured, hazy, or particulates are present ⁴¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Nivolumab 40 mg/4 mL 100 mg/10 mL (BMS) (F)(PFL) do not shake no preservative ⁴²	N/A	10 mg/mL ⁴²	discard unused portion ⁴²	1-10 mg/mL NS , D5W ⁴² 25-100 mL† mix by gentle inversion; do not shake ⁴² OR undiluted in empty infusion bag or glass container ⁴²	complete administration within 7 days F, including max 8 h at RT ⁴² **(PFL) ⁴² (can be in room light when at RT) ⁴²	- do not shake ⁴² - administer with 0.2 micron in-line filter ⁴² - may contain a few amorphous particles ⁴² - discard if cloudy, has pronounced colour change (should be clear to pale yellow) ⁴²
Nivolumab-relatlimab 240 mg-80 mg/20 mL (BMS) (F)(PFL) do not shake no preservative ⁴³	N/A	12 mg/mL nivolumab- 4 mg/mL relatlimab	discard unused portion ⁴³	3-12 mg/mL nivolumab 50-100 mL† NS , D5W ⁴³ mix by gentle inversion; do not shake ⁴³ (OR undiluted in empty infusion bag or glass container ⁴³)	complete administration within 24 h F, 8 h RT ⁴³ **(PFL) ⁴³ (can be in room light when at RT) ⁴³	- do not shake ⁴³ - administer with a 0.2 micron in-line filter ⁴³ - discard if cloudy, discoloured or contains particulates ⁴³ - may contain a few translucent-to- white particles ⁴³

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
oBINutuzumab 1000 mg/40 mL (Roche) (F)(PFL) do not shake no preservative ⁴⁴	N/A	25 mg/mL ⁴⁴	discard unused portion ⁴⁵	NS 100 mg: 100 mL ⁴⁴ 900 mg: 250 mL ⁴⁴ 1000 mg: 250 mL ⁴⁴ do NOT use dextrose containing solutions ⁴⁴	24 h F, 48 h RT ^{44,46}	- once removed from the fridge, diluted product is stable for an additional 48 h RT ^{44,46} - do NOT shake ⁴⁴
Octreotide 50 mcg/1 mL 100 mcg/1 mL 500 mcg/1 mL (Omega) (F)(PFL) no preservative ⁴⁷	N/A	50 mcg/mL ⁴⁷	discard unused portion ⁴⁷	NS ⁴⁷ volume adjusted to ensure a continuous infusion of octreotide at 25 mcg/h ⁴⁷	24 h RT ⁴⁷	
		100 mcg/mL ⁴⁷				
		500 mcg/mL ⁴⁷				

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Octreotide multidose vial: 1000 mcg/5 mL (Omega) (F)(PFL) preservative ⁴⁷	N/A	200 mcg/mL ⁴⁷	15 d F ⁴⁷	NS ⁴⁷ volume adjusted to ensure a continuous infusion of octreotide at 25 mcg/h ⁴⁷	24 h RT ⁴⁷	
Octreotide (SANDOSTATIN®) 50 mcg/1 mL 100 mcg/1 mL 500 mcg/1 mL (Novartis) (F)(PFL) no preservative ⁴⁸	N/A	50 mcg/mL ⁴⁸	discard unused portion ⁴⁸	NS ⁴⁸ volume adjusted to ensure a continuous infusion of octreotide at 25 mcg/h ⁴⁸	24 h RT ⁴⁸	
		100 mcg/mL ⁴⁸				
		500 mcg/mL ⁴⁸				
Octreotide (SANDOSTATIN®) multi-dose vial: 1000 mcg/5 mL (Novartis) (F)(PFL) preservative ⁴⁸	N/A	200 mcg/mL ⁴⁸	14 d F, RT ⁴⁸	NS ⁴⁸ volume adjusted to ensure a continuous infusion of octreotide at 25 mcg/h ⁴⁸	24 h RT ⁴⁸	
Octreotide (SANDOSTATIN LAR®) (long acting) 10 mg	2 mL supplied diluent ⁴⁸	10 mg: 5 mg/mL ⁴⁸	discard unused portion ⁴⁸	syringe (for deep intragluteal administration only) ⁴⁸	use within 4 h of initial	- do NOT shake ⁴⁸

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
20 mg 30 mg (Novartis) (F)(PFL) no preservative ⁴⁸	add diluent: gently run diluent down sides of vial ⁴⁸ do NOT disturb for 2–5 min; then swirl moderately ⁴⁸ record time of reconstitution	20 mg: 10 mg/mL ⁴⁸			reconstitution ^{10,48}	
		30 mg: 15 mg/mL ⁴⁸				
Octreotide suspension (long acting) 10 mg 20 mg 30 mg (Teva) (F)(PFL)	2 mL supplied diluent let stand at RT for 30 min prior to reconstitution ⁴⁹	10 mg: 5 mg/mL ⁴⁹	discard unused portion ⁴⁹	syringe (for deep intragluteal administration only) ⁴⁹	use within 4 h of initial reconstitution ^{10,49}	- gently shake to resuspend before administration ⁴⁹ - delay in administration may result in sedimentation ⁴⁹
		20 mg: 10 mg/mL ⁴⁹				

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
no preservative ⁴⁹	<p>add supplied diluent ⁴⁹</p> <p>let vial stand for 5 min after adding diluent to saturate powder ⁴⁹</p> <p>shake moderately in horizontal direction for ≥30 sec to create suspension ⁴⁹</p> <p>record time of reconstitution</p>	<p>30 mg: 15 mg/mL ⁴⁹</p>				

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Oxaliplatin 50 mg/10 mL 100 mg/20 mL 200 mg/40 mL (Dr. Reddy's) (RT)(PFL) no preservative ⁵⁰	N/A	5 mg/mL ⁵⁰	discard unused portion ⁵⁰	0.2-0.7 mg/mL D5W ⁵⁰ 100-500 mL† do NOT use NS or other chloride- containing solution ⁵⁰ do NOT use aluminum-containing needle and syringe ⁵⁰	0.2-2 mg/mL: 48 h F, 24 h RT ⁵⁰	- do NOT use aluminum- containing needle, syringe, or tubing ⁵⁰
Oxaliplatin 50 mg/10 mL 100 mg/20 mL 200 mg/40 mL (Pfizer/Hospira) (RT) no preservative ⁵¹	N/A	5 mg/mL ⁵¹	discard unused portion ⁵¹	0.2-0.7 mg/mL D5W ⁵¹ 100-500 mL† do NOT use NS or other chloride- containing solutions ⁵¹ do NOT use aluminum-containing needle and syringe ⁵¹	0.2-0.4 mg/mL: 24 h RT ⁵¹ or 5 d F plus an additional 8 h RT ⁵² 0.5–2 mg/mL: 24 h RT ⁵¹ or 10 d F, plus an additional 8 h RT ^{10,52} **(PFL) when stored in F ⁵²	- do NOT use aluminum- containing needle, syringe, tubing ⁵¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Oxaliplatin 50 mg/10 mL 100 mg/20 mL 150 mg/30 mL 200 mg/40 mL (Sandoz) (RT)(PFL) no preservative ⁵³	N/A	5 mg/mL ⁵³	12 h F, RT ^{10,54}	0.2-0.7 mg/mL D5W ⁵³ 100-500 mL† do NOT use NS or other chloride- containing solution ⁵³ do NOT use aluminum-containing needle and syringe ⁵³	0.2-2 mg/mL: 48 h F, 24 h RT ⁵³	- do NOT use aluminum- containing needle, syringe, tubing ⁵³
Oxaliplatin 50 mg/10 mL 100 mg/20 mL 200 mg/40 mL (Teva) (RT)(PFL) no preservative ⁵⁵	N/A	5 mg/mL ⁵⁵	discard unused portion ⁵⁵	0.2-0.7 mg/mL D5W ⁵⁵ 100-500 mL† do NOT use NS or other chloride- containing solution ⁵⁵ do NOT use aluminum-containing needle and syringe ⁵⁵	0.2-2 mg/mL: 48 h F, 24 h RT ⁵⁵	- do NOT use aluminum- containing needle, syringe or tubing ⁵⁵

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
PACLitaxel 30 mg/5 mL 100 mg/16.7 mL 300 mg/50 mL (Accord) (RT)(PFL) preservative ⁵⁶	N/A	6 mg/mL ⁵⁶	30 mg, 100 mg: 28 d RT ^{10,56} 300 mg: 24 h RT ^{10,56}	0.3-1.2 mg/mL NS , D5W, D5NS, D5LR ⁵⁶ 50-500 mL†	complete administration within 27 h RT ⁵⁶	- use non-DEHP bag and tubing ⁵⁶ - administer with 0.2 micron in-line filter ⁵⁶ - avoid excessive shaking ⁵⁶
				0.1 mg/mL NS ⁵⁷	44 h F , RT ⁵⁷	
PACLitaxel 30 mg/5 mL 100 mg/16.7 mL 300 mg/50 mL (Biolyse) (RT) preservative ⁵⁸	N/A	6 mg/mL ⁵⁸	28 d RT ⁵⁹	0.3-1.2 mg/mL NS , D5W ⁵⁸ 50-500 mL†	complete administration within 27 h RT ^{60,61}	- use non-DEHP bag and tubing ⁵⁸ - administer with 0.2 micron in-line filter ⁵⁸
				0.1 mg/mL NS ⁵⁷	44 h F , RT ⁵⁷	
				0.012-0.12 mg/mL NS ⁶²	16 h RT ⁶⁰	
				devices with spikes (e.g., chemo dispensing pins) may be used with vials ⁶³		

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Paclitaxel 30 mg/5 mL 100 mg/16.7 mL 300 mg/50 mL (Sandoz) (RT)(PFL) preservative ⁶⁴	N/A	6 mg/mL ⁶⁴	30 mg, 100 mg: 28 d RT ^{10,64} 300 mg: 24 h RT ^{10,64}	0.3-1.2 mg/mL NS , D5W, D5NS ⁶⁴ 50-500 mL†*	complete administration within 27 h RT ⁶⁴	- use non-DEHP bag and tubing ⁶⁴ - administer with 0.2 micron inline filter ⁶⁴ - avoid excessive shaking ⁶⁴
				0.1 mg/mL NS ⁵⁷	44 h F , RT ⁵⁷	
PACLitaxel, nanoparticle, albumin- bound (NAB) 100 mg (BMS/Celgene) (RT)(PFL) no preservative ⁶⁵	20 mL NS ⁶⁵ slowly direct diluent against side of vial (i.e., ≥1 min) during reconstitution ⁶⁵ let stand for ≥5 min to wet powder ⁶⁵ gently swirl or invert for ≥2 min ⁶⁵	5 mg/mL ⁶⁵	use immediately (RT) or 8 h F ⁶⁵ **(PFL) ⁶⁵	undiluted in empty PVC, non-PVC, or non-DEHP infusion bag ⁶⁵	48 h F plus an additional 8 h RT ⁶⁶	- each vial contains 900 mg human albumin ⁶⁵ - to prevent foaming, do NOT inject NS directly onto the powder ⁶⁵ - some settling may occur; use mild agitation to resuspend ⁶⁵ - administer with 15 micron filter ONLY ⁶⁵ (NOTE: filters with pore size less than 15 microns may cause filter blockage) ⁶⁵

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
PACLitaxel, nanoparticle, albumin- bound (NAB) 100 mg (Panacea/Apo) (RT)(PFL) no preservative ⁶⁷	20 mL NS ⁶⁷ slowly direct diluent against side of vial (i.e., ≥1 min) during reconstitution ⁶⁷ let stand for ≥5 min to wet powder ⁶⁷ gently swirl or invert for ≥2 min ⁶⁷ (if foaming occurs, let stand for ≥15 min) ⁶⁷	5 mg/mL ⁶⁷	use immediately (RT) or 8 h F ⁶⁷ **(PFL) ⁶⁷	undiluted in empty PVC, non-PVC, or non-DEHP infusion bag ⁶⁷	56 h F plus an additional 4 h RT ⁶⁸	- each vial contains 900 mg human albumin ⁶⁷ - to prevent foaming, do NOT inject NS directly onto the powder ⁶⁷ - some settling may occur; use gentle inversion to resuspend ⁶⁷ - discard if visible particulates are present ⁶⁷ - administer with 15 micron filter ONLY ⁶⁷
Pamidronate 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Fresenius Kabi) (RT) no preservative ⁶⁹	N/A	3 mg/mL ⁶⁹	discard unused portion ⁶⁹	≤0.36 mg/mL ⁶⁹ NS , D5W ⁶⁹ 250 mL†	24 h RT ⁶⁹	- do NOT mix with calcium containing solutions (e.g., LR) ⁶⁹
		6 mg/mL ⁶⁹				
		9 mg/mL ⁶⁹				

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Pamidronate 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Hospira) (RT) no preservative ⁷⁰	N/A	3 mg/mL ⁷⁰	discard unused portion ⁷⁰	0.06–0.36 mg/mL NS , D5W ⁷⁰ 250 mL†	24 h F plus an additional 24 h RT (total 48 h) ⁷⁰ **(PFL) ⁷⁰	- do NOT mix with calcium containing solution (e.g., LR) ⁷⁰
		6 mg/mL ⁷⁰				
		9 mg/mL ⁷⁰				
Pamidronate 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Omega) (RT) no preservative ⁷¹	N/A ⁷¹	3 mg/mL ⁷¹	discard unused portion ⁷¹	0.06–0.36 mg/mL NS , D5W ⁷¹ 250 mL†	24 h F plus an additional 24 h RT (total 48 h) ⁷¹ **(PFL) ⁷¹	- do NOT mix with calcium containing solution (e.g., LR) ⁷¹
		6 mg/mL ⁷¹				
		9 mg/mL ⁷¹				
Pamidronate 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Pfizer) (RT) no preservative ⁷²	N/A	3 mg/mL ⁷²	discard unused portion ⁷²	0.06–0.36 mg/mL NS , D5W ⁷² 250 mL†	24 h F plus an additional 24 h RT (total 48 h) ⁷² **(PFL) ⁷²	- do NOT mix with calcium containing solution (e.g., LR) ⁷²
		6 mg/mL ⁷²				
		9 mg/mL ⁷²				

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Pamidronate 30 mg/10 mL 60mg/10 mL 90 mg/10 mL (Sandoz Canada) RT no preservative ⁷³	N/A	3 mg/mL ⁷³	discard unused portion ^{73,74}	NS ; D5W ⁷³ 250 mL†	24 h RT ⁷³	- do NOT mix with calcium containing solution (e.g., LR) ⁷³
		6 mg/mL ⁷³				
		9 mg/mL ⁷³				
PANitumumab 100 mg/5 mL 400 mg/20 mL (Amgen) (F)(PFL) do not shake no preservative ⁷⁵	N/A	20 mg/mL ⁷⁵	discard unused portion ⁷⁵	1-10mg/mL NS ⁷⁵ 100 mL†	24 h F, 6 h RT ⁷⁵⁻⁷⁸	- administer with 0.2 micron in-line filter ⁷⁵ - solution may contain particulates which do not affect product quality ⁷⁵ - do not administer if discoloured ⁷⁵

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Pegaspargase (pegylated asparaginase <i>E. coli</i>) 3750 units/5 mL (Servier) (F)(PFL) do not shake no preservative ⁷⁹	N/A	750 units/mL ⁷⁹	discard unused portion ⁷⁹	IM ⁷⁹ : max volume: 2 mL in children and adolescents; 3 mL in adults if volume greater than above, use multiple sites ⁷⁹	syringe: use within 4 h of vial puncture ^{2,79}	- do NOT shake ⁷⁹
				IV ⁷⁹ : 100 mL NS , D5W	bag: use within 4 h of vial puncture ^{2,79}	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Pegaspargase (pegylated asparaginase <i>E. coli</i>) 3750 units/5 mL (Servier USA) (F)(PFL) do not shake no preservative ^{79,80}	N/A	750 units/mL ⁷⁹	discard unused portion ⁷⁹	IM ⁷⁹ : max volume: 2 mL in children and adolescents; 3 mL in adults if volume greater than above, use multiple sites ⁷⁹	syringe: use within 4 h of vial puncture ^{2,79}	- do NOT shake ⁷⁹
				IV ⁷⁹ : 100 mL NS , D5W	bag: use within 4 h of vial puncture ^{2,79}	
Pembrolizumab 100 mg/4 mL (Merck) (F)(PFL) do not shake no preservatives ⁸¹	N/A	25 mg/mL ⁸¹	discard unused portion ^{2,81}	1-10 mg/mL NS , D5W ⁸¹ 50 mL* mix by gentle inversion ⁸¹	complete administration within 96 h F, 6 h RT ⁸¹	- administer with 0.2 micron in-line filter ⁸¹ - bring vials and diluted solutions to RT prior to use ⁸¹ - vials contain 0.25 mL overfill ⁸¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Pemetrexed 100 mg 500 mg (Accord) (RT) no preservative ⁸²	100 mg: 4.2 mL NS ⁸² 500 mg: 20 mL NS ⁸²	25 mg/mL ⁸²	12 h F, RT ^{10,82}	100 mL NS ⁸²	24 h F, RT ⁸²	- do NOT mix with calcium containing solution (e.g., LR) ⁸²
Pemetrexed 100 mg/4 mL 500 mg/20 mL 1000 mg/40 mL (Accord) (RT)(PFL) no preservative ⁸³	N/A	25 mg/mL ⁸³	discard unused portion ⁸³	100 mL NS ⁸³	24 h F ⁸³	- do NOT mix with calcium containing solution (e.g., LR) ⁸³
Pemetrexed 100 mg 500 mg (Dr. Reddy's) (RT) no preservative ⁸⁴	100 mg: 4.2 mL NS ⁸⁴ 500 mg: 20 mL NS ⁸⁴	25 mg/mL ⁸⁴	12 h F, RT ^{10,85-87}	100 mL NS ⁸⁴	24 h F, RT ⁸⁵⁻⁸⁷	- do NOT mix with calcium containing solution (e.g., LR) ⁸⁴
Pemetrexed 100 mg 500 mg (Lilly) (RT) no preservative ⁸⁸	100 mg: 4.2 mL NS ⁸⁸ 500 mg: 20 mL NS ⁸⁸	25 mg/mL ⁸⁸	12 h F ^{10,88}	100 mL NS ⁸⁸	24 h F ⁸⁸	- do NOT mix with calcium containing solution (e.g., LR) ⁸⁸

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Pemetrexed 100 mg 500 mg 1000 mg (Taro) (RT) no preservative ⁸⁹	100 mg: 4.2 mL NS ⁸⁹ 500 mg: 20 mL NS ⁸⁹ 1000 mg: 40 mL NS ⁸⁹	25 mg/mL ⁸⁹	12 h F ^{10,89}	100 mL NS ⁸⁹	24 h F ⁸⁹	- do NOT mix with calcium containing solution (e.g., LR) ⁸⁹
Pentostatin 10 mg (Hospira/Pfizer) (F) no preservative ⁹⁰	5 mL SWI ⁹⁰	2 mg/mL ⁹⁰	8 h RT ⁹⁰	0.18-0.33 mg/mL ⁹⁰ 25-50 mL NS , D5W ⁹⁰	8 h RT ⁹⁰	
PERTuzumab 420 mg/14 mL (Roche) (F)(PFL) no preservative ⁹¹	N/A	30 mg/mL ⁹¹ do NOT shake ⁹¹	discard unused portion ^{45,91}	250 mL NS only ⁹¹ mix by gentle inversion to avoid foaming ⁹¹ do NOT use dextrose containing solutions ⁹¹	24 h F, RT ⁹¹	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
PERTuzumab- trastuzumab 1200 mg-600 mg/15 mL 600 mg-600 mg/10 mL (Roche) (F)(PFL) do not shake no preservative ⁹²	N/A	1200 mg -600 mg ⁹²: 80 mg/mL pertuzumab and 40 mg/mL trastuzumab 600 mg -600 mg ⁹²: 60 mg/mL pertuzumab and 60 mg/mL trastuzumab	discard unused portion ⁹²	SC syringe ⁹²	10 d F, 24 h RT ^{10,92}	- do not shake ⁹² - contains recombinant human hyaluronidase ⁹²
Plerixafor 24 mg/1.2 mL (sanofi-aventis) (RT) no preservative ⁹³	N/A	20 mg/mL ⁹³	discard unused portion ⁹³	SC syringe ⁹³	48 h RT ^{74,94}	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Polatuzumab vedotin 30 mg 140 mg (Hoffman-La Roche) (F)(PFL) do not shake no preservative ⁹⁵	30 mg: 1.8 mL SWI ⁹⁵ 140 mg: 7.2 mL SWI ⁹⁵ direct diluent against side of vial during reconstitution ⁹⁵ swirl gently to mix ⁹⁵	20 mg/mL ⁹⁵ (PFL)	12 h F, RT ^{10,95}	0.72-2.7 mg/mL NS , D5W, ½NS ⁹⁵ (dilute to a minimum volume of 50 mL) ⁹⁵ gently invert bag to mix ⁹⁵	in NS: 72 h F, 4 h RT ⁹⁵ in D5W or ½NS: 72 h F, 8 h RT ⁹⁵	- do NOT shake ⁹⁵ - administer with 0.2 micron in-line filter ⁹⁵ -discard if discolouration or visible particulates are present ⁹⁵
Pralatrexate 20 mg/1 mL 40 mg/2 mL (Servier) (F)(PFL) no preservative ⁹⁶	N/A	20 mg/mL ⁹⁶	discard unused portion ²	syringe ⁹⁶	24 h F, RT ⁹⁷ **(PFL) ⁹⁷	- do NOT dilute ⁹⁶
Raltitrexed 2 mg (Pfizer) (F,RT)(PFL) no preservative ⁹⁸	4 mL SWI ⁹⁸	0.5 mg/mL ⁹⁸	12 h F, RT ^{10,98}	50-250 mL NS , D5W ⁹⁸	complete administration within 24 h F, RT ⁹⁸	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Ramucirumab 100 mg/10 mL 500 mg/50 mL (Eli Lilly) (F)(PFL) (do not shake) no preservative ⁹⁹	N/A	10 mg/mL ⁹⁹	discard unused portion ⁹⁹	0.4–4 mg/mL NS ^{99,100} 250-500 mL† gently invert to mix ⁹⁹ do NOT shake ⁹⁹ do NOT use dextrose containing solutions ⁹⁹	24 h F, 4 h RT ⁹⁹	- administer with 0.2 micron in-line filter ⁹⁹
Retifanlimab 500 mg/20 mL (Incyte) (F)(PFL) do not shake no preservative ¹⁰¹ (SAP)	N/A	25 mg/mL ¹⁰¹	discard unused portion ¹⁰¹	1.4-10 mg/mL NS , D5W ¹⁰¹ 50 mL* mix by gentle inversion ¹⁰¹ ; do not shake ¹⁰¹	complete administration within 24 h F or 8 h RT ¹⁰¹ if refrigerated, bring to RT prior to use and complete administration within 4 h RT after removing from fridge ¹⁰¹ **(PFL) ¹⁰¹	- discard if cloudy, discoloured, or contains particulates ¹⁰¹ - administer using 0.2 micron filter ¹⁰¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
riTUXimab (RITUXAN®) 100 mg/10 mL 500 mg/50 mL (Roche) (F)(PFL) no preservative ¹⁰²	N/A	10 mg/mL ¹⁰²	discard unused portion ¹⁰²	1-4 mg/mL NS , D5W ¹⁰² 250-500 mL†	NS: 10 d F plus an additional 24 h RT ^{10,102} D5W: 24 h F plus an additional 12 h RT ¹⁰²	
riTUXimab intravitreal injection (RITUXAN®) 100 mg/10 mL (Roche) (F)(PFL) no preservative ¹⁰²	N/A	10 mg/mL ¹⁰²	discard unused portion ¹⁰²	syringe for intravitreal use	use within 4 h of initial puncture ¹⁰	
riTUXimab subcutaneous (RITUXAN® SC) 1400 mg/11.7 mL 1600 mg/13.4 mL (Roche) (F)(PFL) no preservative ¹⁰³	N/A	120 mg/mL ¹⁰³	discard unused portion ¹⁰³	SC syringe ¹⁰³	48 h F plus 8 h RT ¹⁰³	- contains hyaluronidase ¹⁰³ - formulations are NOT interchangeable ¹⁰³

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
riTUXimab (RIXIMYO®) 100 mg/10 mL 500 mg/50 mL (Sandoz) (F)(PFL) (do NOT shake) no preservative ¹⁰⁴	N/A	10 mg/mL ¹⁰⁴	discard unused portion ¹⁰⁴	1-4 mg/mL NS , D5W ¹⁰⁴ 250-500 mL† gently invert to mix	NS: 10 d F plus an additional 24 h RT ^{10,104} D5W: 24 h F plus an additional 12 h RT ¹⁰⁴	
riTUXimab <u>intravitreal injection</u> (RIXIMYO®) 100 mg/10 mL (Sandoz) (F)(PFL) (do NOT shake) no preservative ¹⁰⁴	N/A	10 mg/mL ¹⁰⁴	discard unused portion ¹⁰⁴	syringe for intravitreal use	use within 4 h of initial puncture ¹⁰	
riTUXimab (RUXIENCE®) 100 mg/10 mL 500 mg/50 mL (Pfizer) (F)(PFL) no preservative ¹⁰⁵	N/A	10 mg/mL ¹⁰⁵	discard unused portion ¹⁰⁵	1-4 mg/mL NS , D5W ¹⁰⁵ 250-500 mL† gently invert to mix	24 h F plus an additional 24 h RT ¹⁰⁵	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
riTUXimab <u>intravitreal injection</u> (RUXIENCE®) 100 mg/10 mL 500 mg/50 mL (Pfizer) (F)(PFL) no preservative ¹⁰⁵	N/A	10 mg/mL ¹⁰⁵	discard unused portion ¹⁰⁵	syringe for intravitreal use	use within 4 h of initial puncture ¹⁰	
riTUXimab (TRUXIMA®) 100 mg/10 mL 500 mg/50 mL (Teva/Celltrion) (F)(PFL) no preservative ¹⁰⁶	N/A	10 mg/mL ¹⁰⁶	discard unused portion ¹⁰⁶	1-4 mg/mL NS , D5W ¹⁰⁶ 250-500 mL† gently invert to mix	24 h F plus an additional 12 h RT ¹⁰⁶	
riTUXimab <u>intravitreal injection</u> (TRUXIMA®) 100 mg/10 mL 500 mg/50 mL (Teva/Celltrion) (F)(PFL) no preservative ¹⁰⁶	N/A	10 mg/mL ¹⁰⁶	discard unused portion ¹⁰⁶	syringe for intravitreal use	use within 4 h of initial puncture ¹⁰	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
romiDEPsin 10 mg (Celgene Australia) (RT) no preservative ¹⁰⁷	2.2 mL supplied diluent ¹⁰⁷ swirl to mix ¹⁰⁷	5 mg/mL ¹⁰⁷	discard unused portion ¹⁰⁷	500 mL NS ¹⁰⁷	24 h F ¹⁰⁷	- vials contain overfill to allow full drug recovery (drug vial contains 11 mg romidepsin; diluent vial has 2.4 mL diluent) ¹⁰⁷ - solvent contains 80% propylene glycol and 20% anhydrous ethanol ¹⁰⁷
Sacituzumab govitecan 180 mg (Gilead) (F)(PFL) no preservative ¹⁰⁸	20 mL NS ¹⁰⁸ bring vials to RT before reconstitution ¹⁰⁸ slowly add diluent to vial and gently swirl; allow to dissolve for up to 15 min ¹⁰⁸ do not shake ¹⁰⁸	10 mg/mL ¹⁰⁸	use immediately after reconstitution to prepare infusion solution ¹⁰⁸ discard unused portion ¹⁰⁸	1.1-3.4 mg/mL NS ¹⁰⁸ 100-1000 mL† slowly inject solution to bag to minimize foaming; do not shake ¹⁰⁸	24 h F ¹⁰⁸ , plus an additional 8 h RT including infusion time ¹⁰⁸ **(PFL) ¹⁰⁸	- do not shake ¹⁰⁸ - protect container from light during administration ¹⁰⁸ - vials contain overfill (~20 mg per vial) ¹⁰⁹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Siltuximab 100 mg 400 mg (Recordati/EUSA) (F)(PFL) no preservative ^{110,111}	100 mg: 5.2 mL SWI ¹¹⁰ 400 mg: 20 mL SWI ¹¹⁰ bring vial to RT prior to use (~30 min) ¹¹⁰ gently swirl, do NOT shake ¹¹⁰	20 mg/mL ¹¹⁰	2 h RT ¹¹⁰	250 mL D5W ¹¹⁰ dilute to final volume by withdrawing volume from bag equal to volume of drug to be added ¹¹⁰ gently mix ¹¹⁰	complete administration within 6 h RT ¹¹⁰	- administer with 0.2 micron in-line filter ¹¹⁰ - do not use if visibly opaque, discoloured, or contains particles ¹¹⁰

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Sirolimus, nanoparticle, albumin- bound (NAB) 100 mg (Aadi) (F)(PFL) no preservative ¹¹² (SAP)	20 mL NS ¹¹² slowly direct diluent against side of vial (over ≥1 min) ¹¹² let stand for ≥5 min to wet powder ¹¹² gently swirl or invert for ≥2 min to avoid foaming ¹¹² if foaming/clumping occurs, let stand until foam subsides (≥15 min) ¹¹²	5 mg/mL ¹¹²	4 h F ^{113,114} **(PFL) ¹¹²	undiluted in empty PVC or non-PVC infusion bag ¹¹²	9 h F, followed by max 4 h RT ¹¹² **(PFL) ¹¹²	- each vial contains ~800-900 mg human albumin ^{112,115} - to prevent foaming, do NOT inject NS directly onto the powder ¹¹² - if powder is visible after reconstitution, gently invert to resuspend powder ¹¹² - to prevent administration of proteinaceous strands, administer with 15 micron filter ONLY ¹¹²

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Streptozocin 1g (Keocyt) (F)(PFL) no preservative ¹¹⁶⁻¹¹⁹ (SAP)	9.5mL NS , SWI, D5W ¹¹⁶⁻¹¹⁹	100 mg/mL ¹¹⁶⁻¹¹⁹	12 h F ^{10,117-119}	syringe ¹¹⁷⁻¹¹⁹	48 h F ^{10,117-119}	
				100-500 mL NS , D5W, SWI ¹¹⁶⁻¹¹⁹	24 h F ¹¹⁷⁻¹¹⁹	
Tarlatamab 1 mg (Amgen) (F)(PFL) no preservative ¹²⁰	1.3 mL SWI do NOT use supplied IV solution stabilizer to reconstitute vials ¹²⁰ direct diluent against side of vial ¹²⁰ gently swirl to mix; do not shake ¹²⁰	0.9 mg/mL ¹²⁰	discard unused portion ¹⁰	250 mL NS ¹²⁰ add 13 mL supplied IV solution stabilizer to NS bag and gently mix to avoid foaming; do not shake ¹²⁰ add 1.1 mL reconstituted drug to IV bag following addition of IV solution stabilizer ¹²⁰ gently mix by inverting bag; do not shake ¹²⁰	complete administration within 7 d F, 8 h RT ¹²⁰	- CAUTION: two vial sizes are available; final vial concentrations are different after reconstitution - due to low extractable volume, use filtered venting needle (e.g., Chemo-Vent®) in place of CSTD for compounding ¹²¹ -vials contain overfill to allow full drug recovery ¹²⁰ - discard if cloudy or has particulates ¹²⁰

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Tarlatamab 10 mg (Amgen) (F)(PFL) no preservative ¹²⁰	4.4 mL SWI ¹²⁰ do NOT use supplied IV solution stabilizer to reconstitute vials ¹²⁰ direct diluent against side of vial ¹²⁰ gently swirl to mix; do not shake ¹²⁰	2.4 mg/mL ¹²⁰	discard unused portion ¹⁰	250 mL NS ¹²⁰ add 13 mL supplied IV solution stabilizer to NS bag and gently mix to avoid foaming; do not shake ¹²⁰ add 4.2 mL reconstituted drug to IV bag following addition of IV solution stabilizer ¹²⁰ gently mix by inverting bag; do not shake ¹²⁰	complete administration within 7 d F, 8 h RT ¹²⁰	- CAUTION: two vial sizes are available; final vial concentrations are different after reconstitution - due to low extractable volume, use filtered venting needle (e.g., Chemo-Vent®) in place of CSTD for compounding ¹²¹ -vials contain overfill to allow full drug recovery ¹²⁰ - discard if cloudy or has particulates ¹²⁰

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Tebentafusp 100 mcg/0.5 mL (Immunocore/Medison) (F)(PFL) do not shake no preservative ¹²²	N/A	200 mcg/mL ¹²²	discard unused portion ¹²²	100 mL NS ¹²² Step 1: add calculated volume of human albumin 5% to provide 225-275 mcg/mL final concentration ¹²² to mix: invert the bag and gently rotate ≥5 times; do NOT shake bag (repeat x3) ¹²² Step 2: add calculated volume of drug ¹²² to mix: invert the bag and gently rotate ≥5 times; do NOT shake bag (repeat x3) ¹²²	complete administration within 24 h F, 4 h RT ¹²² bring to RT prior to administration ¹²²	- do not use CSTD or filters during preparation ¹²² ; use filtered venting needle (e.g., Chemo-Vent®) for preparation - CSTD can be used for administration ¹²³ - administer using 0.2 micron in-line filter ¹²² - once the bag has been removed from fridge, it must remain at RT ¹²²

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Teclistamab 30 mg/3 mL 153 mg/1.7 mL (Janssen) (F)(PFL) do not shake no preservative ¹²⁴	N/A	30 mg ¹²⁴ : 10 mg/mL (use for 2.1-52.9 mg doses)*	discard unused portion ¹²⁴	SC syringe ¹²⁴ if drug volume >2 mL, divide volume into separate syringes for administration ¹²⁴	20 h F, RT ¹²⁴ if stored in fridge, bring to RT prior to administration ¹²⁴	- CAUTION: two concentrations are available ¹²⁴ - do not use CSTD for volumes less than 1 mL ¹²⁵ ; use filtered venting needle (e.g., Chemo-Vent®) for preparation ¹²⁶
		153 mg ¹²⁴ : 90 mg/mL (use for 53-375 mg doses)*				
		bring to RT before use (~15 min) ¹²⁴ swirl gently for 10 sec to mix; do NOT shake ¹²⁴				

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Temsirolimus 30 mg/1.2 mL (Pfizer/Wyeth) (F)(PFL) ¹²⁷ no preservative ¹²⁸	1.8 mL supplied diluent ¹²⁷	10 mg/mL ¹²⁷	12 h RT ^{10,127} **(PFL) ¹²⁷	250 mL NS ¹²⁷ record time of dilution ¹²⁷	complete administration within 6 h ¹²⁷ mix by gentle inversion to avoid foaming ¹²⁷	- use non-DEHP bag and tubing - administer with 0.2 micron in-line filter ¹²⁷
Teniposide 50 mg/5 mL (BMS) (RT) preservative ¹²⁹	N/A	10 mg/mL ¹²⁹	discard unused portion	0.1-1 mg/mL NS , D5W ¹²⁹ 50–500 mL *	0.1-0.4 mg/mL: 24 h RT ¹²⁹ 1 mg/mL: complete administration within 4 h RT of preparation ^{129,130}	- do not refrigerate - use non-DEHP bag and tubing ¹²⁹ - do not use if precipitates ^{129,130} - contains DMA*** - excessive agitation may cause precipitation ¹²⁹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Thiotepa 15 mg 100 mg (Hikma) (F, PFL) no preservative ¹³¹	15 mg ¹³¹ : 1.5 mL SWI 100 mg ¹³¹ : 10 mL SWI to remove haze, filter through 0.22 micron filter disc after reconstitution ¹³² record time of reconstitution	10 mg/mL ¹³¹	8 h F ¹³¹	0.5-1 mg/mL NS ¹³¹ ≤500 mg ¹³¹ : 500 mL >500 mg ¹³¹ : 1000 mL reconstituted solution is hypotonic and must be further diluted with NS prior to use ¹³¹	24 h F, 4 h RT ¹³¹	- discard if precipitates are present ¹³¹ - reconstituted solution may be used if opalescent ¹³¹ - administer with 0.2 micron in-line filter ¹³¹ - use non-PVC bag and administration set ¹³¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Thiotepa IT injection 15 mg 100mg (Hikma) (F, PFL) no preservative ¹³¹	15 mg ¹³¹: 1.5 mL SWI 100 mg ¹³¹: 10 mL SWI diluents containing preservatives should NOT be used for intrathecal administration ²⁵ to remove haze, filter through 0.22 micron filter disc after reconstitution ¹³² record time of reconstitution	10 mg/mL ¹³¹	8 h F ¹³¹	IT syringe qs to 6 mL with preservative free NS ¹³³ diluents containing preservatives should NOT be used for intrathecal administration ²⁵	use within 4 h of initial reconstitution ¹⁰	- auxiliary info ¹³⁴ : IT - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag ¹³⁴ - discard if precipitates are present ¹³¹ - reconstituted solution may be used if opalescent ¹³¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Thiotepa 15 mg 100 mg (SteriMax) (F, PFL) no preservative ¹³⁵	15 mg ¹³⁵ : 1.5 mL SWI 100 mg ¹³⁵ : 10 mL SWI to remove haze, filter through 0.22 micron filter disc after reconstitution ¹³² record time of reconstitution	10 mg/mL ¹³⁵	8 h F ¹³⁵	0.5-1 mg/mL NS ¹³⁵ ≤500 mg ¹³⁵ : 500 mL >500 mg ¹³⁵ : 1000 mL reconstituted solution is hypotonic and must be further diluted with NS prior to use ¹³⁵	24 h F, 4 h RT ¹³⁵	- discard if precipitates are present ¹³⁵ - reconstituted solution may be used if opalescent ¹³⁵ - administer with 0.2 micron in-line filter ¹³⁵ - use non-PVC bag and administration set ¹³⁵

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Thiotepa IT injection 15 mg 100mg (SteriMax) (F, PFL) no preservative ¹³⁵	15 mg ¹³⁵ : 1.5 mL SWI 100 mg ¹³⁵ : 10 mL SWI diluents containing preservatives should NOT be used for intrathecal administration ²⁵ to remove haze, filter through 0.22 micron filter disc after reconstitution ¹³² record time of reconstitution	10 mg/mL ¹³⁵	8 h F ¹³⁵	IT syringe qs to 6 mL with preservative free NS ¹³³ diluents containing preservatives should NOT be used for intrathecal administration ²⁵	use within 4 h of initial reconstitution ¹⁰	- auxiliary info ¹³⁴ : IT - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag ¹³⁴ - discard if precipitates are present ¹³⁵ - reconstituted solution may be used if opalescent ¹³⁵
Thyrotropin alfa 1.1 mg (Genzyme) (F)(PFL) no preservative ¹³⁶	1.2 mL SWI ¹³⁶ swirl gently to mix ¹³⁶ do NOT shake ¹³⁶	0.9 mg/mL ¹³⁶	12 h F ^{10,136}	syringe ¹³⁶	24 h F ^{10,136}	- do not use if particulates are present ¹³⁶

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Tislelizumab 100 mg/10 mL (BeiGene) (F)(PFL) (do not shake) no preservative ¹³⁷ (SAP)	N/A	10 mg/mL ¹³⁷	discard unused portion ¹³⁷	2-5 mg/mL NS ¹³⁷ 50 mL* mix by gentle inversion; do not shake ¹³⁷	complete administration within 20 h F, 4 h RT (max 24 h from preparation) ¹³⁷ bring to RT prior to administration ¹³⁷	- discard if has visible particulates, or is discoloured or cloudy ¹³⁷ - administer with 0.2 micron in-line filter ¹³⁷
Tocilizumab 80 mg/4 mL 200 mg/10 mL 400 mg/20 mL (Roche) (F)(PFL) no preservative ¹³⁸	N/A	20 mg/mL ¹³⁸	discard unused portion ¹³⁸	100 mL NS ¹³⁸ dilute to final volume by withdrawing volume from bag equal to volume of drug to be added ¹³⁸ gently invert to mix ¹³⁸	complete administration within 24 h F, RT ¹³⁸ bring to RT prior to administration ¹³⁸	- to prevent foaming: slowly add drug to infusion bag and gently invert bag to mix ¹³⁸
Topotecan 4 mg/4 mL (Accord) (RT)(PFL) no preservative ¹³⁹	N/A	1 mg/mL ¹³⁹	12 h F, RT ^{10,139}	0.025-0.5 mg/mL NS, D5W ¹³⁹ 25-50 mL†	10 d F, 4 d RT ^{10,139}	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Topotecan IT injection 4 mg/4 mL (Accord) (RT)(PFL) no preservative ¹³⁹	N/A	1 mg/mL ¹³⁹	use within 4 h of initial puncture ¹⁰	IT syringe qs to 10 mL with preservative free NS ^{26,140,141} diluent containing preservatives should NOT be used for intrathecal administration ²⁵	use within 4 h of initial puncture ¹⁰	- auxiliary info ¹⁰ : IT - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag ²⁶
Topotecan 4 mg/4 mL (Pfizer/Hospira) (F)(PFL) no preservative ¹⁴²	N/A	1 mg/mL ¹⁴²	discard unused portion ¹⁴²	0.02-0.5 mg/mL NS , D5W ¹⁴² 25-50 mL†	24 h F, RT ¹⁴²	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Topotecan IT injection 4 mg/4 mL (Pfizer/Hospira) (F)(PFL) no preservative ¹⁴²	N/A	1 mg/mL ¹⁴²	use within 4 h of initial puncture ¹⁰	IT syringe qs to 10 mL with preservative free NS ^{26,140,141} diluent containing preservatives should NOT be used for intrathecal administration ²⁵	use within 4 h of initial puncture ¹⁰	- auxiliary info ¹⁰ : IT - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag ²⁶
Topotecan 4 mg/4 mL (Sandoz) (F)(PFL) no preservative ¹⁴³	N/A	1 mg/mL ¹⁴³	discard unused portion ¹⁴³	0.02-0.5 mg/mL NS , D5W ¹⁴³ 25-50 mL†	24 h F ¹⁴³ **(PFL) ¹⁴³	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Topotecan IT injection 4 mg/4 mL (Sandoz) (F)(PFL) no preservative ¹⁴³	N/A	1 mg/mL ¹⁴³	use within 4 h of initial puncture ¹⁰	IT syringe qs to 10 mL with preservative free NS ^{26,140,141} diluent containing preservatives should NOT be used for intrathecal administration ²⁵	use within 4 h of initial puncture ¹⁰ **(PFL) ¹⁴³	- auxiliary info ¹⁰ : IT - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag ²⁶
Trastuzumab (HERCEPTIN®) 440 mg (Roche) (F) no preservative ¹⁴⁴	20 mL supplied BWI ¹⁴⁴ swirl vial gently; allow to stand undisturbed for 5 min ¹⁴⁴	21 mg/mL ¹⁴⁴	28 d F ¹⁴⁴	250 mL NS only ¹⁴⁴ do NOT use dextrose containing solutions ¹⁴⁴	24 h F, RT ¹⁴⁴	- do NOT shake ¹⁴⁴

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Trastuzumab (HERZUMA®) 150 mg 440 mg (Teva/Celltrion) (F) no preservative ¹⁴⁵	150 mg: 7.2 mL SWI ¹⁴⁵	21 mg/mL ¹⁴⁵	discard unused portion ¹⁴⁵	250 mL NS only ¹⁴⁵ do NOT use dextrose containing solutions ¹⁴⁵	24 h F, RT ¹⁴⁵	- do NOT shake ¹⁴⁵ - supplied BWI contains benzyl alcohol ¹⁴⁵
	440 mg: 20 mL supplied BWI ¹⁴⁵		28 d F ¹⁴⁵			
	swirl vial gently; allow to stand undisturbed for 5 min ¹⁴⁵					
Trastuzumab (OGIVRI®) 150 mg 440 mg (BGP) (F) no preservative ¹⁴⁶	150 mg: 7.2 mL SWI ¹⁴⁶	21 mg/mL ¹⁴⁶	discard unused portion ¹⁴⁶	250 mL NS only ¹⁴⁶ do NOT use dextrose containing solutions ¹⁴⁶	24 h F, RT ¹⁴⁶	- do NOT shake ¹⁴⁶ - supplied BWI contains benzyl alcohol ¹⁴⁶
	440 mg: 20 mL supplied BWI ¹⁴⁶		28 d F ¹⁴⁶			
	swirl vial gently; allow to stand undisturbed for 5 min ¹⁴⁶					

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Trastuzumab (TRAZIMERA®) 150 mg 440 mg (Pfizer) (F) no preservative ¹⁴⁷	150 mg: 7.2 mL SWI ¹⁴⁷	21 mg/mL ¹⁴⁷	discard unused portion ¹⁴⁷	250 mL NS only ¹⁴⁷ do NOT use dextrose containing solutions ¹⁴⁷	24 h F, RT ¹⁴⁷	- do NOT shake ¹⁴⁷ - supplied BWI contains benzyl alcohol ¹⁴⁷
	440 mg: 20 mL supplied BWI ¹⁴⁷		28 d F ¹⁴⁷			
	swirl vial gently; allow to stand undisturbed for 5 min ¹⁴⁷					
Trastuzumab deruxtecan (ENHERTU®) 100 mg (AstraZeneca) (F)(PFL) no preservative ¹⁴⁸	5 mL SWI ¹⁴⁸ swirl gently until completely dissolved ¹⁴⁸ do NOT shake ¹⁴⁸	20 mg/mL ¹⁴⁸	12 h F ^{10,148} **(PFL) ¹⁴⁸	100 mL D5W only ¹⁴⁸ gently invert to mix ¹⁴⁸ do NOT shake ¹⁴⁸ do NOT use sodium chloride solution ¹⁴⁸	complete administration within 24 h F, 4 h RT ¹⁴⁸ **(PFL) ¹⁴⁸	- do not use if reconstituted solution contains visible particulates or is cloudy or discoloured ¹⁴⁸ - protect container from light during administration ¹⁴⁹ - administer with 0.2 micron in-line filter ¹⁴⁸ - if stored in fridge, bring bag to RT prior to use ¹⁴⁸

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Trastuzumab emtansine (KADCYLA®) 100 mg 160 mg (Roche) (F)(PFL) no preservative ¹⁵⁰	100 mg: 5 mL SWI ¹⁵⁰ 160 mg: 8 mL SWI ¹⁵⁰ swirl gently until completely dissolved do NOT shake ¹⁵⁰	20 mg/mL ¹⁵⁰	12 h F ^{10,151}	250 mL NS or ½NS only ¹⁵⁰ do NOT shake ¹⁵⁰ do NOT use dextrose containing solutions ¹⁵⁰	24 h F ¹⁵⁰	- do not use if reconstituted solution contains visible particulates or is cloudy or discolored ¹⁵⁰ - D5W causes aggregation of the protein ¹⁵⁰ - for infusions prepared in NS: administer with 0.2 micron in-line filter or 0.22 micron polyethersulfane (PES) filter ¹⁵⁰ - for infusions prepared in ½NS: filter is optional for administration ¹⁵⁰

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Tremelimumab 25 mg/1.25 mL 300 mg/15 mL (AstraZeneca) (F)(PFL) (do not shake) no preservative ¹⁵²	N/A	20 mg/mL ¹⁵²	discard unused portion ¹⁵²	0.1-10 mg/mL NS , D5W ¹⁵² 50 mL* mix by gentle inversion; do NOT shake ¹⁵²	24 h F , RT ¹⁵²	- administer with 0.2 micron in-line filter ¹⁵² - discard if visible particles are present ¹⁵²

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Treosulfan 1 g 5 g (Medexus) (RT) no preservative ¹⁵³	1 g ¹⁵³ : 20 mL NS, D5W, SWI, ½NS 5 g ¹⁵³ : 100 mL NS, D5W, SWI, ½NS pre-heat diluent to 25-30°C (max) ¹⁵⁴ shake vial to loosen powder before adding the warmed diluent ¹⁵⁵ vigorous shaking may be required ¹⁵⁵ ; prolonged standing time may improve solubility ¹⁵³	50 mg/mL ¹⁵³	12 h RT ^{10,153}	undiluted in empty infusion bag ^{153,154}	3 d RT ¹⁵³	- do NOT refrigerate as may precipitate ¹⁵³

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Treosulfan 1 g 5 g (medac) (RT) no preservative ^{156,157} (SAP)	1 g ^{156,157} : 20 mL SWI, ½NS 5 g ^{156,157} : 100 mL SWI, ½NS pre-heat diluent to 25-30°C (max) ^{156,157} shake vial carefully to loosen powder before adding the warmed diluent ^{156,157} gently shake while adding diluent ^{156,157} (takes ~2 min to reconstitute) ^{156,157}	50 mg/mL ^{156,157}	12 h RT ^{10,156,158}	undiluted ¹⁵⁹ or dilute with NS or D5W in empty infusion bag to final concentration of 20 mg/mL ¹⁵⁸	4 d RT ^{156,158}	- compatible with polytetrafluoroethyl ene filters ¹⁶⁰ - may sometimes require vigorous shaking to reconstitute ^{156,157} - do NOT refrigerate as may cause precipitation ^{156,157}
vinBLASTine 10 mg/10 mL (Pfizer) (F)(PFL) no preservative ¹⁶¹	N/A	1 mg/mL ¹⁶¹	discard unused portion ^{2,161}	25-50 mL NS , D5W ¹⁶²	use within 4 h of initial vial puncture ^{2,161}	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{163,164}

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
vinBLASTine 10 mg/10 mL (Teva) (F)(PFL) no preservative ¹⁶⁵	N/A	1 mg/mL ¹⁶⁵	discard unused portion ^{2,165}	25-50 mL NS , D5W ¹⁶²	use within 4 h of initial vial puncture ^{2,165}	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{163,164}
vinCRISTine 2 mg/2 mL 5 mg/5 mL (Pfizer/Hospira) (F)(PFL) no preservative ¹⁶⁶	N/A	1 mg/mL ¹⁶⁶	8 h F, RT ¹⁶⁶	0.01-0.1 mg/mL NS , D5W ¹⁶⁶ 50 mL†	24 h F, RT ¹⁶⁶ **(PFL) ¹⁶⁶	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{163,164} - for LYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing either etoposide or etoposide phosphate AND DOXOrubicin and vinCRISTine)

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
vinCRISTine 1 mg/1 mL 2 mg/2 mL 5 mg/5 mL (Teva) (F)(PFL) no preservative ¹⁶⁷	N/A	1 mg/mL ¹⁶⁷	8 h F, RT ¹⁶⁷	0.01-0.1 mg/mL NS , D5W ¹⁶⁷ 50 mL†	24 h F, RT ¹⁶⁷	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{163,164} - for LYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing either etoposide or etoposide phosphate AND DOXOrubicin and vinCRISTine)
Vinorelbine 10 mg/1 mL 50 mg/5mL (Fresenius Kabi) (F)(PFL) no preservative ¹⁶⁸	N/A	10 mg/mL ¹⁶⁸	discard unused portion ¹⁶⁸	0.5-2.0 mg/mL NS , D5W, ½NS, D5-½NS, LR, Ringer's ¹⁶⁸ 50 mL†	24 h F, RT ¹⁶⁸	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{163,164}

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Vinorelbine 10 mg/1 mL 50 mg/5 mL (GMP) (F)(PFL) no preservative ¹⁶⁹	N/A	10 mg/mL ¹⁶⁹	discard unused portion ²	0.5-2.0 mg/mL NS , D5W, ½NS, D5-½NS, LR, Ringer's ¹⁶⁹ 50 mL†	24 h F , RT ¹⁶⁹	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{163,164}
Vinorelbine 10 mg/1 mL 50 mg/5 mL (Teva) (F)(PFL) no preservative ¹⁷⁰	N/A	10 mg/mL ¹⁷⁰	discard unused portion ¹⁷⁰	0.5–2.0 mg/mL NS , D5W, ½NS, D5-½NS, LR, Ringer's ¹⁷⁰ 50 mL†	24 h F , RT ¹⁷⁰	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{163,164}
Zoledronic acid 4 mg/5 mL (Dr Reddy's) (RT) no preservative ¹⁷¹	N/A	0.8 mg/mL ¹⁷¹	discard unused portion ¹⁷¹	100 mL NS , D5W ¹⁷¹	complete infusion within 24 h of preparation ¹⁷¹ refrigerate diluted product if not used immediately after preparation; bring to RT prior to use ¹⁷¹	- do NOT mix with calcium containing solutions (e.g., LR) ¹⁷¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Zoledronic acid 4 mg/5 mL (Marcan) (RT) no preservative ¹⁷²	N/A	0.8 mg/mL ¹⁷²	discard unused portion ¹⁷²	100 mL NS , D5W ¹⁷²	complete infusion within 24 h of preparation ¹⁷² refrigerate diluted product if not used immediately after preparation; bring to RT prior to use ¹⁷²	- do NOT mix with calcium containing solutions (e.g., LR) ¹⁷²
Zoledronic acid 4 mg/5 mL (MDA) (RT) no preservative ¹⁷³	N/A	0.8 mg/mL ¹⁷³	discard unused portion ¹⁷³	100 mL NS , D5W ¹⁷³	complete infusion within 24 h of preparation ¹⁷³ refrigerate diluted product if not used immediately after preparation; bring to RT prior to use ¹⁷³	- do NOT mix with calcium containing solutions (e.g., LR) ¹⁷³

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Zoledronic acid (ZOMETA) 4 mg/ 5 mL (Novartis) (RT) no preservative ¹⁷⁴	N/A	0.8 mg/mL ¹⁷⁴	discard unused portion ⁴⁵	100 mL NS , D5W ¹⁷⁴	complete infusion within 24 h of preparation ¹⁷⁴ refrigerate diluted product if not used immediately after preparation; bring to RT prior to use ¹⁷⁴	- do NOT mix with calcium containing solutions (e.g., LR) ¹⁷⁴
Zoledronic acid 4 mg/5 mL (Sandoz) (RT) no preservative ¹⁷⁵	N/A	0.8 mg/mL ¹⁷⁵	discard unused portion ¹⁷⁵	100 ml NS , D5W ¹⁷⁵	complete infusion within 24 h of preparation ¹⁷⁵ refrigerate diluted product if not used immediately after preparation; bring to RT prior to use ¹⁷⁵	- do NOT mix with calcium containing solutions (e.g., LR) ¹⁷⁵

* Suggested volume based on usual dose range and any concentration range of stability data

† see [BC Cancer IV Bag Selection table](#); standardized bag sizes are provided for select Benefit Drugs with concentration-dependent stability or large drug volume

** Protect from light means minimizing exposure to direct sunlight over a *storage* period. More specific information on protection from light (eg, protecting container and tubing during *administration*) will be indicated in the Special Precautions/Notes column.

*** Contains DMA (N,N dimethylacetamide). Product may be incompatible with closed system transfer devices (CSTD) such as ChemoLock.

Centres are not to change content locally. All suggestions for change are to be forwarded to the Cancer Drug Manual editor.

Explanatory Notes:

Stability data assumes products prepared using standard aseptic technique in biological safety cabinet at low risk for contamination according to the classification outlined in USP 797. ^{176,177}

Vial stability: Stability of solution after first puncture or reconstituted solution.

Storage temperature: If information states same stability with refrigerator and room temperature storage, then fridge stability is bolded as preferred (ie, to minimize growth of micro-organisms).

Discard unused portion: Unused portion from single use vials should be discarded at the end of the day.

“overfill known” is stated if the manufacturer states overfill that is present is within acceptable limits.

“Complete administration within ___” is stated if the manufacturer specifies that the infusion must be completed in a specific time frame following preparation, usually including entire time required for preparation (from first puncture), storage, and administration of infusion.

Nomenclature for **in-line filters** has been standardized to 0.2 micron filter size. For more information, refer to CDM monograph.

Abbreviations:

BWI = bacteriostatic water for injection

CIVI: ambulatory pump = Continuous Intravenous Infusion (e.g., elastomeric infusor)

CSTD = closed system transfer device

D5W = dextrose 5% in water

DMA = N,N dimethylacetamide

F = refrigerate

LR = lactated ringer's solution

non-DEHP = not containing Di(2-ethylhexyl) phthalate (DEHP)

non-PVC = not containing polyvinylchloride (PVC)

NS = normal saline

PFL = protect from light

RT = room temperature

SAP = drug is approved for use through the Health Canada Special Access Program

SWI = sterile water for injection

References:

1. Generic Medical Partners Inc. Leucovorin calcium injection product monograph. Toronto, Ontario; 6 March, 2020.
2. BC Cancer. Provincial Pharmacy Directive Number II-20: Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer; December 5, 2018.
3. Pfizer Canada Inc. Leucovorin calcium injection product monograph. Kirkland, Quebec; 21 June, 2018.
4. Teva Canada Limited. Leucovorin calcium injection® product monograph. Toronto, Ontario; 5 May, 2014.
5. Hospira Healthcare Corporation. LEUCOVORIN CALCIUM INJECTION® product monograph. Saint-Laurent, Quebec; 7 June, 2007.
6. Novopharm Limited (Teva). LEUCOVORIN CALCIUM® Injection product information package. Toronto, Ontario; undated,

7. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 6 January , 2006.
8. Jenny Yeung. Medical Information Specialist, Teva Canada. Personal communication. 12 April, 2017.
9. Jazz Pharmaceuticals Canada Incorporated. ZEPZELCA® product monograph. Mississauga, Ontario; September 29, 2021.
10. BC Cancer. Provincial Pharmacy Directive Number II-20: Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer; May 1 , 2022.
11. Linda Ewing.BD US Market Manager. WW Infusion Disposables. Personal communication. Feb, 2018.
12. Pharma Mar. Lurbinectedin Compassionate Use: Preparation Guide for Infusion - edition 1. Madrid, Spain; April, 2019.
13. Marcan Pharmaceuticals Inc. Melphalan for injection product monograph. Ottawa, ON; July 26, 2019.
14. Taro Pharmaceuticals Inc. Taro-Melphalan product monograph. Brampton, Ontario; April 5, 2019.
15. Baxter Corporation. UROMITEXAN® product monograph. Mississauga, Ontario; 6 August, 2013.
16. Mona Ghobros BPharm MSc. Medical Information, Baxter Corporation. Personal communication. 29 November, 2018.
17. Trissel's® 2 Clinical Pharmaceutics Database (database on the Internet). Mesna. Lexi-Comp Inc.; created by Lawrence A. Trissel; Accessed 29 November, 2018. Available at: <http://online.lexi.com>
18. Fresenius Kabi Canada Ltd. Mesna for injection product monograph. Richmond Hill, Ontario; 21 December, 2017.
19. Fresenius Kabi Canada Ltd. Mesna for injection product monograph. Richmond Hill, Ontario; 30 March, 2015.
20. Accord Healthcare Inc. Methotrexate injection product monograph. Kirkland, Québec; October 30, 2019.
21. Maia M, Farah ME, Belfort RN, et al. Effects of intravitreal triamcinolone acetonide injection with and without preservative. Br J Ophthalmol ; 2007;91(9):1122–1124
22. Mayne Pharma Canada. Methotrexate Product Monograph. Montreal, Quebec; December, 2003.
23. BC Cancer Miscellaneous Origin Tumour Group. (MOIT) BC Cancer Protocol Summary for Solid Tumours using Intrathecal Methotrexate and/or Thiotepa and/or Cytarabine. Vancouver, British Columbia: BC Cancer; September 1 , 2021.
24. BC Cancer Lymphoma Tumour Group. (LYIT) BC Cancer Protocol Summary for Treatment of Lymphoma using Intrathecal Methotrexate and Cytarabine. Vancouver, British Columbia: BC Cancer; April 1 , 2022.
25. Hematology/Oncology Pharmacy Association. HOPA News Clinical Pearls: Intrathecal Chemotherapy: Focus on Drugs, Dosing, and Preparation. 13(4) ed. Chicago, Illinois, USA: Hematology/Oncology Pharmacy Association; 2016.
26. BC Cancer. Systemic Therapy Policy and Procedure III-50: Administration of High Alert Medications by the Intrathecal Route via Lumbar Puncture or Ommaya Reservoir. Vancouver, British Columbia; May 1, 2019.
27. Fresenius Kabi. Methotrexate for injection (lyophilized powder) package insert. Lake Zurich, Illinois, USA; May, 2020.
28. Ng T, Woo S, Badry N, et al. SBAR: Methotrexate Backorder: Concentration Switch Management and Review. Approved by BC Cancer Provincial Pharmacy Professional Practice Committee (P4C). October 24, 2024
29. Pfizer Canada-ULC. Methotrexate injection product monograph. Kirkland, Québec; July 8, 2019.
30. AbbVie Deutschland GmbH & Co. KG. ELAHERE® Summary of Product Characteristics. Ludwigshafen, Germany; April 18th, 2024.
31. Accord Healthcare Inc. Mitomycin product monograph. Kirkland, Quebec; 16 July, 2018.
32. Au JLS, Badalament RA, Wientjes MG, et al. Methods to improve efficacy of intravesical mitomycin C: results of a randomized phase III trial. J Natl Cancer Inst ; April 18, 2001;93(8):597–604
33. Jessie LS Au PharmD PhD. Professor, Ohio State University. Personal communication. 14 May, 2007.
34. Myers AL, Zhang Y, Kawedia JD, et al. Solubilization and stability of mitomycin C solutions prepared for intravesical administration. Drugs R D ; 2017;17:297–304
35. Beijnen JH, Van Gijn R, Underberg WJM. Chemical stability of the antitumor drug mitomycin C in solutions for intravesical instillation. Journal of Parenteral Science and Technology ; 1990;44(6):332–335
36. Hikma Canada Limited. Mitomycin for injection product monograph. Mississauga, Ontario; October 12, 2022.
37. Teva Canada Limited. Mitomycin for injection® product monograph. Toronto, Ontario; 30 June, 2017.
38. Fresenius Kabi Canada Ltd. Mitoxantrone injection® product monograph. Richmond Hill, Ontario; September 28, 2016.
39. Pfizer Canada Inc. Mitoxantrone injection product monograph. Kirkland, Quebec; 17 October, 2018.
40. Kyowa Kirin (Distributed by Innomar Strategies). POTELIGEO® product monograph. Oakville, Ontario; July 12, 2023.
41. Sandoz Canada Inc. ATRIANCE® product monograph . Boucherville, Quebec; November 30, 2022.
42. Bristol-Myers Squibb Canada Co. OPDIVO® product monograph. Montreal, Canada; February 28, 2022.
43. Bristol-Myers Squibb Canada. OPDUALAG® product monograph. Montreal, Quebec; 13 September, 2023.
44. Hoffmann-La Roche Ltd. GAZYVA® product monograph. Mississauga, Ontario; 21 December , 2015.

45. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 19 September , 2007.
46. Anna Sivojelezova MSc. Drug Information Associate, Hoffmann-La Roche Ltd. Personal communication. 24 April, 2015.
47. Omega Laboratories Limited. OCTREOTIDE ACETATE solution product monograph. Montreal, Quebec; January 25, 2022.
48. Novartis Pharmaceuticals Canada Inc. SANDOSTATIN® and SANDOSTATIN® LAR® product monograph. Dorval, Quebec; April 19, 2021.
49. Teva Canada Limited. OCTREOTIDE for injectable suspension Product Monograph. Toronto, Ontario; Aug 30, 2021.
50. Dr. Reddy's Laboratories Canada Inc. Oxaliplatin injection product monograph. Mississauga, Ontario; February 8, 2022.
51. Pfizer Canada Inc. Oxaliplatin injection product monograph. Kirkland, Quebec; 31 May, 2017.
52. Medical Information Pfizer Canada Inc. Personal communication. 6 June, 2017.
53. Sandoz Canada Inc. Oxaliplatin injection product monograph. Boucherville, Quebec; 12 August, 2015.
54. Katryn Vosburg. Drug Information & Pharmacovigilance Specialist, Sandoz Canada Inc. Personal communication. 26 February, 2016.
55. Teva Canada Limited. Teva-Oxaliplatin injection® product monograph. Toronto, Ontario; 11 September, 2015.
56. Accord Healthcare Inc. Paclitaxel Injection. Kirkland, Quebec; September 14, 2020.
57. Mercure C. Stability of 0.1 mg/mL of paclitaxel for injection in sodium chloride (0.9%) solution. St Catharines, Ontario: Biolyse Pharma; 2 February , 2007.
58. Biolyse. PACLITAXEL FOR INJECTION® product monograph. St. Catharines, Ontario; 12 September, 2013.
59. Claude Mercure. Manager BPC. Personal communication. 24 June, 2014.
60. Zeng Z, Lazakovitch E. Study IR 120: Physical and Chemical Stability Study of Paclitaxel for Injection in 0.9 % Sodium Chloride in concentration range 0.012-0.12 mg/mL. Biolyse Pharma ; March, 2010
61. Biolyse. PACLITAXEL FOR INJECTION® product monograph. St. Catharines, Ontario; 2 December , 2005.
62. Xu Q, Trissel LA, Martinez JF. Pharmaceutical Analysis Laboratory, University of Texas M. D. Anderson Cancer Center, Houston 77030. Stability of paclitaxel in 5% dextrose injection or 0.9% sodium chloride injection at 4, 22, or 32 degrees C. Am J Hosp Pharm ; Dec 15, 1994;51(24):3058–60
63. Lisa Tavano. Biolyse Pharma Corporation. Personal communication. 14 May, 2012.
64. Sandoz Canada Inc. Paclitaxel injection USP product monograph . Boucherville, Quebec; October 18, 2021.
65. Bristol-Myers Squibb Canada. ABRAXANE® product monograph. Saint-Laurent, Quebec; March 8, 2024.
66. Aisling Cahill. Drug Safety and Medical Information Specialist. Celgene Inc. Personal communication. April 23, 2015.
67. Panacea Biotec Pharma Limited (distributed by Apotex Inc.). Nanoparticle, albumin-bound paclitaxel product monograph. Toronto, Ontario; November 16, 2021.
68. Panacea Biotec Pharma Limited (R&D L. In-use infusion stability study report for paclitaxel powder for injectable suspension nanoparticle, albumin-bound paclitaxel (100 mg paclitaxel/vial). Toronto, Ontario; October 4, 2023.
69. Pharmaceutical Partners of Canada. Pamidronate Disodium For Injection product monograph. Richmond Hill, Ontario; 18 January , 2010.
70. Mayne Pharma (Canada) Inc. Pamidronate Package Insert. Montreal, Quebec; 2002.
71. Omega Laboratories Ltd. Pamidronate Disodium product monograph. Montreal, Quebec; 06 June , 2005.
72. Pfizer Canada-ULC. Pamidronate disodium for injection product monograph. Kirkland, Quebec; December 11, 2018.
73. Sandoz Canada Inc. Pamidronate injection product monograph. Boucherville, Quebec; 28 February, 2006.
74. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 19 September , 2007.
75. Amgen Canada. VECTIBIX® product monograph. Mississauga, Ontario; 31 March, 2017.
76. Amgen Inc. VECTIBIX® full prescribing information. Thousand Oaks, California USA; June, 2017.
77. Amgen Canada. VECTIBIX® product monograph. Mississauga, Ontario; 5 March , 2009.
78. Diane Lord. Medical Information Department, Amgen Canada Inc. Personal communication. 19 June, 2009.
79. Servier Canada Inc. ONCASPAR® product monograph. Laval, Quebec; November 22, 2021.
80. Servier Canada Inc. Personal Communication: Availability of US-Labelled ONCASPAR® (pegaspargase injection) for Canadian patients. January 27, 2025.
81. Merck Canada Inc. KEYTRUDA® product monograph. Kirkland, Quebec; 6 December, 2019.
82. Accord Healthcare Inc. Pemetrexed disodium for injection product monograph. Kirkland, Quebec; 12 March, 2015.
83. Accord Healthcare Inc. Pemetrexed solution for injection product monograph. Kirkland, QC; April 26, 2021.
84. Dr. Reddy's Laboratories Limited. Pemetrexed for Injection product monograph . Oakville, Ontario; April 12, 2017.
85. Dr. Reddy's Laboratories Ltd. Pemetrexed for Injection product monograph. Mississauga, Ontario; May 3, 2022.

86. Dr. Reddy's Laboratories Inc. Pemetrexed for injection full prescribing information (package insert). Princeton, NJ, USA; Mar, 2022.
87. Md Aslam. Medical information Associate. Drug information. Dr Reddy's Laboratories. Personal Communication (verbal). December 1, 2022.
88. Eli Lilly and Company. ALIMTA® product monograph. Indianapolis, Indiana, USA; September, 2013.
89. Taro Pharmaceuticals Inc. Pemetrexed disodium for injection product monograph. Brampton, Ontario; 30 August, 2018.
90. Hospira Inc. NIPENT® full prescribing information. Lake Forest, Illinois USA; October, 2019.
91. Hoffmann-La Roche Limited. PERJETA® product monograph. Mississauga, Ontario; April 12, 2013.
92. Hoffmann-La Roche Limited. PHESGO® product monograph. Mississauga, Ontario; January 5, 2022.
93. sanofi-aventis Canada Inc. MOZOBIL® product monograph. Laval, Quebec; 8 October , 2014.
94. Maureen Coughlin BSc Pharm. Solutions in Health Inc. (acting as an authorized agent of sanofi-aventis and). Personal communication. 24 May, 2017.
95. Hoffman-La Roche Limited. POLIVY® product monograph. Mississauga, Ontario; April 27, 2021.
96. Servier Canada Inc. FOLOTYN® product monograph. Laval, Quebec; 19 October, 2018.
97. Marjolaine Migeon PharmD. Servier Canada Medical Information. Personal communication. 24 September, 2019.
98. Pfizer Canada Inc. TOMUDEX® product monograph. Kirkland, Quebec; 8 August, 2017.
99. Eli Lilly Canada Inc. CYRAMZA® product monograph. Toronto, Ontario; 16 July, 2015.
100. Marilyn Bain BScN. Senior Medical Information Associate, Eli Lilly Canada Inc. Personal communication. 16 January, 2017.
101. Innomar Strategies Inc for Incyte Corporation. ZYNYZ® product monograph (undated). Oakville, Ontario;
102. Hoffmann-La Roche Ltd. RITUXAN® product monograph. Mississauga, Ontario; October 10, 2019.
103. Hoffmann-La Roche Ltd. RITUXAN® SC product monograph. Mississauga, Ontario; March 21, 2018.
104. Sandoz Canada Inc. RIXIMYO® product monograph. Boucherville, Quebec; April 28, 2020.
105. Pfizer Canada-ULC. RUXIENCE® product monograph. Kirkland, Quebec; May 4, 2020.
106. Teva Canada Limited for Celltrion Healthcare Co Ltd. TRUXIMA® product monograph. Toronto, Ontario; July 22, 2019.
107. Celgene Pty Limited. ISTODAX® Australian product information. Mulgrave, Victoria, Australia; February 1, 2022.
108. Gilead Sciences Canada Inc. TRODELVY® product monograph. Mississauga, ON; August 2, 2022.
109. Gilead Medical Information. Gilead Sciences Inc. Personal Communication: Trodelvy® (sacituzumab govitecan) Concentration After Reconstitution. Jan 24, 2022.
110. Recordati Rare Diseases Canada Inc. SYLVANT® product monograph. Toronto, Ontario; July 30, 2024.
111. Recordati Industria Chimica E Farmaceutica (Milan, Italy). Personal Communication: News Release - Recordati completes the acquisition of EUSA Pharma (UK). March 16, 2022.
112. Aadi Bioscience Inc. Pharmacy Manual Protocol PEX-002 (Version 4.0) Expanded Access for an Intermediate-size Population for ABI-009 (Sunitinib Albumin-bound Nanoparticles for Injectable Suspension) in Patients with Perivascular Epithelioid Cell Tumors (PEComa) or Patients with Relevant Genetic Mutations for mTOR Pathway Activation. Pacific Palisades, CA, USA; July 16, 2021.
113. Aadi Bioscience Inc. ABI-009 sunitinib albumin-bound nanoparticle (packaging information - BOX). Pacific Palisades, CA; Feb 1 , 2022.
114. Aadi Bioscience Inc. ABI-009 sunitinib albumin-bound nanoparticle (packaging information - VIAL). Pacific Palisades, CA; Feb 1, 2022.
115. Aadi Bioscience Inc. ABI-009 (*nab*-Sunitinib) Investigator's Brochure - version 10.0. Pacific Palisades, CA, USA; July 27, 2021.
116. Keocyt. Streptozocin Keocyt summary of product characteristics, version 5. Montrouge, France; February, 2016.
117. Keocyt. ZANOSAR® package leaflet. Montrouge, France; 24 February, 2016.
118. Keocyt. ZANOSAR® summary of product characteristics, version 3. Montrouge, France; 31 October, 2017.
119. Keocyt-Riemsers Pharma. Marie-Laure Vedel, Sales Order Handling and Hospital Relationship Manager. 16 September, 2021.
120. Amgen Canada Inc. IMDELLTRA® product monograph. Mississauga, Ontario; September 11, 2024.
121. de Lemos, M. BC Cancer Provincial Pharmacy Professional Practice Leader and Drug Information Coordinator. SBAR: Preparing Tarlatamab - BC Cancer Recommendation vs. Product Monograph. BC Cancer - Systemic Therapy Program ; September 12, 2024
122. Medison Pharma Canada Inc. Immunocore Ireland Limited. KIMMTRAK® product monograph. Toronto, Ontario; June 7, 2022.
123. Pfeiffer, Connie. Head of Medical Affairs Immunocore. Personal Communication. Feb 16, 2022.
124. Janssen Inc. TECVAYLI® product monograph. Toronto, Ontario; July 26, 2023.
125. Janssen Research & Development LLC. Investigational product preparation and Administration instructions for subcutaneous Administration of 10 mg/mL and 90 mg/mL Teclistamab (JNJ-64007957) For Weight-Based Dosing in mcg/kg for Managed Access Programs - Version 6.0 . Beerse, Belgium; June 9, 2023.
126. de Lemos, M. BC Cancer Provincial Pharmacy Professional Practice Leader and Drug Information Coordinator. SBAR: Teclistamab Subcutaneous Injection for Multiple Myeloma at BC Cancer. (draft). BC Cancer - Systemic Therapy Program ; September 1, 2023

127. Pfizer Canada Inc (for Wyeth Canada). TORISEL® product monograph. Kirkland, Quebec; 21 December , 2016.
128. Anna Sivojelezova M.Sc. Medical Information Associate, Wyeth. Personal communication. 6 January, 2010.
129. Bristol-Myers Squibb Canada. VUMON® product monograph. St. Laurent, Quebec; 26 October , 2004.
130. Trissel's®2 IV Compatibility (database on the Internet). Teniposide. Thomson Reuters MICROMEDEX® 2.0; Accessed 27 April, 2011. Available at: <http://www.micromedex.com>
131. Hikma Canada Limited. Thiotepea for injection product monograph. Mississauga, Ontario; March 14, 2023.
132. AHFS DI® (Adult and Pediatric) (database on the Internet). Thiotepea. UpToDate® Lexidrug®; Accessed October 16, 2024. Updated March 11, 2024. Available at: <http://online.lexi.com>
133. BC Cancer Miscellaneous Origin Tumour Group. (MOIT) BC Cancer Protocol Summary for Solid Tumours using Intrathecal Methotrexate and/or Thiotepea and/or Cytarabine. Vancouver, British Columbia: BC Cancer; September 1 , 2021.
134. BC Cancer. Systemic Therapy Policy and Procedure III-50: Administration of High Alert Medications by the Intrathecal Route via Lumbar Puncture or Ommaya Reservoir. Vancouver, British Columbia; December 29, 2022.
135. SteriMax Inc. Thiotepea for injection product monograph. Oakville, Ontario; April 5, 2023.
136. Sanofi Genzyme. THYROGEN® product monograph. Mississauga, Ontario; July 29, 2021.
137. BeiGene Ireland Ltd. TEVIMBRA® Summary of product characteristics. Dublin, Ireland; January, 2023.
138. Hoffmann-La Roche Limited. ACTEMRA® product monograph. Mississauga, Ontario; 27 October, 2017.
139. Accord Healthcare Inc. Topotecan hydrochloride for injection product monograph. Kirkland, Quebec; May 9, 2019.
140. BC Children's Hospital (Distribution - Sterile Compounding). Intrathecal or Intra-OMAYA Compounding Worksheet: Topotecan IT or Intra-Ommaya Syringe. Vancouver, British Columbia; May, 2020.
141. Poplack Potter SL, Berg S, Ingle AM, et al. Phase 2 clinical trial of intrathecal topotecan in children with refractory leptomeningeal leukemia: a Children's Oncology Group trial (P9962). *Pediatr Blood Cancer* ; 2012;58(3):362–365
142. Pfizer Canada Inc. Topotecan hydrochloride for injection product monograph. Kirkland, Quebec; April 9, 2021.
143. Sandoz Canada Inc. Topotecan injection product monograph. Boucherville, Quebec; March 29, 2019.
144. Hoffmann-La Roche Limited. HERCEPTIN® product monograph. Mississauga, Ontario; May 7, 2020.
145. Celltrion Healthcare Co Ltd (distributed by Teva Canada Limited). HERZUMA® product monograph. Toronto, Ontario; 20 October, 2020.
146. BGP Pharmacy ULC. OGIVRI® product monograph. Etobicoke, Ontario; 3 May, 2019.
147. Pfizer Canada-ULC. TRAZIMERA® product monograph. Kirkland, Quebec; August 15, 2019.
148. AstraZeneca Canada Inc. ENHERTU® product monograph. Mississauga, Ontario; 15 April, 2021.
149. AstraZeneca Canada Inc. Medical Information. 5 October, 2021.
150. Hoffmann-La Roche Limited. KADCYLA® product monograph. Mississauga, Ontario; 11 September, 2013.
151. Hoffmann-La Roche Limited. KADCYLA® product monograph. Mississauga, Ontario; July 3, 2020.
152. AstraZeneca Canada Inc. IMJUDO® product monograph. Mississauga, Ontario; January 26, 2024.
153. Medexus Inc. TRECONDYV® product monograph. Bolton, Ontario; June 25, 2021.
154. Erin Wallace. Ontario Territory Manager, Medexus Pharmaceuticals Inc. Personal Communication - TRECONDYV®. November 2, 2021.
155. Medical Information team. Medexus Pharmaceuticals Inc. Personal Communication - TRECONDYV®. November 1, 2021.
156. medac GmbH. TRECONDI® summary of product characteristics. Wedel, Germany;
157. medac-UK. TREOSULFAN injection® summary of product characteristics. Wedel, Germany; 26 Jan, 2017.
158. medac-UK. TREOSULFAN injection® Details about Handling and Stability. Hamburg, Germany; August, 2008.
159. Henrik Fenger. Management Associate, International Division medac. Personal communication. 03 March, 2010.
160. medac-UK. TREOSULFAN injection® summary of product characteristics. Hamburg, Germany; 24 Jun, 2008.
161. Pfizer Canada-ULC. Vinblastine sulfate injection product monograph. Kirkland, Quebec; April 18, 2019.
162. Lexi-Drugs® (database on the Internet). VinBLASTine. Lexi-Comp Inc.; Accessed 30 January, 2020. Available at: <http://online.lexi.com>
163. BC Cancer Provincial Systemic Therapy Program. Policy V-40: Dispensing and Labelling of Vinca Alkaloid Preparations. Vancouver, British Columbia: BC Cancer; 1 April , 2015.
164. World Health Organization. Information Exchange System - Vincristine (and other vinca alkaloids) should only be given intravenously via a minibag. Alert No. 115 ed. Geneva, Switzerland: World Health Organization; 18 July , 2007.
165. Teva Canada Limited. Vinblastine sulfate injection product monograph. Toronto, Ontario; 22 October, 2019.
166. Pfizer Canada-ULC. Vincristine sulfate injection product monograph. Kirkland, Québec; July 26, 2021.

167. Teva Canada Limited. Vincristine sulfate injection® product monograph. Scarborough, Ontario; March 27, 2014.
168. Pharmaceutical Partners of Canada. Vinorelbine Injection product monograph. Richmond Hill, Ontario; 15 January , 2008.
169. Generic Medical Partners Inc. Vinorelbine Injection product monograph. Toronto, Ontario; 3 September, 2014.
170. Teva Canada Limited. Vinorelbine tartrate for Injection product monograph. Toronto, Ontario; 20 March, 2014.
171. Innomar Strategies Inc. (for Dr. Reddy's Laboratories Limited). Zoledronic acid for injection concentrate® product monograph. Oakville, Ontario; 11 March, 2015.
172. Marcan Pharmaceuticals Inc. Zoledronic acid concentrate for injection product monograph. Ottawa, Ontario; 5 February, 2018.
173. MDA Inc. Zoledronic acid for injection product monograph. Mississauga, Ontario; 11 August, 2015.
174. Novartis Pharmaceuticals Canada Inc. ZOMETA® product monograph. Dorval, Quebec; 26 July, 2013.
175. Sandoz Canada Inc. Zoledronic Acid - Z® product monograph. Boucherville, Quebec; 02 December, 2016.
176. The United States Pharmacopeia (. General Chapter 797: Pharmaceutical compounding - sterile preparations. USP 27-NF 22. Rockville, Maryland: The United States Pharmacopeial Convention, Inc.; 2004.
177. Kastango ES. The ASHP discussion guide for compounding sterile preparations. Bethesda (MD): American Society of Health-System Pharmacists, Inc.; 2004. p. 5