

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Lambrolizumab – see Pembrolizumab						
Leucovorin 50 mg/5 mL 500 mg/50 mL (Hospira) (F)(PFL) no preservative ¹	N/A	10 mg/mL ¹	5 mL vial: discard unused portion ² 50 mL vial: 8 h ¹	syringe ³ 0.05-10 mg/mL NS, D5W, Ringer's, Lactated Ringer's, D10W, D5NS ^{1,2} (e.g., 50-250 mL*)	7 d F, ³ 48 h RT ^{3,4} 24 h RT in NS , D5W, Lactated Ringer's, Ringer's ¹ 8 h RT in D10W, D5- NS ¹	

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Leucovorin 50 mg/5 mL 500 mg/50 mL (Teva) (F)(PFL) no preservative ⁵	N/A	10 mg/mL ¹	discard unused portion ¹	syringe	8 h ^{6,7}	
				0.4 - 4.8 mg/mL NS , D5W ⁸ (e.g., 50-250 mL*)	72 h F , RT ⁸	
				0.06 - 0.4 mg/mL NS , D5W ⁵	24 h RT in NS ⁵ 12 h RT in D5W ⁵	
				0.06 - 1 mg/mL Ringer's, Lactated Ringer's, D10W, D10NS ⁵	24 h RT in Ringer's, Lactated Ringer's ⁵ 12 h RT in D10W ⁵ 6 h RT in D10NS ⁵	

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Mechlorethamine 10 mg (Ovation Pharmaceuticals/Merck) (RT,PFL) no preservative ⁹	do NOT use if discoloured or water droplets form in vial before reconstitution ⁹ 10 mL SWI or NS ⁹ record time of reconstitution	1 mg/mL ⁹	use within 4 h of reconstitution RT ^{7,10}	syringe ⁹	complete administration within 4 h of reconstitution RT ^{7,9,10}	
				100 mL NS ^{9,11}	complete administration within 4 h of reconstitution RT ^{7,9,11}	
Melphalan 50 mg (GSK) (RT)(PFL) no preservative ¹²	10mL supplied diluent ¹² immediately after adding diluent, shake vigorously ¹² record time of reconstitution	5 mg/mL ¹²	2 h RT ¹² do NOT refrigerate	0.1 – 0.45 mg/mL in NS only ¹² (e.g., greater than 45 mg and less than or equal to 110 mg in 250 mL NS)*	complete administration within 60 min from time of initial reconstitution at RT ¹⁰	

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Mesna 1000 mg/10mL (Fresenius Kabi) (RT) preservative ¹³	N/A	100 mg/mL ¹³	14 d F , RT ^{7,13}	Greater than or equal to 1mg/mL ¹³ NS or D5W	48 h F , 24 h RT ¹³	
Methotrexate 50 mg/2mL 500 mg/20mL 1 g/40mL 5 g/200mL (Hospira) (RT)(PFL) no preservative ¹⁴	N/A	25 mg/mL ¹⁴	50mg: discard unused portion ¹⁴ 500mg, 1 g, 5 g: 8 h F , RT ¹⁴	syringe	2 d F , RT ^{10,15,16}	- for high-dose regimens (e.g., 1- 12 g/m ² as a single dose) ¹⁷⁻²¹ : use preservative-free methotrexate ¹⁰ - do not use for IT injection
				0.4–2 mg/mL ²² 100 mL* NS , D5W	24 h RT ²²	
				high dose (e.g., 1-12 g/m ² as a single dose) ¹⁷⁻²¹ : 1000 mL* NS	24 h RT ^{2,22,23}	
Methotrexate IT Injection¹⁴ : Only preservative free methotrexate may be administered by the intrathecal route ¹⁴ 50 mg/2mL ²⁴ (Hospira) (RT)(PFL) no preservative ¹⁴	N/A	25 mg/mL ¹⁴	discard unused portion ¹⁴	qs to 6 mL with preservative free NS ²⁵	use within 4 h of initial puncture ^{7,10}	- auxiliary label ²⁶ : "IT" - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag ²⁶

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Methotrexate 50 mg/2mL 500 mg/20mL (Hospira) (RT)(PFL) preservative ¹⁴	N/A	25 mg/mL ¹⁴	14 d F ^{27,28}	syringe	14 d F ^{4,15}	- for high-dose regimens (e.g., 1- 12 g/m ² as a single dose) ¹⁷⁻²¹ : use preservative-free methotrexate ¹⁰ - do not use for IT injection
				0.4–2 mg/mL ²² 100 mL* NS, D5W ¹⁴	24 h RT ²⁷	
Mitomycin 5 mg 20 mg (Teva/Novopharm) (RT)(PFL) no preservative ²⁹	SWI 5 mg: 10 mL 20 mg: 40 mL shake well ²⁹	0.5 mg/mL ²⁹	48 h F, RT ^{7,29} **(PFL) ²⁹	syringe ⁷	14 d F, 48 h RT ^{7,29}	
Mitomycin intravesical 5 mg 20 mg (Teva/Novopharm) (RT)(PFL) no preservative ²⁹	SWI 5 mg: 10 mL 20 mg: 40 mL shake well ²⁹	0.5 mg/mL ²⁹	48 h F, RT ^{7,29} **(PFL) ²⁹	syringe ⁷	14 d F, 48 h RT ^{7,29}	

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Mitomycin intraoperative 5 mg 20 mg (Teva/Novopharm) (RT)(PFL) no preservative ²⁹	SWI 5 mg: 10 mL 20 mg: 40 mL shake well ²⁹	0.5 mg/mL ²⁹	48 h F, RT ^{7,29} **(PFL) ²⁹	0.02-0.04 mg/mL ²⁹ NS , D5W, sodium lactate ²⁹	3 h RT: D5W 12 h RT: NS 24 h RT: sodium lactate ²⁹	
Mitomycin 5 mg 20 mg (BMS) (RT)(PFL) no preservative ³⁰	SWI 5 mg: 10 mL 20 mg: 40 mL shake well ³⁰	0.5 mg/mL ³⁰	48 h F, RT ^{7,30} **(PFL) ³⁰	syringe ¹⁰	14 d F, 48 h RT ^{10,31}	
Mitomycin intravesical 5 mg 20 mg (BMS) (RT)(PFL) no preservative ³⁰	SWI 5 mg: 10 mL 20 mg: 40 mL shake well ³⁰	0.5 mg/mL ³⁰	48 h F, RT ^{7,30} **(PFL) ³⁰	syringe ¹⁰	14 d F, 48 h RT ^{10,31}	

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Mitomycin intraperitoneal 5 mg 20 mg (BMS) (RT)(PFL) no preservative ³⁰	SWI 5 mg: 10 mL 20 mg: 40 mL shake well ³⁰	0.5 mg/mL ³⁰	48 h F, RT ^{7,30} **(PFL) ³⁰	0.02–0.04 mg/mL NS , D5W, sodium lactate ³⁰	12 h RT: NS 3h: D5W 24 h: sodium lactate ³⁰	
mitoXANTRONE 20 mg/10 mL (Fresenius Kabi) (RT) no preservative ³²	N/A	2 mg/mL ³²	discard unused portion ³²	NS , D5W ³² Greater than or equal to *50 mL ³²	24 h RT ³²	
mitoXANTRONE 20 mg/10 mL 25 mg/12.5 mL (Hospira) (RT)(PFL) no preservative ³³	N/A	2 mg/mL ³³	discard unused portion ³³	0.2-0.6 mg/mL ³³ NS , D5W ³³ Greater than or equal to *50 mL ³³	NS: 24 h F, RT ³³ **(PFL) ³³	
mitoXANTRONE 20 mg/10 mL (Teva/Novopharm) (RT)(PFL) no preservative ³⁴	N/A	2 mg/mL ³⁴	discard unused portion ³⁴	NS , D5W ³⁴ Greater than or equal to *50 mL ³⁴	24 h RT ³⁴ **(PFL) ³⁵	

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p>Nivolumab 40 mg/4 mL 100 mg/10 mL (BMS) (F)(PFL) do not shake no preservative³⁶</p>	<p align="center">N/A</p>	<p align="center">10 mg/mL³⁶</p>	<p align="center">discard unused portion³⁶</p>	<p align="center">1-10 mg/mL³⁶ NS, D5W³⁶ mix by gentle inversion³⁶</p>	<p align="center">complete administration within 8 h RT or 24 h F³⁷</p>	<ul style="list-style-type: none"> - administer with a 0.2 to 1.2 micron low protein binding in-line filter³⁶ - flush line with NS or D5W following infusion³⁶ - discard if cloudy or has pronounced colour change (should be clear to pale yellow)³⁶
<p>oBINutuzumab 1000 mg/40 mL (Hoffman-La Roche) (F)(PFL)^{**} do not shake no preservative³⁸</p>	<p align="center">N/A</p>	<p align="center">25 mg/mL³⁸</p>	<p align="center">discard unused portion²</p>	<p align="center">100 mg: in 100 mL NS³⁸ 900 mg: in 250 mL NS³⁸ 1000 mg: in 250 mL NS³⁸</p>	<p align="center">24 h F, 48 h RT^{38,39}</p>	<ul style="list-style-type: none"> -once removed from the fridge, diluted product is stable for an additional 48 h RT^{38,39} - do NOT shake³⁸ - do NOT use dextrose containing solutions³⁸

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Octreotide 50 mcg/mL 100 mcg/mL 500 mcg/mL (Omega) (F)(PFL) no preservative ⁴⁰ multidose vial: 1000 mcg/5 mL (Omega) (F)(PFL) preservative ⁴⁰	N/A	50 mcg/mL 100 mcg/mL 500 mcg/mL ⁴⁰	Use within 4 h ⁴⁰	NS ⁴⁰ volume adjusted to ensure a continuous infusion of octreotide at 25 mcg/hour ⁴⁰	24 h RT ⁴⁰	
		200 mcg/mL ⁴⁰	15 d F ⁴⁰	NS ⁴⁰ volume adjusted to ensure a continuous infusion of octreotide at 25 mcg/hour ⁴⁰	24 h RT ⁴⁰	
Octreotide 50 mcg/mL 100 mcg/mL 500 mcg/mL (Teva/Novopharm) (F)(PFL) no preservative ⁴¹ multidose vial: 1000 mcg/5 mL (Teva/Novopharm) (F)(PFL) preservative ⁴¹	N/A	50 mcg/mL 100 mcg/mL 500 mcg/mL ⁴¹	discard unused portion ⁴¹	SC syringe ⁴¹	single use vials: use within 4 h multidose vials: use within 14 d F ^{28,41}	
		200 mcg/mL ⁴¹	14 d F ^{28,41}	infusion: NS ⁴¹	single use or multidose vials: 24 h RT ⁴¹	

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Octreotide (SANDOSTATIN®) 1000 mcg/5 mL (Novartis) (F)(PFL) preservative ⁴²	N/A	200 mcg/mL ⁴²	discard unused portion ⁴³	50–200 mL NS ^{10,44,45} SC infusion: adjust volume to ensure infusion rate of 25 mcg/h ⁴⁴	24 h RT ⁴⁴	
Octreotide (SANDOSTATIN®) 50 mcg/1 mL 100 mcg/1 mL 500 mcg/1 mL (Novartis) (F)(PFL) no preservative ⁴²	N/A	50 mcg/mL 100 mcg/mL 500 mcg/mL ⁴²	discard unused portion ⁴⁴	50-100 mL ^{10,45} NS ⁴⁴ SC infusion: adjust volume to ensure infusion rate of 25 mcg/h ⁴⁴	24 h RT ⁴⁴	
Octreotide (SANDOSTATIN LAR®) 10 mg 20 mg 30 mg (Novartis) (F)(PFL) no preservative ⁴³	2 mL supplied diluent gently run 2 mL down sides of the vial; do NOT disturb for 2–5 min, then swirl moderately ⁴⁴ record time of reconstitution	10 mg: 5 mg/mL 20 mg: 10 mg/mL 30 mg: 15 mg/mL ⁴⁴	discard unused portion ⁴⁴	deep intragluteal administration only ⁴⁴	use within 4 h of initial reconstitution ^{7,44}	- do NOT shake

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p>oFatumumab 100 mg/ 5 mL 1000 mg/50 mL (GlaxoSmithKline) (F)(PFL) no preservative⁴⁶</p>	<p align="center">N/A</p>	<p align="center">20 mg/mL⁴⁶</p>	<p align="center">discard unused portion²</p>	<p align="center">1000 mL NS⁴⁶ or alternatively, 2000 mg doses may be supplied in 2 x 500 mL NS⁴⁶ withdraw volume from bag equal to volume of drug to be added⁴⁶</p>	<p align="center">48 h RT⁴⁶</p>	<p>- administer with 0.2 micron low protein binding in- line filter⁴⁶ - do NOT shake; mix by slow inversion to avoid formation of foam⁴⁶ - solution may contain a small quantity of drug particles; do not administer if solution is cloudy or discoloured⁴⁶</p>
<p>Oxaliplatin 50 mg 100 mg (Actavis) (RT)(PFL) no preservative⁴⁷</p>	<p align="center">50 mg: 10 mL SWI, D5W⁴⁷ 100 mg: 20 mL SWI, D5W⁴⁷</p>	<p align="center">5 mg/mL⁴⁷</p>	<p align="center">discard unused portion⁴⁷</p>	<p align="center">250-500 mL D5W (0.2-0.7 mg/mL)⁴⁷ do NOT use NS or other chloride- containing solution⁴⁷ do NOT use aluminum-containing needle and syringe⁴⁷</p>	<p align="center">0.2-2 mg/mL: 24 h F⁴⁷</p>	<p>- do NOT use aluminum- containing needle, syringe or tubing⁴⁷</p>

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Oxaliplatin 50 mg/10 mL 100 mg/20 mL 200 mg/40 mL (Hospira/Pfizer) (RT) no preservative ⁴⁸	N/A	5 mg/mL ⁴⁸	discard unused portion ⁴⁸	250-500 mL D5W (0.2 - 0.7 mg/mL ⁴⁸) do NOT use NS or other chloride- containing solution ⁴⁹ do NOT use aluminum-containing needle and syringe ⁴⁹	0.2 - 0.4 mg/mL: 24 h RT or 5 d F plus an additional 8 h RT ^{48,50} 0.5 – 2 mg/mL: 24 h RT or 14 d F plus an additional 8 h RT ^{48,50}	- do NOT use aluminum- containing needle, syringe or tubing ⁴⁸
Oxaliplatin 50 mg/10 mL 100 mg/20 mL 200 mg/40 mL (sanofi-aventis) (RT)(PFL) no preservative ⁵¹	N/A	5 mg/mL ⁵¹	discard unused portion ⁵¹	250–500 mL D5W ⁵¹ (0.2-2 mg/mL) ^{51,52} do NOT use NS or other chloride- containing solution ⁵¹ do NOT use aluminum-containing needle and syringe ⁵¹	0.2-1.3 mg/mL: 48 h RT, 14 d F ^{4,52,53} 1.3-2 mg/mL: 24 h RT, 48 h F ⁵¹	- do NOT use aluminum- containing needle, syringe or tubing ⁵¹

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Oxaliplatin 50 mg/10 mL 100 mg/20 mL 150 mg/30 mL 200 mg/40 mL (Sandoz) (RT)(PFL) no preservative ⁴⁹	N/A	5 mg/mL ⁴⁹	2 d F, RT ^{2,54}	0.2-0.7 mg/mL ⁴⁹ 250-500 mL D5W ⁴⁹ do NOT use NS or other chloride- containing solution ⁴⁹ do NOT use aluminum-containing needle and syringe ⁴⁹	0.2-2 mg/mL: 24 h RT, 48 h F ⁴⁹	
Oxaliplatin 50 mg/10 mL 100 mg/20 mL 200 mg/40 mL (Teva) (RT)(PFL) no preservative ⁵⁵	N/A	5 mg/mL ⁵⁵	discard unused portion ⁵⁵	250-500 mL D5W (0.2-0.7 mg/mL) ⁵⁵ do NOT use NS or other chloride- containing solution ⁵⁵ do NOT use aluminum-containing needle and syringe ⁵⁵	0.2-2 mg/mL: 24 h RT, 48 h F ⁵⁵	- do NOT use aluminum- containing needle, syringe or tubing ⁵⁵

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
PACLitaxel 30 mg/5 mL 100 mg/16.7 mL 300 mg/50 mL (Accord) (RT)(PFL) no preservative ⁵⁶	N/A	6 mg/ mL ⁵⁶	30 mg/5 mL or 100 mg/16.7 mL vial: 48 h RT ^{2,56} 300 mg/50 mL vial: 24 h RT ⁵⁶	0.3-1.2 mg/mL in NS , D5W, D5NS, D5LR ⁵⁶ (e.g., 100-1000 mL)*	complete administration within 27 h RT ⁵⁶	- use non-DEHP bag and tubing with 0.22 micron in-line filter ⁵⁶ - avoid excessive shaking ⁵⁶
PACLitaxel 30 mg/5 mL 100 mg/16.7 mL 300 mg/50 mL (Biolyse) (RT) ⁵⁷ no preservative ⁵⁸	N/A	6 mg/mL ⁵⁸	48 h RT ^{2,59}	0.3-1.2 mg/mL in NS , D5W ⁵⁸ (e.g., 100-1000 mL)*	complete administration within 27 h RT ^{58,60}	- use non-DEHP bag and tubing with 0.22 micron in-line filter ⁵⁸
				0.1 mg/mL in NS ⁶¹	44 h F , RT ⁶¹	
				0.012-0.12 mg/mL in NS ⁶²	16 h RT ⁶⁰	
				devices with spikes (e.g., chemo dispensing pins) may be used with vials ⁶³		

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
PACLitaxel 30 mg/5 mL 100 mg/16.7 mL 150 mg/25 mL 300 mg/50 mL (Hospira) (RT)(PFL) preservative ⁶⁴	N/A	6 mg/mL ⁶⁵	48 h RT ^{2,65,66}	0.3-1.2 mg/mL in NS , D5W, D5NS, D5LR ⁶⁵ (e.g., 100-1000 mL)*	complete administration within 27 h RT ⁶⁵	- use non-DEHP bag and tubing with 0.22 micron in-line filter ⁶⁵
PACLitaxel, nanoparticle, albumin- bound (nab) 100 mg (Celgene) (RT)(PFL) no preservative ⁶⁷	20 mL NS ⁶⁷ - slowly direct diluent against side of vial (i.e., greater than or equal to 1 min) during reconstitution ⁶⁷ - let stand for greater than or equal to 5 min to wet powder ⁶⁷ - gently swirl or invert for greater than or equal to 2 min ⁶⁷	5 mg/mL ⁶⁷	use immediately (RT) or 8 h F ⁶⁷ **(PFL) ⁶⁷	in empty sterile PVC, non-PVC, or non- DEHP infusion bag ⁶⁷	48 h F plus an additional 8 h RT ⁶⁸	- each vial contains 900 mg human albumin ⁶⁷ - to prevent foaming, do NOT inject NS directly onto the powder ⁶⁷ - some settling may occur; use mild agitation to resuspend ⁶⁷ - administer using a 15 micron filter ONLY (NOTE: use of filters with a pore size less than 15 microns may cause filter blockage) ^{69,70}

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Pamidronate 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Hospira) (RT) no preservative ⁷¹	N/A	3 mg/mL 6 mg/mL 9 mg/mL ⁷¹	discard unused portion ⁷¹	0.06–0.36 mg/mL NS , D5W ⁷¹ e.g., 250 mL* NS ⁷²	24 h F followed by 24 h RT (total 48 h) ⁷¹ **(PFL) ⁷¹	- do NOT mix with calcium containing solution (e.g., Ringer's) ⁷¹
Pamidronate 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Omega) (RT) no preservative ⁷³	N/A ⁷³	3 mg/mL 6 mg/mL 9 mg/mL ⁷³	discard unused portion ⁷³	0.06–0.36 mg/mL NS , D5W ⁷³ e.g., 250 mL* NS ⁷²	24 h F followed by 24 h RT (total 48 h) ⁷³ **(PFL) ⁷³	- do NOT mix with calcium containing solution (e.g., Ringer's) ⁷³
Pamidronate 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Fresenius Kabi) (RT) no preservative ⁷⁴	N/A ⁷⁴	3 mg/mL 6 mg/mL 9 mg/mL ⁷⁴	discard unused portion ⁷⁴	NS ; D5W ⁷⁴ Less than or equal to 0.36 mg/mL ⁷⁴	24 h RT ⁷⁴	- do NOT mix with calcium containing solutions ⁷⁴
Pamidronate 30 mg/10 mL 60mg/10 mL 90 mg/10 mL (Sandoz Canada) RT no preservative ⁷⁵	N/A ⁷⁵	3 mg/mL 6 mg/mL 9 mg/mL ⁷⁵	discard unused portion ^{75,28}	NS ; D5W ⁷⁵	24 h RT ⁷⁵	- do NOT mix with calcium containing solution (e.g., Ringer's) ⁷⁵

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p>PANitumumab 100 mg/5 mL 400 mg/20 mL (Amgen) (F)(PFL) do not shake no preservative⁷⁶</p>	<p align="center">N/A</p>	<p align="center">20 mg/mL⁷⁶</p>	<p align="center">discard unused portion⁷⁶</p>	<p>Less than or equal to 1000 mg: 100 mL NS⁷⁶</p> <p>Greater than 1000mg: 150 mL NS⁷⁶</p> <p>1-10mg/mL^{76,77}</p>	<p align="center">24 h F, 6 h RT^{76,77}</p>	<ul style="list-style-type: none"> - administer with 0.2 or 0.22 micron low protein binding in-line filter⁷⁶ - solution may contain particulates which do not affect product quality⁷⁶ - do not administer if discoloured⁷⁶

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p>pegaspargase (PEG-asparaginase) (pegylated asparaginase <i>E. coli</i>) 3750 units/5 mL (Baxalta) (F)(PFL) no preservative⁷⁸</p>	<p align="center">N/A</p>	<p align="center">750 units/mL⁷⁸</p>	<p align="center">discard unused portion⁷⁸</p>	<p align="center">IM: max volume: 2 mL in children and adolescents; 3 mL in adults</p> <p align="center">if volume greater than above, use multiple sites⁷⁸</p>	<p align="center">syringe: discard at end of day^{2,78}</p>	<p>- discard cloudy solution⁷⁸ - do not shake⁷⁸ - do not use if stored out of refrigerator for greater than 48 h⁷⁹ - do not use if previously frozen⁷⁹</p>
<p align="center">IV: 100 mL NS, D5W⁷⁸</p>	<p align="center">bag: complete administration within 48 h F⁷⁹</p> <p align="center">protect bag from direct sunlight during infusion⁷⁹</p>					

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Pembrolizumab 50 mg (Merck) (F) no preservative ⁸⁰	2.3 mL SWI ⁸⁰ direct diluent against side of vial during reconstitution to avoid foaming ⁸⁰ allow up to 5 minutes for bubbles to clear ⁸⁰ do NOT shake ⁸⁰	25 mg/mL ⁸⁰	6 h RT, 24 h F ⁸⁰	1-10 mg/mL NS ⁸⁰	6 h RT, 24 h F ⁸⁰	- use 0.2 to 5 micron in-line filter ⁸⁰ - allow vials/bags to come to RT prior to use ⁸⁰
Pemetrexed 100 mg 500 mg (Eli Lilly) (RT) no preservative ⁸¹	100 mg: 4.2 mL preservative-free NS 500 mg: 20 mL preservative-free NS ⁸¹	25 mg/mL ⁸¹	24 h F, RT ⁸¹	100 mL preservative-free NS ⁸¹	24 h F, RT ⁸¹	- do NOT mix with calcium containing solution (e.g., Ringer's) ⁸²
PERTuzumab 420 mg/14 mL (Roche) (F)(PFL) no preservative ⁸³	N/A	30 mg/mL ⁸³ do NOT shake ⁸³	discard unused portion ²⁸³	250 mL NS only ⁸³ mix by gentle inversion to avoid foaming ⁸³	24 h F, RT ⁸³	- do NOT use dextrose containing solutions ⁸³

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Plerixafor 24 mg/1.2 mL (sanofi-aventis) (RT) no preservative ⁸⁴	N/A	20 mg/mL ⁸⁴	discard unused portion ⁸⁴	SC syringe ⁸⁴	48 hours RT ^{28,85}	
Porfimer 15 mg 75 mg (Axcan) (RT)(PFL) no preservative ⁸⁶	15 mg: 6.6 mL D5W 75 mg: 31.8 mL D5W ⁸⁶ record time of reconstitution	2.5 mg/mL ⁸⁶	24 h F **(PFL) ⁸⁶	syringe ⁸⁶	use within 4 h of initial reconstitution ^{7,87} **(PFL) ⁸⁶	- avoid contact with skin and eyes; protect exposed area from light ⁸⁶
Raltitrexed 2 mg (Hospira) (F, RT)(PFL) no preservative ⁸⁸	4 mL SWI ⁸⁸	0.5 mg/mL ⁸⁸	24 h F, RT ⁸⁸	50–250 mL NS , D5W ⁸⁸	24 h F, RT ⁸⁸	
Ramucirumab 100 mg/10 mL 500 mg/50 mL (Eli Lilly) (F)(PFL) (do not shake) no preservative ⁸⁹	N/A	10 mg/mL ⁸⁹	discard unused portion ⁸⁹	250 mL* NS ⁸⁹ (0.4 – 4 mg/mL) ⁹⁰ gently invert to mix ⁸⁹ do NOT shake ⁸⁹	4 h RT, 24 h F ⁸⁹	- use 0.22 micron filter ⁸⁹ - flush line with NS following administration ⁸⁹ - do NOT use dextrose containing solutions ⁸⁹

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
riTUXimab 100 mg/10 mL 500 mg/50 mL (Roche) (F)(PFL) no preservative ⁹¹	N/A	10 mg/mL ⁹¹	discard unused portion ⁹¹	1-4 mg/mL NS , D5W ⁹¹ (e.g., 250-500 mL)*	24 h F , 12 h RT ^{92,93}	- once removed from the fridge, compounded product is stable for 12h RT ^{92,93}
riTUXimab subcutaneous 1400 mg/11.7 mL (Roche) (F)(PFL) no preservative ⁹⁴	N/A	120 mg/mL ⁹⁴	discard unused portion ⁹⁴	SC syringe ⁹⁴	48 h F plus 8 h RT ⁹⁴	- contains hyaluronidase ⁹⁴
romiDEPsin 10 mg (Celgene Inc.) (RT) ⁹⁵ no preservative ²	2.2 mL of supplied diluent ^{95,96} swirl gently to mix ⁹⁵	5 mg/mL ⁹⁵	8 h RT ⁹⁵	500 mL NS ⁹⁵	24 h RT ⁹⁵	- reconstituted solution will be slightly viscous ⁹⁷ - vials contain overflow to allow for full drug recovery (drug vial contains 11 mg romidepsin; diluent vial contains 2.4 mL diluent) ⁹⁵

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Siltuximab 100 mg 400 mg (Janssen) (F)(PFL) no preservative ⁹⁸	100 mg: 5.2 mL SWI ⁹⁸ 400 mg: 20 mL SWI ⁹⁸ allow vial to come to room temperature prior to use (~30 minutes) ⁹⁸ gently swirl, do not shake ⁹⁸	20 mg/mL ⁹⁸	2 h RT ⁹⁸	250 mL D5W ⁹⁸ dilute to 250 mL final volume by withdrawing volume from bag equal to volume of drug to be added ⁹⁸	complete administration within 6 h RT ⁹⁸	- use 0.2 micron in- line filter ⁹⁸
Streptozocin 1g (Pfizer) (F)(PFL) no preservative ⁹⁹	9.5mL NS , SWI, D5W ⁹⁹	100 mg/mL ⁹⁹	48 h F, ⁹⁹ 24 h RT	syringe ⁹⁹	48 h F, 24 h RT ⁹⁹	
				50-500 mL* NS , D5W, SWI ⁹⁹	48 h F, 24 h RT ⁹⁹	
Temozolomide 100 mg (Schering) (F) do not dilute do not shake no preservative ¹⁰⁰	41 mL SWI ¹⁰⁰	2.5 mg/mL ¹⁰⁰	14 h F, RT ¹⁰⁰	empty 250 mL PVC bag ¹⁰⁰	14 h RT ¹⁰⁰	

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Temsirolimus 30 mg/1.2 mL (Wyeth) (F)(PFL) ^{101,102} no preservative ¹⁰³	1.8 mL supplied diluent ^{101,102}	10 mg/mL ^{101,102}	24 h RT ^{101,102} **(PFL) ¹⁰¹	250 mL NS ^{101,102}	complete administration within 6 h ^{101,102}	- use non-DEHP bag and tubing with in-line filter ^{101,102}
Teniposide 50 mg/5 mL (BMS) (RT) preservative ¹⁰⁴	N/A	10 mg/mL ¹⁰⁴	discard unused portion	50 – 500 mL NS or D5W for a final concentration of 0.1-1 mg/mL ¹⁰⁴	0.1-0.4 mg/mL: 24 h RT ¹⁰⁴ 1 mg/mL: complete administration within 4 h of preparation RT ^{104,105}	- do not refrigerate - use non-DEHP bag and tubing ¹⁰⁴ - do not use if precipitates ^{104,105} - contains DMA*** - excessive agitation may cause precipitation ¹⁰⁴

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Thiotepa 15 mg (Bedford) (F)(PFL) no preservative ¹⁰⁶	1.5 mL SWI ¹⁰⁶ filter through 0.22 micron filter ¹⁰⁶ record time of reconstitution	10 mg/mL ¹⁰⁶	8 h F ¹⁰⁶	50 mL * NS ¹⁰⁶	use within 4 h of initial reconstitution ^{106,107} **(PFL) ^{106,108}	- do not use if precipitates or remains opaque ¹⁰⁶ - do not use for IT injection
				syringe: reconstituted solution is hypotonic and must be further diluted with NS prior to use ¹⁰⁶ (final concentration of 0.5-1 mg/ml is nearly isotonic) ¹⁰⁹	use within 4 h of initial reconstitution ^{106,107} **(PFL) ^{106,108}	
Thiotepa IT injection: 15 mg (Bedford) (F)(PFL) no preservative ¹⁰⁶	diluents containing preservatives should NOT be used for intrathecal administration 1.5 mL SWI ¹⁰⁶ filter through 0.22 micron filter ¹⁰⁶ record time of reconstitution	10 mg/mL ¹⁰⁶	8 h F ¹⁰⁶	qs to 6 mL with preservative free NS ¹¹⁰	use within 4 h of initial reconstitution ^{106,107} **(PFL) ^{106,108}	- auxiliary label ²⁵ : "IT" - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag ²⁵ - do not use if precipitates or remains opaque ¹⁰⁶

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Thyrotropin alfa 1.1 mg (Genzyme) (F)(PFL) no preservative ¹¹¹	1.2 mL SWI ¹¹¹ swirl contents ¹¹¹ do not shake	0.9 mg/mL ¹¹¹	24 h F ¹¹¹	syringe ¹¹¹	24 h F ¹¹¹	
Topotecan 4 mg (Hospira) (F)(PFL) no preservative ¹¹²	N/A ¹¹²	1 mg/mL ¹¹²	discard unused portion ¹¹³	0.02–0.5 mg/mL ¹¹² 50-100 mL NS , D5W ¹¹³	24 h F , RT ¹¹²	
Topotecan 4 mg (Mylan) (RT)(PFL) no preservative ¹¹⁴	4 mL SWI ¹¹⁴	1 mg/mL ¹¹⁴	24 h F , RT ¹¹⁴	0.02 – 0.5 mg/mL ¹¹⁴ 50-100 mL NS , D5W ¹¹⁴	24 h F , RT ¹¹⁴	
Topotecan 4 mg/4 mL (Sandoz) (F)(PFL) no preservative ¹¹⁵	N/A	1 mg/mL ¹¹⁵	discard unused portion ¹¹⁵	0.02-0.5 mg/mL 50-100 mL NS , D5W ¹¹⁵	24 h F ¹¹⁵ **(PFL) ¹¹⁵	

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Trastuzumab (HERCEPTIN®) 440 mg (Roche) (F) preservative ¹¹⁶	20 mL supplied BWI ¹¹⁶ swirl vial gently; allow to stand undisturbed for 5 min ¹¹⁶	21 mg/mL ¹¹⁶	14 d F ²	250 mL NS only ¹¹⁶ Do NOT use dextrose containing solutions ¹¹⁶	24 h F, RT ¹¹⁶	- do NOT shake ¹¹⁶

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p>Trastuzumab Emtansine (KADCYLA®) 100 mg 160 mg (Hoffmann-La Roche) (F)(PFL) no preservative¹¹⁷</p>	<p>100 mg vial: 5 mL SWI¹¹⁷</p> <p>160 mg vial: 8 mL SWI¹¹⁷</p> <p>swirl gently until completely dissolved; do NOT shake¹¹⁷</p>	<p>20 mg/mL¹¹⁷</p>	<p>24 h F¹¹⁷ do NOT freeze¹¹⁷</p>	<p>250 mL NS or 0.45% sodium chloride¹¹⁷</p> <p>do NOT shake¹¹⁷</p>	<p>24 h F¹¹⁷ do NOT freeze¹¹⁷</p>	<p>- do not use if reconstituted solution contains visible particulates or is cloudy or discolored¹¹⁷ - dextrose 5% solutions cause aggregation of the protein; do not dilute with dextrose containing solutions¹¹⁷ - use a 0.2 micron in-line filter or 0.22 micron polyethersulfane (PES) filter to administer infusions prepared in NS; filter is optional for solutions in 0.45% NS¹¹⁷</p>

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
TRC105 (Carotuximab) 100 mg/4 mL 200 mg/8 mL 400 mg/16 mL (Tracon) (F)(PFL) no preservative ¹¹⁸	N/A	25 mg/mL ¹¹⁸	discard unused portion ²	0.6 – 10 mg/mL NS ¹¹⁹ invert gently to mix	complete infusion within 8 h RT, 24 h F ^{118,119}	- use a 0.2 micron in-line filter for administration ¹¹⁸

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p>Treosulfan 1 g 5 g (medac) (RT) no preservative¹²⁰</p>	<p>pre-heat SWI to 30°C (not higher) shake vial carefully before adding the warmed SWI 1 g vial: 20 mL SWI, while slightly shaking vial and syringe; continue shaking the reconstituted solution for another 2 min¹²⁰ 5 g vial: 100 mL SWI, while slightly shaking vial and syringe; continue shaking the reconstituted solution for another 2 min¹²⁰</p>	<p>50 mg/mL¹²⁰</p>	<p>48 h RT^{7,120}</p>	<p>undiluted¹²¹ dilute with NS or D5W in empty infusion bag for final concentration = 20 mg/mL¹²⁰</p>	<p>48 h RT^{7,120}</p>	<p>- compatible with polytetrafluoroethylene filters¹²⁰ - may require vigorous shaking to reconstitute¹²⁰</p>

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
vinBLASTine 10 mg/10 mL (Hospira) (F)(PFL) no preservative ¹²²	N/A	1 mg/mL ¹²²	discard unused portion ¹²²	25-50 mL NS , D5W ¹²³	24 h F , RT ^{124,125}	- auxiliary label: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{126,127}
vinBLASTine 10 mg/10 mL (Teva) (F)(PFL) no preservative ¹²⁸	N/A	1 mg/mL ¹²⁸	discard unused portion ¹²⁸	25-50 mL NS , D5W ^{123,129}	use within 4 h of initial puncture ²	- auxiliary label: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{126,127}

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
vinCRISTine 2 mg/2 mL 5 mg/5 mL (Hospira) (F)(PFL) no preservative ¹³⁰	N/A	1 mg/mL ¹³⁰	8 h F, RT ¹³⁰	50 mL * NS , D5W ¹³⁰	24 h F, 6 h RT ¹³⁰ **(PFL) ¹³⁰	- auxiliary label: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{126,127} - for ULYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing etoposide, DOXOrubicin, vinCRISTine)

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
vinCRISStine 1 mg/1 mL 2 mg/2 mL 5 mg/5 mL (Teva) (F)(PFL) no preservative ¹³¹	N/A	1 mg/mL ¹³¹	8 h F, RT ¹³¹	0.01-0.1 mg/mL NS or D5W ¹³¹ 25-50 mL NS , D5W ¹³²	24 h F, RT ¹³¹	- auxiliary label: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{126,127} - for ULYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing etoposide, DOXOrubicin, vinCRISStine)
Vinorelbine 10 mg/1 mL 50 mg/5mL (Fresenius Kabi) (F)(PFL) no preservative ¹³³	N/A	10 mg/mL ¹³³	discard unused portion ¹³³	0.5-2.0 mg/mL ¹³³ NS , D5W, ½NS, D5½NS, Ringer's, Ringer's Lactate ¹³³	24 h F, RT ¹³³	- auxiliary label: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{126,127}

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Vinorelbine 10 mg/1 mL 50 mg/5 mL (Hospira) (F)(PFL) no preservative ¹³⁴	N/A	10 mg/mL ¹³⁴	discard unused portion ¹³⁴	0.5–2.0 mg/mL ¹³⁴ 50 mL* NS , D5W, ½NS, D5½NS, Ringer's, Ringer's Lactate ¹³⁴	24 h F , RT ¹³⁴	- auxiliary label: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{126,127}
Vinorelbine 10 mg/1 mL 50 mg/5 mL (Teva) (F)(PFL) no preservative ¹³⁵	N/A	10 mg/mL ¹³⁵	discard unused portion ¹³⁵	0.5–2.0 mg/mL ¹³⁵ 50 mL* NS , D5W, ½NS, D5½NS, Ringer's, Ringer's Lactate ¹³⁵	24 h F , RT ¹³⁵	- auxiliary label: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{126,127}

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Zoledronic acid 4 mg/5 mL (Dr Reddy's) (RT) no preservative ¹³⁶	N/A	0.8 mg/mL ¹³⁶	discard unused portion ¹³⁶	100 mL NS , D5W ¹³⁶	complete infusion within 24 h of preparation ¹³⁶ Refrigerate diluted product if not used immediately after preparation; bring to RT prior to administration ¹³⁶	- do NOT mix with calcium containing solutions ¹³⁶
Zoledronic acid (ZOMETA) 4 mg/ 5 mL (Novartis) (RT) no preservative ¹³⁷	N/A ¹³⁷	0.8 mg/ mL ¹³⁷	discard unused portion ²	100 mL NS , D5W ¹³⁷	complete infusion within 24 h of preparation ¹³⁷ Refrigerate diluted product if not used immediately after preparation; bring to RT prior to administration ¹³⁷	- do NOT mix with calcium containing solutions ¹³⁷

BC CANCER AGENCY CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
Zoledronic acid 4 mg/5 mL (Sandoz) (RT) no preservative ¹³⁸	N/A	0.8 mg/mL ¹³⁸	discard unused portion ¹³⁸	100 ml NS , D5W ¹³⁸	complete infusion within 24 h of preparation ¹³⁸ Refrigerate diluted product if not used immediately after preparation; bring to RT prior to administration ¹³⁸	- do NOT mix with calcium- or other divalent cation- containing infusion solutions (e.g., Lactated Ringer's) ¹³⁸

* Suggested volume based on usual dose range and any concentration range of stability data

** Protect from light means minimizing exposure to direct sunlight over a *storage* period. More specific information on protection from light (eg, protecting container and tubing during *administration*) will be indicated in the Under the Special Precautions/Notes column.

*** Contains DMA (N,N dimethylacetamide). Product may be incompatible with closed system transfer devices such as ChemoLock.

Centres are not to change the content locally but should forward suggestions to the Cancer Drug Manual staff.

Explanatory Notes

Stability data assumes products prepared using standard aseptic technique in biological safety cabinet at low risk for contamination according to the classification outlined in USP 797.^{31,139}

Vial stability: Stability of solution after first puncture or reconstituted solution.

Storage temperature: If information states same stability with refrigerator and room temperature storage, then fridge stability is bolded as preferred (ie, to minimize growth of micro-organisms).

Discard unused portion: Unused portion from single use vials should be discarded at the end of the day.

“overflow known” is stated if the manufacturer states overflow that is present is within acceptable limits.

“Complete administration within ___” is stated if the manufacturer specifies that the infusion must be completed in a specific time frame following preparation, usually including entire time required for preparation (from first puncture), storage, and administration of infusion.

Abbreviations

BWI = bacteriostatic water for injection
CIVI: ambulatory pump = Continuous Intravenous Infusion (e.g., elastomeric infusor)
D5W = dextrose 5% in water
DMA = N,N dimethylacetamide
F = refrigerate
Non-DEHP = not containing Di(2-ethylhexyl) phthalate (DEHP)
NS = normal saline
PFL = protect from light
RT = room temperature
SWI = sterile water for injection

References

1. Hospira Healthcare Corporation. LEUCOVORIN CALCIUM INJECTION® product monograph. Saint-Laurent, Quebec; 7 June 2007.
2. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 19 September 2007.
3. John Korontzis. Personal communication. Regulatory Affairs Associate, Leucovorin Calcium, Mayne Pharma Canada; February 2005.
4. The United States Pharmacopeial Convention, Inc. General Chapter 797: Pharmaceutical compounding - sterile preparations. USP 27-NF 22. Rockville, Maryland: The United States Pharmacopeial Convention, Inc.; 2003.
5. Teva Canada Limited. Leucovorin calcium injection® product monograph. Toronto, Ontario; 5 May 2014.
6. Novopharm Limited (Teva). LEUCOVORIN CALCIUM® Injection product information package. Toronto, Ontario; undated.
7. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 6 January 2006.
8. Jenny Yeung. Personal communication. Medical Information Specialist, Teva Canada; 12 April 2017.
9. Ovation Pharmaceuticals Inc. MUSTARGEN® Package Insert. Deerfield, Illinois; October 2005.
10. Trissel LA. Handbook on Injectable Drugs. 13th ed. Bethesda, MD: American Society of Health-System Pharmacists, Inc.; 2005.
11. Trissel LA. Handbook on Injectable Drugs. 12th ed. Bethesda, MD: American Society of Health-System Pharmacists, Inc.; 2003.
12. GlaxoSmithKline Inc. Alkeran Package Insert. Mississauga, Ontario; Montreal, Quebec; 2004.
13. Pharmaceutical Partners of Canada Inc. Mesna Package Insert. Richmond Hill, Ontario; November 2001.
14. Mayne Pharma Canada. Methotrexate Product Monograph. Montreal, Quebec; December 2003.
15. John Korontzis. Personal communication. Regulatory Affairs Associate, Methotrexate, Mayne Pharma Canada; February 2005.
16. John Korontzis. Personal communication. Regulatory Affairs Associate, Fluorouracil, Methotrexate, Vincristine, Mayne Pharma Canada; 14 March 2005.
17. BC Cancer Agency Miscellaneous Origins Tumour Group. (MOHDMTX) BCCA Protocol Summary for Treatment of Meningeal Disease (Miscellaneous Tumour Origins) using High Dose Methotrexate with Leucovorin Rescue. Vancouver, British Columbia: BC Cancer Agency; 1 Jan 2013.
18. BC Cancer Agency Sarcoma Tumour Group. (SAHDMTX) BCCA Protocol Summary for Treatment of Osteosarcoma Using High Dose Methotrexate with Leucovorin Rescue. Vancouver, British Columbia: BC Cancer Agency; 1 Nov 2012.
19. BC Cancer Agency Lymphoma Tumour Group. (LYHDMRP) BCCA Protocol Summary for Treatment of Primary Intracerebral Lymphoma with High Dose Methotrexate and rituximab. Vancouver, British Columbia: BC Cancer Agency; 1 Jun 2014.
20. BC Cancer Agency Lymphoma Tumour Group. (LYHDMTXP) BCCA Protocol Summary for Treatment of Primary Intracerebral Lymphoma with High Dose Methotrexate. Vancouver, British Columbia: BC Cancer Agency; 1 Jun 2014.
21. BC Cancer Agency Lymphoma Tumour Group. (LYHDMTXR) BCCA Protocol Summary for Treatment of Leptomeningeal Lymphoma or Recurrent Intracerebral Lymphoma with High Dose Methotrexate. Vancouver, British Columbia: BC Cancer Agency; 1 Jun 2014.
22. Hospira Healthcare Corporation. Methotrexate Injection USP® product monograph. Saint-Laurent, Quebec; 27 June 2012.

23. Trissel's®2 IV Compatibility (database on the Internet). Methotrexate sodium. Thomson Reuters MICROMEDEX® 2.0, updated periodically. Available at: <http://www.micromedex.com>. Accessed 9 March 2015.
24. BC Cancer Agency. Pharmacy Policy Number III-50-02: Methotrexate Intrathecal Injection. Vancouver, British Columbia: BC Cancer Agency; 27 January 2006.
25. BC Cancer Agency Miscellaneous Origin Tumour Group. BCCA protocol summary for solid tumours using intrathecal methotrexate and/or thiotepa and/or cytarabine (MOIT). Vancouver, British Columbia: BC Cancer Agency; 1 July 2005.
26. BC Cancer Agency. Systemic Therapy Policy III-50: Administration of Cytotoxic Drugs by the Intrathecal Route via Lumbar Puncture or Ommaya Reservoir. Vancouver, British Columbia; 31 October 2002.
27. Hospira Healthcare Corporation. Methotrexate Injection USP product monograph. Saint-Laurent, Quebec; 05 May 2009.
28. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 19 September 2007.
29. Novopharm Limited. Mitomycin for Injection Product Monograph. Scarborough, Ontario; 23 December 1996.
30. Bristol Laboratories of Canada. Mutamycin Package Insert. Montreal, Canada; 2000.
31. The United States Pharmacopeia (USP). General Chapter 797: Pharmaceutical compounding - sterile preparations. USP 27-NF 22. Rockville, Maryland: The United States Pharmacopeial Convention, Inc.; 2004.
32. Fresenius Kabi Canada Ltd. Mitoxantrone injection® product monograph. Richmond Hill, Ontario; 28 September 2016.
33. Hospira Healthcare Corporation. Mitoxantrone Injection, USP Product Monograph. Saint-Laurent, Quebec; 13 June 2007.
34. Novopharm Limited. Mitoxantrone Injection Product Monograph. Toronto, Ontario; 19 May 2005.
35. Angie Chan. Personal communication. Drug Information Pharmacist, Novopharm; 7 Mar 2008.
36. Bristol-Myers Squibb Canada. OPDIVO® product monograph. Montreal, Quebec; 24 September 2015.
37. Bristol-Myers Squibb Canada. OPDIVO® product monograph. Montreal, Quebec; 12 May 2017.
38. Hoffmann-La Roche Ltd. GAZYVA® product monograph. Mississauga, Ontario; 21 December 2015.
39. Anna Sivojelezova MSc. Personal communication. Drug Information Associate; Hoffmann-La Roche Ltd Drug Information; 24 April 2015.
40. Omega Laboratories Ltd. Octreotide Acetate Injection product monograph. Montreal, Quebec; 23 July 2010.
41. Novopharm Limited. Octreotide Injection Product Monograph. Scarborough, Ontario; 15 March 2007.
42. Novartis Pharmaceuticals Canada Inc. SANDOSTATIN® Product Monograph. Dorval, Quebec; 9 January 2001.
43. Repchinsky C, editor. Sandostatin LAR monograph, Compendium of Pharmaceuticals and Specialties. Ottawa, Ontario: Canadian Pharmacists Association; 2005. p. 1912-1916.
44. Repchinsky C editor. Compendium of Pharmaceuticals and Specialties. 12th ed. Ottawa, Ontario: Canadian Pharmacists Association; 2004.
45. Vancouver Hospital and Health Sciences Centre Pharmacy Department. Octreotide. Parenteral drug therapy manual. Vancouver, BC; February 2002.
46. GlaxoSmithKline. Guidance Booklet: compassionate use access of ofatumumab for patients with chronic lymphocytic leukemia (CLL) who are refractory to fludarabine and alemtuzumab or who are deemed inappropriate for alemtuzumab therapy. version 5; 1 February 2010.
47. Actavis Pharma Company. ACT Oxaliplatin® product monograph. Mississauga, Ontario; 1 June 2015.
48. Pfizer Canada Inc. Oxaliplatin injection product monograph. Kirkland, Quebec; 31 May 2017.
49. Sandoz Canada Inc. Oxaliplatin injection product monograph. Boucherville, Quebec; 12 August 2015.
50. Medical Information. Personal communication. Pfizer Canada Inc.; 6 June 2017.
51. sanofi-aventis Canada Inc. ELOXATIN® product monograph. Laval, Quebec; 13 January 2009.
52. Eiden C, Philibert L, Bekhtari K, et al. Physicochemical stability of oxaliplatin in 5% dextrose injection stored in polyvinyl chloride, polyethylene, and polypropylene infusion bags. Am J Health Syst Pharm 2009;66.
53. Andre P, Cisternino S, Roy A-, et al. Stability of oxaliplatin in infusion bags containing 5% dextrose injection. American Journal of Health-System Pharmacy 2007;64(18):1950-1954.
54. Katryn Vosburg. Personal communication. Drug Information & Pharmacovigilance Specialist, Sandoz Canada Inc.; 26 February 2016.
55. Teva Canada Limited. Teva-Oxaliplatin injection® product monograph. Toronto, Ontario; 11 September 2015.
56. Accord Healthcare Inc. Paclitaxel injection product monograph. Markham, Ontario; 13 August 2012.
57. Claude Mercure. Personal communication. Production Manager Biolyse Pharma Corporation; 21 December 2009.
58. Biolyse. PACLITAXEL FOR INJECTION® product monograph. St. Catherines, Ontario; 2 December 2005.
59. Claude Mercure. Personal communication. Manager, Biolyse Pharma Corporation; 24 June 2014.

60. Zeng Z, Lazakovitch E. Study IR 120: Physical and Chemical Stability Study of Paclitaxel for Injection in 0.9 % Sodium Chloride in concentration range 0.012-0.12 mg/mL. Biolyse Pharma March 2010.
61. Mercure C. Stability of 0.1 mg/mL of paclitaxel for injection in sodium chloride (0.9%) solution. St Catharines, Ontario: Biolyse Pharma; 2 February 2007.
62. Xu Q, Trissel LA, Martinez JF. Stability of paclitaxel in 5% dextrose injection or 0.9% sodium chloride injection at 4, 22, or 32 degrees C. Am J Hosp Pharm 1994;51(24):3058-60.
63. Lisa Tavano. Personal communication. Biolyse Pharma Corporation; 14 May 2012.
64. Robyn Mackenzie. Personal communication. Hospira Healthcare Corporation; 4 April 2012.
65. Hospira Healthcare Corporation. PACLITAXEL FOR INJECTION® product monograph. Saint-Laurent, Quebec; 1 September 2009.
66. Rose Toussaint. Personal communication. Hospira Canada Healthcare Corporation; 4 April 2012.
67. Celgene Inc. ABRAXANE® product monograph. Mississauga, Ontario; 24 July 2014.
68. Aisling Cahill. Drug Safety and Medical Information Specialist. Personal communication. Celgene Inc.; 23 April 2015.
69. Celgene Europe Limited. ABRAXANE® product monograph. Uxbridge, UK; 11 January 2013.
70. Celgene Inc. ABRAXANE® product monograph. Mississauga, Ontario; 18 January 2016.
71. Mayne Pharma (Canada) Inc. Pamidronate Package Insert. Montreal, Quebec; 2002.
72. BC Cancer Agency Breast Tumour Group. BCCA protocol summary for treatment of acute bone pain secondary to breast cancer metastases using pamidronate or IV clodronate (BRAVPAM). Vancouver, British Columbia: BC Cancer Agency; 1 May 2001.
73. Omega Laboratories Ltd. Pamidronate Disodium product monograph. Montreal, Quebec; 06 June 2005.
74. Pharmaceutical Partners of Canada. Pamidronate Disodium For Injection product monograph. Richmond Hill, Ontario; 18 January 2010.
75. Sandoz Canada Inc. Pamidronate injection product monograph. Boucherville, Quebec; 28 February 2006.
76. Amgen Canada. VECTIBIX® product monograph. Mississauga, Ontario; 5 March 2009.
77. Diane Lord. Personal communication. Amgen Canada Inc Medical Information Department; 19 June 2009.
78. Baxalta Canada Corporation. ONCASPAR® product monograph. Mississauga, Ontario; 24 February 2017.
79. Baxalta US Inc. ONCASPAR® product monograph. Cambridge, Massachusetts, USA; 24 February 2017.
80. Merck Canada Inc. KEYTRUDA® product monograph. Kirkland, Quebec; 5 February 2016.
81. Eli Lilly Canada. Pemetrexed product information. Toronto, Ontario; 2008.
82. Eli Lilly Canada Inc. ALIMTA® Product Monograph. Toronto, Ontario; 21 May 2004.
83. Hoffmann-La Roche Limited. PERJETA® product monograph. Mississauga, Ontario; 12 April 2013.
84. sanofi-aventis Canada Inc. MOZOBIL® product monograph. Laval, Quebec; 8 October 2014.
85. Maureen Coughlin BSc Pharm. Personal communication. Solutions in Health Inc. acting as an authorized agent of sanofi-aventis; 24 May 2017.
86. Axcan Pharma Inc. Photofrin Package Insert. Mont-Saint-Hilaire, Quebec; 2004.
87. Marie-Helene Doyon B.Pharm. M.Sc. Personal communication. Medical Information Specialist, Porfimer, Axcan Pharma. 3 March 2005.
88. Hospira Healthcare Corporation. TOMUDEX® product monograph. Saint-Laurent, Quebec; 23 April 2008.
89. Eli Lilly Canada Inc. CYRAMZA® product monograph. Toronto, Ontario; 16 July 2015.
90. Marilyn Bain BScN. Personal communication. Senior Medical Information Associate, Eli Lilly Canada, Inc.; 16 January 2017.
91. Hoffmann-La Roche Ltd. RITUXAN® product monograph. Mississauga, Ontario; 29 March 2012.
92. Hoffmann-La Roche Ltd. RITUXAN® product monograph. Mississauga, Ontario; 29 May 2014.
93. Diana Fung pharmacist. Personal communication. Hoffmann-La Roche Medical Information; 16 July 2014.
94. Hoffmann-La Roche Ltd. RITUXAN® SC product monograph. Mississauga, Ontario; 9 September 2016.
95. Celgene Inc. ISTODAX® product monograph. Mississauga, Ontario; 13 December 2016.
96. Celgene Inc. INFO Rx ISTODAX® (romidepsin) for Injection. Mississauga, Ontario; 10 July 2017.
97. Aisling Cahill, Drug Safety and Medical Information Specialist. Personal communication. Celgene Inc.; 17 July 2015.
98. Janssen Inc. SYLVANT® product monograph. Toronto, Ontario; 6 January 2016.
99. Pharmacia Canada Inc. Zanosar Package Insert. Mississauga, Ontario; March 2003.
100. Schering-Plough Canada. TEMODAL® product monograph. Kirkland, Quebec; 5 January 2009.
101. Wyeth Canada. TORISEL® product monograph. Montreal, Canada; 16 October 2008.
102. McEvoy GK, editor. AHFS 2008 Drug Information. Bethesda, Maryland: American Society of Health-System Pharmacists, Inc. p. 1226-1228.
103. Anna Sivojelezova M.Sc. Personal communication. Medical Information Associate, Wyeth. 6 January 2010.

104. Bristol-Myers Squibb Canada. VUMON® product monograph. St. Laurent, Quebec; 26 October 2004.
105. Trissel's®2 IV Compatibility (database on the Internet). Teniposide. Thomson Reuters MICROMEDEX® 2.0, updated periodically. Available at: <http://www.micromedex.com>. Accessed 27 April 2011.
106. Bedford Laboratories™. Thiotepea for Injection USP Package Insert. Bedford, Ohio; April 2001.
107. Solimando D.A. Updates of melphalan and thiotepea. Hosp Pharm 1997;32(8):1082-1088.
108. Christina Hsu Pharm. D. Personal communication. Bedford Laboratories Professional Services Department; 29 August 2008.
109. Trissel LA. Handbook on Injectable Drugs. 13th ed. Bethesda, Maryland: American Society of Health-System Pharmacists, Inc.; 2005. p. 1395-1403.
110. BC Cancer Agency Miscellaneous Origin Tumour Group. (MOIT) BCCA Protocol Summary for Solid Tumours using Intrathecal Methotrexate and/or Thiotepea and/or Cytarabine. Vancouver, British Columbia: BC Cancer Agency; 1 July 2005.
111. Genzyme Canada. Thyrogen Product Monograph. Mississauga, Ontario; 2004.
112. Hospira Healthcare. Topotecan Hydrochloride for Injection product monograph. Saint-Laurent, QC; 22 June 2012.
113. Hospira Healthcare Corporation. Topotecan hydrochloride for injection product monograph. Saint-Laurent, Quebec; 26 August 2014.
114. Mylan Pharmaceuticals ULC. Topotecan hydrochloride for injection product monograph. Etobicoke, Ontario; 18 February 2015.
115. Sandoz Canada Inc. Topotecan injection product monograph. Boucherville, Quebec; 5 September 2014.
116. Hoffman-La Roche Limited. HERCEPTIN® product monograph. Mississauga, Ontario; 16 November 2012.
117. Hoffmann-La Roche Limited. KADCYLA® product monograph. Mississauga, Ontario; 11 September 2013.
118. Tracoon Pharmaceuticals Inc. TRC105 (carotuximab) investigational brochure. San Diego, California; 14 February 2017 (version 10.0).
119. Tracoon Pharmaceuticals Inc. Clinical Protocol: A Phase 2A Study of TRC105 (with Option to Add Bevacizumab) in Patients with Refractory Gestational Trophoblastic Neoplasia (GTN). San Diego, California; 20 September 2016 Amendment #3.
120. medac UK. TREOSULFAN injection® product monograph. Hamburg, Germany; 24 June 2008.
121. Henrik Fenger. Personal communication. Management Associate, International Division medac; 03 March 2010.
122. Mayne Pharma (Canada) Inc. Vinblastine product monograph. Kirkland, Quebec; 10 August 2003.
123. Lexi-Drugs® (database on the Internet). VinBLASTine. Lexi-Comp Inc., 3 November 2014. Available at: <http://online.lexi.com>. Accessed 20 November 2014.
124. Jan Barrow. Personal communication. Supervisor, Hospira Canada Clinical Support. Saint-Laurent, Quebec; 03 December 2007.
125. Tanya Leduc. Personal communication. Acting editor, BC Cancer Agency Cancer Drug Manual; 18 Dec 2007.
126. World Health Organization. Information Exchange System: Alert No. 115 (QSM/MC/IEA.115). Geneva, Switzerland: World Health Organization; 18 July 2007.
127. BCCA Provincial Systemic Therapy Program. Labeling of vinca alkaloid syringes. Policy # V-40. Vancouver, British Columbia: BC Cancer Agency; 27 May 1999.
128. Teva Canada Limited. Vinblastine sulfate injection® product monograph. Toronto, Ontario; 1 February 2013.
129. Trissel's®2 IV Compatibility (database on the Internet). Vinblastine sulfate. Thomson Reuters MICROMEDEX® 2.0, updated periodically. Available at: <http://www.micromedex.com>. Accessed 8 June 2015.
130. Mayne Pharma (Canada) Inc. Vincristine Package Insert. Montreal, QC; Undated.
131. Teva Canada Limited. Vincristine sulfate injection® product monograph. Scarborough, Ontario; 27 March 2014.
132. Lexi-Drugs® (database on the Internet). VinCRISTine. Lexi-Comp Inc., 3 June 2015. Available at: <http://online.lexi.com>. Accessed 9 June 2015.
133. Pharmaceutical Partners of Canada. Vinorelbine Injection product monograph. Richmond Hill, Ontario; 15 January 2008.
134. Mayne Pharma (Canada) Inc. Vinorelbine Product Monograph. Montreal, QC; Undated.
135. Teva Canada Limited. Vinorelbine tartrate for Injection product monograph. Toronto, Ontario; 20 March 2014.
136. Innomar Strategies Inc. (for Dr. Reddy's Laboratories Limited). Zoledronic acid for injection concentrate® product monograph. Oakville, Ontario; 11 March 2015.
137. Novartis Pharmaceuticals Canada Inc. ZOMETA® product monograph. Dorval, Quebec; 26 July 2013.
138. Sandoz Canada Inc. Zoledronic Acid - Z® product monograph. Boucherville, Quebec; 02 December 2016.
139. Kastango ES. The ASHP discussion guide for compounding sterile preparations. Bethesda (MD): American Society of Health-System Pharmacists, Inc.; 2004. p. 5.