

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Leucovorin</b> 50 mg/5 mL (GMP) (F)(PFL) no preservative <sup>1</sup>	N/A	10 mg/mL <sup>1</sup>	discard unused portion <sup>1</sup>	syringe	8 h RT <sup>1,2</sup>	
				0.05-10 mg/mL NS, D5W, Ringer's, LR, D10W, D5NS <sup>1,2</sup>  (e.g., 50-250 mL*)	<b>NS</b> , D5W, LR, Ringer's: 24 h RT <sup>1</sup>  D10W, D5-NS: 8 h RT <sup>1</sup>	
<b>Leucovorin</b> 50 mg/5 mL 500 mg/50 mL (Pfizer/Hospira) (F)(PFL) no preservative <sup>3</sup>	N/A	10 mg/mL <sup>3</sup>	8 h <sup>3</sup>	syringe	8 h RT <sup>3</sup>	
				0.05 – 10 mg/mL <b>NS</b> , D5W, LR, Ringer's, D10W, D5NS <sup>3</sup>  (e.g., 50-250 mL*)	<b>NS</b> , D5W, LR, Ringer's: 24 h RT <sup>3</sup>  D10W, D5NS: 8 h RT <sup>3</sup>	

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Leucovorin</b> 50 mg/5 mL 500 mg/50 mL (Teva) (F)(PFL) no preservative <sup>4</sup>	N/A	10 mg/mL <sup>5</sup>	discard unused portion <sup>5</sup>	syringe	8 h <sup>6,7</sup>	
				0.4 - 4.8 mg/mL <b>NS</b> , D5W <sup>8</sup>  (e.g., 50-250 mL*)	72 h <b>F</b> , RT <sup>8</sup>	
				0.06 - 0.4 mg/mL <b>NS</b> , D5W <sup>4</sup>	<b>NS:</b> 24 h RT <sup>4</sup>  <b>D5W:</b> 12 h RT <sup>4</sup>	
				0.06 - 1 mg/mL Ringer's, Lactated Ringer's, D10W, D10NS <sup>4</sup>	<b>Ringer's, LR:</b> 24 h RT <sup>4</sup>  <b>D10W:</b> 12 h RT <sup>4</sup>  <b>D10NS:</b> 6 h RT <sup>4</sup>	

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Melphalan</b> 50 mg (GSK) (RT)(PFL) no preservative <sup>9</sup>	10mL supplied diluent <sup>9</sup>  immediately after adding diluent, shake vigorously <sup>9</sup>  record time of reconstitution	5 mg/mL <sup>9</sup>	2 h RT <sup>9</sup>  <b>do NOT refrigerate</b>	0.1 – 0.45 mg/mL in <b>NS</b> only <sup>9</sup>  (e.g., greater than 45 mg and less than or equal to 110 mg in 250 mL NS)*	complete administration within 60 min from time of initial reconstitution at RT <sup>10</sup>	
<b>Mesna</b> 400 mg/4 mL 1000 mg/10 mL (Baxter) (RT) no preservative <sup>11</sup>	N/A	100 mg/mL <sup>11</sup>  (use filter needle to withdraw from ampoule)	discard unused portion <sup>11</sup>	greater than 1 mg/mL in D5W, D5½NS, NS, LR <sup>11-13</sup>	complete administration within 24 h RT <sup>11</sup>	
<b>Mesna</b> 1000 mg/10 mL 5000 mg/50 mL (Baxter) (RT) preservative <sup>11</sup>	N/A	100 mg/mL <sup>11</sup>	8 days RT <sup>11</sup>  (vial may be punctured up to 4 times) <sup>11</sup>	greater than 1 mg/mL in D5W, D5½NS, NS, LR <sup>11-13</sup>	complete administration within 24 h RT <sup>11</sup>	
<b>Mesna</b> 1000 mg/10mL (Fresenius Kabi) (RT) preservative <sup>14</sup>	N/A	100 mg/mL <sup>14</sup>	14 d RT,F <sup>14,15</sup>	greater than or equal to 1 mg/mL in NS or D5W <sup>16</sup>	24 h RT, 48 h F <sup>14</sup>	

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Methotrexate</b> 50 mg/2mL 500 mg/20mL 1 g/40mL (Accord) (RT)(PFL) no preservative <sup>17</sup>	N/A	25 mg/mL <sup>17</sup>	50mg: discard unused portion <sup>17</sup>  500 mg or 1 g: 8 h RT <sup>17</sup>	syringe	use within 8 h RT of initial puncture <sup>17</sup>	- for high-dose regimens (e.g., 1- 12 g/m <sup>2</sup> as a single dose) <sup>18-22</sup> : use preservative-free methotrexate <sup>17</sup> - do not use for IT injection
				0.4–2 mg/mL <b>NS</b> , D5W <sup>17</sup>  (100 mL* <b>NS</b> , D5W)	use within 24 h RT of initial puncture <sup>17</sup>  **(PFL)	
				high dose (e.g., 1-12 g/m <sup>2</sup> as a single dose) <sup>18-22</sup> : 1000 mL* NS	use within 24 h RT of initial puncture <sup>17</sup>  **(PFL)	
<b>Methotrexate IT Injection</b> Only preservative free methotrexate may be administered by the intrathecal route <sup>23</sup> 50 mg/2mL (Accord) (RT)(PFL) no preservative <sup>17</sup>	N/A	25 mg/mL <sup>17</sup>	discard unused portion <sup>17</sup>	qs to 6 mL with preservative free NS <sup>24,25</sup>	use within 4 h of initial puncture <sup>2</sup>	- auxiliary info <sup>2</sup> - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag <sup>26</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Methotrexate</b> 50 mg/2mL 500 mg/20mL (Accord) (RT)(PFL) preservative <sup>17</sup>	N/A	25 mg/mL <sup>17</sup>	14 d F <sup>2,17</sup>	syringe	14 d F <sup>2</sup>	- contains benzyl alcohol <sup>17</sup> - do NOT use for high-dose regimens (e.g., 1-12 g/m <sup>2</sup> as a single dose) <sup>17</sup> - do NOT use for IT injection <sup>17</sup>
				0.4–2 mg/mL <b>NS</b> , D5W <sup>17</sup>  (100 mL* <b>NS</b> , D5W)	24 h RT <sup>17</sup>	
<b>Methotrexate</b> 50 mg/2mL 500 mg/20mL 1 g/40mL 2.5 g/100 mL (Pfizer/Hospira) (RT)(PFL) no preservative <sup>27</sup>	N/A	25 mg/mL <sup>27</sup>	50mg: discard unused portion <sup>27</sup>  500 mg, 1 g, or 2.5 g: 8 h RT <sup>27</sup>	syringe	use within 8 h RT of initial puncture <sup>27</sup>	- for high-dose regimens (e.g., 1-12 g/m <sup>2</sup> as a single dose) <sup>18-22</sup> ; use preservative-free methotrexate <sup>27</sup> - do not use for IT injection
				0.4–2 mg/mL <b>NS</b> , D5W <sup>27</sup>  (100 mL* <b>NS</b> , D5W)	use within 24 h RT of initial puncture <sup>27</sup>  **(PFL)	
				high dose (e.g., 1-12 g/m <sup>2</sup> as a single dose) <sup>18-22</sup> ; 1000 mL* NS	use within 24 h RT of initial puncture <sup>27</sup>  **(PFL)	

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Methotrexate IT Injection</b> Only preservative free methotrexate may be administered by the intrathecal route <sup>23</sup> 50 mg/2mL (Pfizer/Hospira) (RT)(PFL) no preservative <sup>27</sup>	N/A	25 mg/mL <sup>27</sup>	discard unused portion <sup>27</sup>	qs to 6 mL with preservative free NS <sup>24,25</sup>	use within 4 h of initial puncture <sup>15</sup>	- auxiliary info <sup>2</sup> : "IT" - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag <sup>26</sup>
<b>Methotrexate</b> 50 mg/2mL 500 mg/20mL (Pfizer/Hospira) (RT)(PFL) preservative <sup>27</sup>	N/A	25 mg/mL <sup>27</sup>	14 d F <sup>15,27</sup>	syringe	14 d F <sup>15</sup>	- contains benzyl alcohol <sup>27</sup> - do NOT use for high-dose regimens (e.g., 1- 12 g/m <sup>2</sup> as a single dose) <sup>27</sup> - do NOT use for IT injection <sup>27</sup>
				0.4–2 mg/mL <b>NS</b> , D5W <sup>27</sup>  (100 mL * <b>NS</b> , D5W)	24 h RT <sup>27</sup>	
<b>Mitomycin</b> 20 mg (Accord) (RT)(PFL) no preservative <sup>28</sup>	40 mL SWI <sup>28</sup>  shake well <sup>28</sup>	0.5 mg/mL <sup>28</sup>	6 h RT, 72 h F <sup>28</sup>  **(PFL) <sup>28</sup>	syringe	6 h RT, 72 h F <sup>28</sup>  **(PFL) <sup>28</sup>	

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Mitomycin intravesical</b> 20 mg (Accord) (RT)(PFL) no preservative <sup>28</sup>	40 mL SWI <sup>28</sup> shake well <sup>28</sup>	0.5 mg/mL <sup>28</sup>	6 h RT, 72 h F <sup>28</sup>  **(PFL) <sup>28</sup>	syringe	6 h RT, 72 h F <sup>28</sup>  **(PFL) <sup>28</sup>	
	10 mL SWI <sup>29</sup> shake well <sup>28</sup>	2 mg/mL <sup>29</sup>	use immediately after preparation to prevent precipitation <sup>30</sup>	syringe	use immediately after preparation to prevent precipitation <sup>30</sup>	- may precipitate due to low solubility <sup>30,31</sup> - do NOT refrigerate <sup>30</sup>
<b>Mitomycin intraperitoneal</b> 20 mg (Accord) (RT)(PFL) no preservative <sup>28</sup>	40 mL SWI <sup>28</sup> shake well <sup>28</sup>	0.5 mg/mL <sup>28</sup>	6 h RT, 72 h F <sup>28</sup>  **(PFL) <sup>28</sup>	0.02-0.04 mg/mL <sup>28</sup>  <b>NS</b> , sodium lactate <sup>28</sup>	NS: 3 h RT, 18 h F <sup>28</sup>  sodium lactate: 3 h RT, 6 h F <sup>28</sup>	
<b>Mitomycin</b> 20 mg (Teva/Novopharm) (RT)(PFL) no preservative <sup>32</sup>	40 mL SWI <sup>32</sup> shake well <sup>32</sup>	0.5 mg/mL <sup>32</sup>	6 h RT, 72 h F <sup>32</sup>  **(PFL) <sup>32</sup>	syringe	6 h RT, 72 h F <sup>32</sup>  **(PFL) <sup>32</sup>	

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Mitomycin intravesical</b> 20 mg (Teva/Novopharm) (RT)(PFL) no preservative <sup>32</sup>	40 mL SWI <sup>32</sup> shake well <sup>32</sup>	0.5 mg/mL <sup>32</sup>	6 h RT, 72 h F <sup>32</sup>  **(PFL) <sup>32</sup>	syringe	6 h RT, 72 h F <sup>32</sup>  **(PFL) <sup>32</sup>	
	10 mL SWI <sup>29</sup> shake well <sup>32</sup>	2 mg/mL <sup>29</sup>	use immediately after preparation to prevent precipitation <sup>30</sup>	syringe	use immediately after preparation to prevent precipitation <sup>30</sup>	- may precipitate due to low solubility <sup>30,31</sup> - do NOT refrigerate <sup>30</sup>
<b>Mitomycin intraperitoneal</b> 20 mg (Teva/Novopharm) (RT)(PFL) no preservative <sup>32</sup>	40 mL SWI <sup>32</sup> shake well <sup>32</sup>	0.5 mg/mL <sup>32</sup>	6 h RT, 72 h F <sup>32</sup>  **(PFL) <sup>32</sup>	0.02-0.04 mg/mL <sup>32</sup>  <b>NS</b> , sodium lactate <sup>32</sup>	NS: 6 h RT, 18 h F <sup>32</sup>  sodium lactate: 6 h RT, F <sup>32</sup>	
<b>mitoXANTRONE</b> 20 mg/10 mL (Fresenius Kabi) (RT) no preservative <sup>33</sup>	N/A	2 mg/mL <sup>33</sup>	discard unused portion <sup>33</sup>	<b>NS</b> , D5W <sup>33</sup>  Greater than or equal to *50 mL <sup>33</sup>	24 h RT <sup>33</sup>	



**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>mitoXANTRONE</b> 20 mg/10 mL 25 mg/12.5 mL (Hospira) (RT)(PFL) no preservative <sup>34</sup>	N/A	2 mg/mL <sup>34</sup>	discard unused portion <sup>34</sup>	0.2-0.6 mg/mL <sup>34</sup>  <b>NS, D5W<sup>34</sup></b>  Greater than or equal to *50 mL <sup>34</sup>	NS: 24 h F, RT <sup>34</sup>  **(PFL) <sup>34</sup>	
<b>mitoXANTRONE</b> 20 mg/10 mL (Teva/Novopharm) (RT)(PFL) no preservative <sup>35</sup>	N/A	2 mg/mL <sup>35</sup>	discard unused portion <sup>35</sup>	Greater than or equal to *50 mL <b>NS, D5W<sup>35</sup></b>	24 h RT <sup>35</sup>  **(PFL) <sup>36</sup>	
<b>Nivolumab</b> 40 mg/4 mL 100 mg/10 mL (BMS) (F)(PFL) do not shake no preservative <sup>37</sup>	N/A	10 mg/mL <sup>37</sup>	discard unused portion <sup>37</sup>	1-10 mg/mL <b>NS,</b> <b>D5W<sup>37</sup></b>  (50-100* mL)  mix by gentle inversion; do not shake <sup>37</sup>	complete administration within 8 h RT or 24 h F <sup>37</sup>  **(PFL) <sup>37</sup>	- administer with a 0.2 to 1.2 micron in-line filter <sup>37</sup> - discard if cloudy or has pronounced colour change (should be clear to pale yellow) <sup>37</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>oBINutuzumab</b> 1000 mg/40 mL (Hoffman-La Roche) (F)(PFL) do not shake no preservative <sup>38</sup>	N/A	25 mg/mL <sup>38</sup>	discard unused portion <sup>39</sup>	100 mg: in 100 mL <b>NS</b> <sup>38</sup>  900 mg: in 250 mL <b>NS</b> <sup>38</sup>  1000 mg: in 250 mL <b>NS</b> <sup>38</sup>	<b>24 h F</b> , 48 h RT <sup>38,40</sup>	-once removed from the fridge, diluted product is stable for an additional 48 h RT <sup>38,40</sup> - do NOT shake <sup>38</sup> - do NOT use dextrose containing solutions <sup>38</sup>
<b>Octreotide</b> 50 mcg/mL 100 mcg/mL 500 mcg/mL (Omega) (F)(PFL) no preservative <sup>41</sup>	N/A	50 mcg/mL <sup>41</sup>  100 mcg/mL <sup>41</sup>  500 mcg/mL <sup>41</sup>	Use within 4 h <sup>41</sup>	<b>NS</b> <sup>41</sup>  volume adjusted to ensure a continuous infusion of octreotide at 25 mcg/hour <sup>41</sup>	24 h RT <sup>41</sup>	
<b>Octreotide</b> multidose vial: 1000 mcg/5 mL (Omega) (F)(PFL) preservative <sup>41</sup>	N/A	200 mcg/mL <sup>41</sup>	15 d F <sup>41</sup>	<b>NS</b> <sup>41</sup>  volume adjusted to ensure a continuous infusion of octreotide at 25 mcg/hour <sup>41</sup>	24 h RT <sup>41</sup>	

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Octreotide</b> 50 mcg/mL 100 mcg/mL 500 mcg/mL (Teva/Novopharm) (F)(PFL) no preservative <sup>42</sup>	N/A	50 mcg/mL <sup>42</sup>  100 mcg/mL <sup>42</sup>  500 mcg/mL <sup>42</sup>	discard unused portion <sup>42</sup>	SC syringe <sup>42</sup>	use within 4 h <sup>42</sup>	
				infusion: NS <sup>42</sup>	24 h RT <sup>42</sup>	
<b>Octreotide</b> multidose vial: 1000 mcg/5 mL (Teva/Novopharm) (F)(PFL) preservative <sup>42</sup>	N/A	200 mcg/mL <sup>42</sup>	14 d F <sup>42,43</sup>	SC syringe	use within 14 d F <sup>42,43</sup>	
				infusion: NS <sup>42</sup>	24 h RT <sup>42</sup>	
<b>Octreotide</b> (SANDOSTATIN®) 1000 mcg/5 mL (Novartis) (F)(PFL) preservative <sup>44</sup>	N/A	200 mcg/mL <sup>44</sup>	discard unused portion <sup>45</sup>	50–200 mL NS <sup>10,46,47</sup>  SC infusion: adjust volume to ensure infusion rate of 25 mcg/h <sup>46</sup>	24 h RT <sup>46</sup>	

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Octreotide</b> (SANDOSTATIN®) 50 mcg/1 mL 100 mcg/1 mL 500 mcg/1 mL (Novartis) (F)(PFL) no preservative <sup>44</sup>	N/A	50 mcg/mL 100 mcg/mL 500 mcg/mL <sup>44</sup>	discard unused portion <sup>46</sup>	50-100 mL <sup>10,47</sup>  NS <sup>46</sup>  SC infusion: adjust volume to ensure infusion rate of 25 mcg/h <sup>46</sup>	24 h RT <sup>46</sup>	
<b>Octreotide</b> (SANDOSTATIN LAR®) 10 mg 20 mg 30 mg (Novartis) (F)(PFL) no preservative <sup>45</sup>	2 mL supplied diluent  gently run 2 mL down sides of the vial; do NOT disturb for 2–5 min, then swirl moderately <sup>46</sup>  record time of reconstitution	10 mg: 5 mg/mL  20 mg: 10 mg/mL  30 mg: 15 mg/mL <sup>46</sup>	discard unused portion <sup>46</sup>	deep intragluteal administration only <sup>46</sup>	use within 4 h of initial reconstitution <sup>7,46</sup>	- do NOT shake

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p><b>Olaratumab</b> 500 mg/50 mL (Lilly) (F)(PFL) do not shake no preservative<sup>48</sup></p>	<p>N/A</p>	<p>10 mg/mL<sup>48</sup></p>	<p>discard unused portion<sup>39,48</sup></p>	<p>dilute to a final volume of 250 mL NS<sup>48</sup></p> <p>do NOT use D5W or other dextrose containing solutions<sup>48</sup></p> <p>gently invert to mix<sup>48</sup></p>	<p>complete administration within 24 h F, plus an additional 12 h RT<sup>48</sup></p>	<p>- do NOT shake<sup>48</sup></p>
<p><b>Oxaliplatin</b> 50 mg/10 mL 100 mg/20 mL 200 mg/40 mL (Hospira/Pfizer) (RT) no preservative<sup>49</sup></p>	<p>N/A</p>	<p>5 mg/mL<sup>49</sup></p>	<p>discard unused portion<sup>49</sup></p>	<p>250-500 mL D5W (0.2-0.7 mg/mL<sup>49</sup></p> <p>do NOT use NS or other chloride- containing solutions<sup>50</sup></p> <p>do NOT use aluminum-containing needle and syringe<sup>50</sup></p>	<p>0.2-0.4 mg/mL: 24 h RT or 5 d F plus an additional 8 h RT<sup>49,51</sup></p> <p>0.5–2 mg/mL: 24 h RT or 14 d F plus an additional 8 h RT<sup>49,51</sup></p>	<p>- do NOT use aluminum- containing needle, syringe or tubing<sup>49</sup></p>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Oxaliplatin</b> 50 mg/10 mL 100 mg/20 mL 150 mg/30 mL 200 mg/40 mL (Sandoz) (RT)(PFL) no preservative <sup>50</sup>	N/A	5 mg/mL <sup>50</sup>	2 d F, RT <sup>39,52</sup>	0.2-0.7 mg/mL <sup>50</sup> 250-500 mL D5W <sup>50</sup> do NOT use NS or other chloride- containing solution <sup>50</sup>  do NOT use aluminum-containing needle and syringe <sup>50</sup>	0.2-2 mg/mL: 24 h RT, 48 h F <sup>50</sup>	
<b>Oxaliplatin</b> 50 mg/10 mL 100 mg/20 mL 200 mg/40 mL (Teva) (RT)(PFL) no preservative <sup>53</sup>	N/A	5 mg/mL <sup>53</sup>	discard unused portion <sup>53</sup>	250-500 mL D5W (0.2-0.7 mg/mL) <sup>53</sup> do <b>NOT</b> use NS or other chloride- containing solution <sup>53</sup>  do NOT use aluminum-containing needle and syringe <sup>53</sup>	0.2-2 mg/mL: 24 h RT, 48 h F <sup>53</sup>	- do NOT use aluminum- containing needle, syringe or tubing <sup>53</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>PACLitaxel</b> 30 mg/5 mL 100 mg/16.7 mL 300 mg/50 mL (Accord) (RT)(PFL) no preservative <sup>54</sup>	N/A	6 mg/ mL <sup>54</sup>	30 mg: 48 h RT <sup>39,54</sup>  100 mg: 48 h RT <sup>39,54</sup>  300 mg: 24 h RT <sup>54</sup>	0.3-1.2 mg/mL in <b>NS</b> , D5W, D5NS, D5LR <sup>54</sup>  (e.g., 100-1000 mL)*	complete administration within 27 h RT <sup>54</sup>	- use non-DEHP bag and tubing with 0.22 micron in-line filter <sup>54</sup> - avoid excessive shaking <sup>54</sup>
<b>PACLitaxel</b> 30 mg/5 mL 100 mg/16.7 mL 300 mg/50 mL (Biolyse) (RT) <sup>55</sup> no preservative <sup>56</sup>	N/A	6 mg/mL <sup>56</sup>	48 h RT <sup>39,57</sup>	0.3-1.2 mg/mL in <b>NS</b> , D5W <sup>56</sup>  (e.g., 100-1000 mL)*	complete administration within 27 h RT <sup>56,58</sup>	- use non-DEHP bag and tubing with 0.22 micron in-line filter <sup>56</sup>
				0.1 mg/mL in <b>NS</b> <sup>59</sup>	44 h <b>F</b> , RT <sup>59</sup>	
				0.012-0.12 mg/mL in <b>NS</b> <sup>60</sup>	16 h RT <sup>58</sup>	
				devices with spikes (e.g., chemo dispensing pins) <b>may be used</b> with vials <sup>61</sup>		

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>PACLitaxel</b> 30 mg/5 mL 100 mg/16.7 mL 150 mg/25 mL 300 mg/50 mL (Hospira) (RT)(PFL) preservative <sup>62</sup>	N/A	6 mg/mL <sup>63</sup>	48 h RT <sup>39,63,64</sup>	0.3-1.2 mg/mL in <b>NS</b> , D5W, D5NS, D5LR <sup>63</sup>  (e.g., 100-1000 mL)*	complete administration within 27 h RT <sup>63</sup>	- use non-DEHP bag and tubing with 0.22 micron in-line filter <sup>63</sup>
<b>PACLitaxel, nanoparticle, albumin- bound (nab)</b> 100 mg (Celgene) (RT)(PFL) no preservative <sup>65</sup>	20 mL NS <sup>65</sup>  - slowly direct diluent against side of vial (i.e., greater than or equal to 1 min) during reconstitution <sup>65</sup>  - let stand for greater than or equal to 5 min to wet powder <sup>65</sup>  - gently swirl or invert for greater than or equal to 2 min <sup>65</sup>	5 mg/mL <sup>65</sup>	use immediately (RT) or 8 h F <sup>65</sup>  **(PFL) <sup>65</sup>	in empty sterile PVC, non-PVC, or non- DEHP infusion bag <sup>65</sup>	48 h F plus an additional 8 h RT <sup>66</sup>	- each vial contains 900 mg human albumin <sup>65</sup> - to prevent foaming, do NOT inject NS directly onto the powder <sup>65</sup> - some settling may occur; use mild agitation to resuspend <sup>65</sup> - administer using a 15 micron filter ONLY (NOTE:filters with a pore size less than 15 microns may cause filter blockage) <sup>67,68</sup>



**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Pamidronate</b> 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Fresenius Kabi) (RT) no preservative <sup>69</sup>	N/A	3 mg/mL <sup>69</sup>	discard unused portion <sup>69</sup>	Less than or equal to 0.36 mg/mL <sup>69</sup> <b>NS</b> , D5W <sup>69</sup>	24 h RT <sup>69</sup>	- do <b>NOT</b> mix with calcium containing solutions <sup>69</sup>
		6 mg/mL <sup>69</sup>				
		9 mg/mL <sup>69</sup>				
<b>Pamidronate</b> 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Hospira) (RT) no preservative <sup>70</sup>	N/A	3 mg/mL <sup>70</sup>	discard unused portion <sup>70</sup>	0.06–0.36 mg/mL in <b>NS</b> , D5W <sup>70</sup>	24 h F followed by 24 h RT (total 48 h) <sup>70</sup>  **(PFL) <sup>70</sup>	- do <b>NOT</b> mix with calcium containing solution (e.g., Ringer's) <sup>70</sup>
		6 mg/mL <sup>70</sup>				
		9 mg/mL <sup>70</sup>				
<b>Pamidronate</b> 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Omega) (RT) no preservative <sup>71</sup>	N/A <sup>71</sup>	3 mg/mL <sup>71</sup>	discard unused portion <sup>71</sup>	0.06–0.36 mg/mL in <b>NS</b> , D5W <sup>71</sup>	24 h F followed by 24 h RT (total 48 h) <sup>71</sup>  **(PFL) <sup>71</sup>	- do <b>NOT</b> mix with calcium containing solution (e.g., Ringer's) <sup>71</sup>
		6 mg/mL <sup>71</sup>				
		9 mg/mL <sup>71</sup>				

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Pamidronate</b> 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Pfizer) (RT) no preservative <sup>72</sup>	N/A	3 mg/mL <sup>72</sup>	discard unused portion <sup>72</sup>	0.06-0.36 mg/mL in <b>NS</b> , D5W <sup>72</sup>	24 h F followed by 24 h RT (total 48 h) <sup>72</sup>  **(PFL) <sup>72</sup>	- do <b>NOT</b> mix with calcium containing solution (e.g., Ringer's) <sup>72</sup>
		6 mg/mL <sup>72</sup>				
		9 mg/mL <sup>72</sup>				
<b>Pamidronate</b> 30 mg/10 mL 60mg/10 mL 90 mg/10 mL (Sandoz Canada) RT no preservative <sup>73</sup>	N/A	3 mg/mL <sup>73</sup>	discard unused portion <sup>43,73</sup>	<b>NS</b> ; D5W <sup>73</sup>	24 h RT <sup>73</sup>	- do <b>NOT</b> mix with calcium containing solution (e.g., Ringer's) <sup>73</sup>
		6 mg/mL <sup>73</sup>				
		9 mg/mL <sup>73</sup>				
<b>PANitumumab</b> 100 mg/5 mL 400 mg/20 mL (Amgen) (F)(PFL) do not shake no preservative <sup>74</sup>	N/A	20 mg/mL <sup>74</sup>	discard unused portion <sup>74</sup>	Less than or equal to 1000 mg: 100 mL NS <sup>74</sup>  Greater than 1000mg: 150 mL NS <sup>74</sup>  1-10mg/mL <sup>74,75</sup>	24 h F, 6 h RT <sup>74,75</sup>	- administer with 0.2 or 0.22 micron in-line filter <sup>74</sup> - solution may contain particulates which do not affect product quality <sup>74</sup> - do not administer if discoloured <sup>74</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>pegaspargase</b> (pegylated asparaginase <i>E. coli</i> ) 3750 units/5 mL (Shire) (F)(PFL) do not shake no preservative <sup>76</sup>	N/A	750 units/mL <sup>76</sup>	discard unused portion <sup>76</sup>	IM: max volume: 2 mL in children and adolescents; 3 mL in adults  if volume greater than above, use multiple sites <sup>76</sup>	syringe: use within 4 h of vial puncture <sup>39,76</sup>	- do NOT shake <sup>76</sup>
				IV: 100 mL <b>NS</b> , D5W <sup>76</sup>	bag: use within 4 h of vial puncture <sup>39,76</sup>	
<b>Pembrolizumab</b> 100 mg/4 mL (Merck) (F)(PFL) do not shake no preservatives <sup>77</sup>	N/A	25 mg/mL <sup>77</sup>	discard unused portion <sup>39,77</sup>	1-10 mg/mL <b>NS</b> , D5W <sup>77</sup>  mix by gentle inversion <sup>77</sup>	complete administration within 6 h RT, 24 h F <sup>77</sup>	- use a 0.2 to 5 micron in-line filter <sup>77</sup> - allow vials and diluted solutions to come to RT prior to use <sup>77</sup> - vials contain 0.25 mL overfill <sup>77</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Pembrolizumab</b> 50 mg (Merck) (F) no preservative <sup>77</sup>	2.3 mL SWI <sup>77</sup>  direct diluent against side of vial during reconstitution to avoid foaming <sup>77</sup>  allow up to 5 minutes for bubbles to clear <sup>77</sup>  do NOT shake <sup>77</sup>	25 mg/mL <sup>77</sup>	6 h RT, 24 h F <sup>77</sup>	1-10 mg/mL NS, D5W <sup>77</sup>  mix by gentle inversion <sup>77</sup>	complete administration within 6 h RT, 24 h F <sup>77</sup>	- use 0.2 to 5 micron in-line filter <sup>78</sup> - allow reconstituted vials and diluted solutions to come to RT prior to use <sup>77</sup> - vials can be at RT for up to 24 h prior to use <sup>77</sup> - vials contain 20% overfill <sup>77</sup>
<b>Pemetrexed</b> 100 mg 500 mg (Accord) (RT) no preservative <sup>79</sup>	100 mg: 4.2 mL NS <sup>79</sup>  500 mg: 20 mL NS <sup>79</sup>	25 mg/mL <sup>79</sup>	24 h F, RT <sup>79</sup>	100 mL NS <sup>79</sup>	24 h F, RT <sup>79</sup>	- do <b>NOT</b> mix with calcium containing solution (e.g., Ringer's) <sup>79</sup>
<b>Pemetrexed</b> 100 mg 500 mg (Eli Lilly) (RT) no preservative <sup>80</sup>	100 mg: 4.2 mL NS <sup>80</sup>  500 mg: 20 mL preservative- free NS <sup>80</sup>	25 mg/mL <sup>80</sup>	24 h F, RT <sup>80</sup>	100 mL NS <sup>80</sup>	24 h F, RT <sup>80</sup>	- do <b>NOT</b> mix with calcium containing solution (e.g., Ringer's) <sup>81</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>PERTuzumab</b> 420 mg/14 mL (Roche) (F)(PFL) no preservative <sup>82</sup>	N/A	30 mg/mL <sup>82</sup>  do NOT shake <sup>82</sup>	discard unused portion <sup>39,82</sup>	250 mL NS only <sup>82</sup>  mix by gentle inversion to avoid foaming <sup>82</sup>	24 h F, RT <sup>82</sup>	- do NOT use dextrose containing solutions <sup>82</sup>
<b>Plerixafor</b> 24 mg/1.2 mL (sanofi-aventis) (RT) no preservative <sup>83</sup>	N/A	20 mg/mL <sup>83</sup>	discard unused portion <sup>83</sup>	SC syringe <sup>83</sup>	48 hours RT <sup>43,84</sup>	
<b>Porfimer</b> 15 mg 75 mg (Axcan) (RT)(PFL) no preservative <sup>85</sup>	15 mg: 6.6 mL D5W <sup>85</sup>  75 mg: 31.8 mL D5W <sup>85</sup>  record time of reconstitution	2.5 mg/mL <sup>85</sup>	24 h F  **(PFL) <sup>85</sup>	syringe <sup>85</sup>	use within 4 h of initial reconstitution <sup>7,86</sup>  **(PFL) <sup>85</sup>	- avoid contact with skin and eyes; protect exposed area from light <sup>85</sup>
<b>Raltitrexed</b> 2 mg (Pfizer) (F,RT)(PFL) (no preservative) <sup>87</sup>	4 mL SWI <sup>87</sup>	0.5 mg/mL <sup>87</sup>	24 h F, RT <sup>87</sup>	50-250 mL NS, D5W <sup>87</sup>	complete administration within 24 h F, RT <sup>87</sup>	

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Ramucirumab</b> 100 mg/10 mL 500 mg/50 mL (Eli Lilly) (F)(PFL) (do not shake) no preservative <sup>88</sup>	N/A	10 mg/mL <sup>88</sup>	discard unused portion <sup>88</sup>	250 mL* NS <sup>88</sup> (0.4 – 4 mg/mL) <sup>89</sup> gently invert to mix <sup>88</sup> do NOT shake <sup>88</sup>	4 h RT, 24 h F <sup>88</sup>	- use 0.22 micron filter <sup>88</sup> - do NOT use dextrose containing solutions <sup>88</sup>
<b>riTUXimab</b> 100 mg/10 mL 500 mg/50 mL (Roche) (F)(PFL) no preservative <sup>90</sup>	N/A	10 mg/mL <sup>90</sup>	discard unused portion <sup>90</sup>	1-4 mg/mL <b>NS</b> , D5W <sup>90</sup> (e.g., 250-500 mL)*	24 h F, 12 h RT <sup>91,92</sup>	- once removed from the fridge, compounded product is stable for 12h RT <sup>91,92</sup>
<b>riTUXimab subcutaneous</b> 1400 mg/11.7 mL 1600 mg/13.4 mL (Roche) (F)(PFL) no preservative <sup>93</sup>	N/A	120 mg/mL <sup>93</sup>	discard unused portion <sup>93</sup>	SC syringe <sup>93</sup>	48 h F plus 8 h RT <sup>93</sup>	- contains hyaluronidase <sup>93</sup> - formulations are NOT interchangeable <sup>93</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>romiDEPsin</b> 10 mg (Celgene Inc.) (RT) <sup>94</sup> no preservative <sup>39</sup>	2.2 mL of supplied diluent <sup>94,95</sup>  swirl gently to mix <sup>94</sup>	5 mg/mL <sup>94</sup>	8 h RT <sup>94</sup>	500 mL NS <sup>94</sup>	24 h RT <sup>94</sup>	- reconstituted solution will be slightly viscous <sup>96</sup> - vials contain overfill to allow for full drug recovery (drug vial contains 11 mg romidepsin; diluent vial contains 2.4 mL diluent) <sup>94</sup>
<b>Siltuximab</b> 100 mg 400 mg (Janssen) (F)(PFL) no preservative <sup>97</sup>	100 mg: 5.2 mL SWI <sup>97</sup>  400 mg: 20 mL SWI <sup>97</sup>  allow vial to come to room temperature prior to use (~30 minutes) <sup>97</sup>  gently swirl, do NOT shake <sup>97</sup>	20 mg/mL <sup>97</sup>	2 h RT <sup>97</sup>	250 mL <b>D5W</b> <sup>97</sup>  dilute to 250 mL final volume by withdrawing volume from bag equal to volume of drug to be added <sup>97</sup>	complete administration within 6 h RT <sup>97</sup>	- use 0.2 micron in-line filter <sup>97</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Streptozocin</b> 1g (Pfizer) (F)(PFL) no preservative <sup>98</sup>	9.5mL <b>NS</b> , SWI, D5W <sup>98</sup>	100 mg/mL <sup>98</sup>	48 h F, <sup>98</sup> 24 h RT	syringe <sup>98</sup>	48 h F, 24 h RT <sup>98</sup>	
				50-500 mL * <b>NS</b> , D5W, SWI <sup>98</sup>	48 h F, 24 h RT <sup>98</sup>	
<b>Temsirolimus</b> 30 mg/1.2 mL (Wyeth) (F)(PFL) <sup>99,100</sup> no preservative <sup>101</sup>	1.8 mL supplied diluent <sup>99,100</sup>	10 mg/mL <sup>99,100</sup>	24 h RT <sup>99,100</sup>  **(PFL) <sup>99</sup>	250 mL NS <sup>99,100</sup>	complete administration within 6 h <sup>99,100</sup>	- use non-DEHP bag and tubing with in-line filter <sup>99,100</sup>
<b>Teniposide</b> 50 mg/5 mL (BMS) (RT) preservative <sup>102</sup>	N/A	10 mg/mL <sup>102</sup>	discard unused portion	50 – 500 mL <b>NS</b> or D5W for a final concentration of 0.1-1 mg/mL <sup>102</sup>	0.1-0.4 mg/mL: 24 h RT <sup>102</sup>  1 mg/mL: complete administration within 4 h of preparation RT <sup>102,103</sup>	- do not refrigerate - use non-DEHP bag and tubing <sup>102</sup> - do not use if precipitates <sup>102,103</sup> - contains DMA*** - excessive agitation may cause precipitation <sup>102</sup>



**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p><b>Thiotepa</b> 15 mg 100 mg (Adienne/Methapharm) (F) no preservative<sup>104</sup></p>	<p><b>15 mg:</b> 1.5 mL SWI<sup>104</sup></p> <p><b>100 mg:</b> 10 mL SWI<sup>104</sup></p> <p>to remove haze, filter through 0.22 micron filter after reconstitution<sup>105</sup></p> <p>record time of reconstitution</p>	<p>10 mg/mL<sup>104</sup></p>	<p>8 h F<sup>104</sup></p>	<p>reconstituted solution is hypotonic and must be further diluted with NS prior to use<sup>104</sup></p> <p>doses ≤ 500 mg: 500 mL NS or with an appropriate volume to achieve 0.5-1 mg/mL concentration<sup>104</sup></p> <p>doses &gt; 500 mg: 1000 mL NS<sup>104</sup></p>	<p>4 h RT, 24 h F<sup>104</sup></p>	<p>- do not use if precipitates are present<sup>104</sup> - reconstituted solution may be used if opalescent<sup>104</sup> - administer with 0.2 micron inline filter<sup>104</sup></p>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p><b>Thiotepa IT injection</b> 15 mg 100mg (Adienne/Methapharm) (F) no preservative<sup>104</sup></p>	<p>diluents containing preservatives should <b>NOT</b> be used for intrathecal administration<sup>106</sup></p> <p><b>15 mg:</b> 1.5 mL SWI<sup>104</sup></p> <p><b>100 mg:</b> 10 mL SWI<sup>104</sup></p> <p>to remove haze, filter through 0.22 micron filter after reconstitution<sup>105</sup></p> <p>record time of reconstitution</p>	<p>10 mg/mL<sup>104</sup></p>	<p>8 h F<sup>104</sup></p>	<p>qs to 6 mL with preservative free NS<sup>25</sup></p>	<p>use within 4 h of initial reconstitution<sup>2</sup></p>	<p>- auxiliary info<sup>26</sup>: "IT" - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag<sup>26</sup> - do not use if precipitates are present<sup>104</sup> - reconstituted solution may be used if opalescent<sup>104</sup></p>
<p><b>Thyrotropin alfa</b> 1.1 mg (Genzyme) (F)(PFL) no preservative<sup>107</sup></p>	<p>1.2 mL SWI<sup>107</sup></p> <p>swirl contents<sup>107</sup></p> <p>do NOT shake</p>	<p>0.9 mg/mL<sup>107</sup></p>	<p>24 h F<sup>107</sup></p>	<p>syringe<sup>107</sup></p>	<p>24 h F<sup>107</sup></p>	

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Tocilizumab</b> 80 mg/4 mL 200 mg/10 mL 400 mg/20 mL (Roche) (F)(PFL) no preservative <sup>108</sup>	N/A	20 mg/mL <sup>108</sup>	discard unused portion <sup>108</sup>	100 mL NS <sup>108</sup>  dilute to final volume by withdrawing volume from bag equal to volume of drug to be added <sup>108</sup>  gently invert to mix <sup>108</sup>	complete administration within 24 h F, RT <sup>108</sup>  bring to room temperature prior to administration <sup>108</sup>	- to prevent foaming: slowly add drug to infusion bag and gently invert bag to mix <sup>108</sup>
<b>Topotecan</b> 4 mg/4 mL (Accord) (RT)(PFL) no preservative <sup>109</sup>	N/A	1 mg/mL <sup>109</sup>	discard unused portion <sup>2,109</sup>	0.025-0.5 mg/mL  50-100 mL NS, D5W <sup>109</sup>	14 d F, 48 h RT <sup>2,109</sup>	
<b>Topotecan</b> 1 mg 4 mg (Actavis) (RT)(PFL) no preservative <sup>110</sup>	<b>1 mg:</b> 1.1 mL SWI <sup>110</sup>  <b>4 mg:</b> 4 mL SWI <sup>110</sup>	1 mg/mL <sup>110</sup>	24 h F, RT <sup>110</sup>	0.02-0.5 mg/mL  50-100 mL NS, D5W <sup>110</sup>	24 F, RT <sup>110</sup>	
<b>Topotecan</b> 4 mg/4 mL (Pfizer/Hospira) (F)(PFL) no preservative <sup>111</sup>	N/A	1 mg/mL <sup>111</sup>	discard unused portion <sup>2,111</sup>	0.02-0.5 mg/mL  50-100 mL NS, D5W <sup>111</sup>	24 h F, RT <sup>111</sup>	

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Topotecan</b> 4 mg/4 mL (Sandoz) (F)(PFL) no preservative <sup>112</sup>	N/A	1 mg/mL <sup>112</sup>	discard unused portion <sup>112</sup>	0.02-0.5 mg/mL  50-100 mL <b>NS</b> , D5W <sup>112</sup>	24 h F <sup>112</sup>  **(PFL) <sup>112</sup>	
<b>Trastuzumab (HERCEPTIN®)</b> 440 mg (Roche) (F) preservative <sup>113</sup>	20 mL supplied BWI <sup>113</sup>  swirl vial gently; allow to stand undisturbed for 5 min <sup>113</sup>	21 mg/mL <sup>113</sup>	14 d F <sup>39</sup>	250 mL <b>NS</b> only <sup>113</sup>  do NOT use dextrose containing solutions <sup>113</sup>	24 h <b>F</b> , RT <sup>113</sup>	- do NOT shake <sup>113</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<p><b>Trastuzumab Emtansine (KADCYLA®)</b> 100 mg 160 mg (Roche) (F)(PFL) no preservative<sup>114</sup></p>	<p>100 mg vial: 5 mL SWI<sup>114</sup></p> <p>160 mg vial: 8 mL SWI<sup>114</sup></p> <p>swirl gently until completely dissolved</p> <p>do NOT shake<sup>114</sup></p>	<p>20 mg/mL<sup>114</sup></p>	<p>24 h F<sup>114</sup></p> <p>do NOT freeze<sup>114</sup></p>	<p>250 mL <b>NS</b> or 0.45% sodium chloride only<sup>114</sup></p> <p>do NOT shake<sup>114</sup></p>	<p>24 h F<sup>114</sup></p> <p>do NOT freeze<sup>114</sup></p>	<p>- do not use if reconstituted solution contains visible particulates or is cloudy or discolored<sup>114</sup></p> <p>- dextrose 5% solutions cause aggregation of the protein; do not dilute with dextrose containing solutions<sup>114</sup></p> <p>- use a 0.2 micron in-line filter or 0.22 micron polyethersulfane (PES) filter to administer infusions prepared in NS; filter is optional for solutions in 0.45% NS<sup>114</sup></p>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>TRC105 (Carotuximab)</b> 100 mg/4 mL 200 mg/8 mL 400 mg/16 mL (Tracon) (F)(PFL) no preservative <sup>115</sup>	N/A	25 mg/mL <sup>115</sup>	discard unused portion <sup>39</sup>	0.6 – 10 mg/mL NS <sup>116</sup>  invert gently to mix	complete infusion within 8 h RT, 24 h F <sup>115,116</sup>	- use a 0.2 micron in-line filter for administration <sup>115</sup>
<b>Treosulfan</b> 1 g 5 g (medac) (RT) no preservative <sup>117</sup>	<b>pre-heat</b> SWI to 30°C (not higher) <b>shake vial</b> carefully before adding the warmed SWI <b>1 g vial:</b> 20 mL SWI, while slightly shaking vial and syringe; continue shaking the reconstituted solution for another 2 min <sup>117</sup> <b>5 g vial:</b> 100 mL SWI, while slightly shaking vial and syringe; continue shaking the reconstituted solution for another 2 min <sup>117</sup>	50 mg/mL <sup>117</sup>	48 h RT <sup>7,117</sup>	undiluted <sup>118</sup>  dilute with <b>NS</b> or D5W in empty infusion bag for final concentration = 20 mg/mL <sup>117</sup>	48 h RT <sup>7,117</sup>	- compatible with polytetrafluoroethyl ene filters <sup>117</sup> - may require vigorous shaking to reconstitute <sup>117</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>vinBLAStine</b> 10 mg/10 mL (Hospira) (F)(PFL) no preservative <sup>119</sup>	N/A	1 mg/mL <sup>119</sup>	discard unused portion <sup>119</sup>	25-50 mL <b>NS</b> , D5W <sup>120</sup>	24 h <b>F</b> , RT <sup>121,122</sup>	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES <sup>123,124</sup>
<b>vinBLAStine</b> 10 mg/10 mL (Teva) (F)(PFL) no preservative <sup>125</sup>	N/A	1 mg/mL <sup>125</sup>	discard unused portion <sup>125</sup>	25-50 mL <b>NS</b> , D5W <sup>120,126</sup>	use within 4 h of initial puncture <sup>39</sup>	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES <sup>123,124</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>vinCRISTine</b> 2 mg/2 mL 5 mg/5 mL (Hospira) (F)(PFL) no preservative <sup>127</sup>	N/A	1 mg/mL <sup>127</sup>	8 h F, RT <sup>127</sup>	50 mL * NS, D5W <sup>127</sup>	24 h F, 6 h RT <sup>127</sup>  **(PFL) <sup>127</sup>	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES <sup>123,124</sup> - for ULYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing etoposide, DOXOrubicin, vinCRISTine)



**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>vinCRISTine</b> 1 mg/1 mL 2 mg/2 mL 5 mg/5 mL (Teva) (F)(PFL) no preservative <sup>128</sup>	N/A	1 mg/mL <sup>128</sup>	8 h F, RT <sup>128</sup>	0.01-0.1 mg/mL <b>NS</b> , D5W <sup>128</sup>  25-50 mL <b>NS</b> , D5W <sup>129</sup>	24 h F, RT <sup>128</sup>	- auxiliary info: <b>WARNING: FOR            INTRAVENOUS            USE ONLY –            FATAL IF GIVEN            BY OTHER            ROUTES</b> <sup>123,124</sup> - for ULYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing etoposide, DOXOrubicin, vinCRISTine)
<b>Vinorelbine</b> 10 mg/1 mL 50 mg/5mL (Fresenius Kabi) (F)(PFL) no preservative <sup>130</sup>	N/A	10 mg/mL <sup>130</sup>	discard unused portion <sup>130</sup>	0.5-2.0 mg/mL <sup>130</sup>  <b>NS</b> , D5W, ½NS, D5½NS, Ringer's, Ringer's Lactate <sup>130</sup>	24 h F, RT <sup>130</sup>	- auxiliary info: <b>WARNING: FOR            INTRAVENOUS            USE ONLY –            FATAL IF GIVEN            BY OTHER            ROUTES</b> <sup>123,131</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Vinorelbine</b> 10 mg/1 mL 50 mg/5 mL (GMP) (F)(PFL) no preservative <sup>132</sup>	N/A	10 mg/mL <sup>132</sup>	discard unused portion <sup>2</sup>	0.5-2.0 mg/mL <sup>132</sup>  50 mL* <b>NS</b> , D5W, ½NS, D5½NS, Ringer's, Ringer's Lactate <sup>132</sup>	24 h <b>F</b> , RT <sup>132</sup>	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES <sup>123,131</sup>
<b>Vinorelbine</b> 10 mg/1 mL 50 mg/5 mL (Pfizer/Hospira) (F)(PFL) no preservative <sup>133</sup>	N/A	10 mg/mL <sup>133</sup>	discard unused portion <sup>133</sup>	0.5–2.0 mg/mL <sup>133</sup>  50 mL* <b>NS</b> , D5W, ½NS, D5½NS, Ringer's, Ringer's Lactate <sup>133</sup>	24 h <b>F</b> , RT <sup>133</sup>	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES <sup>123,131</sup>
<b>Vinorelbine</b> 10 mg/1 mL 50 mg/5 mL (Teva) (F)(PFL) no preservative <sup>134</sup>	N/A	10 mg/mL <sup>134</sup>	discard unused portion <sup>134</sup>	0.5–2.0 mg/mL <sup>134</sup>  50 mL* <b>NS</b> , D5W, ½NS, D5½NS, Ringer's, Ringer's Lactate <sup>134</sup>	24 h <b>F</b> , RT <sup>134</sup>	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES <sup>123,131</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Zoledronic acid</b> 4 mg/5 mL (Dr Reddy's) (RT) no preservative <sup>135</sup>	N/A	0.8 mg/mL <sup>135</sup>	discard unused portion <sup>135</sup>	100 mL <b>NS</b> , D5W <sup>135</sup>	complete infusion within 24 h of preparation <sup>135</sup>  <b>Refrigerate</b> diluted product if not used immediately after preparation; bring to RT prior to administration <sup>135</sup>	- do <b>NOT</b> mix with calcium containing solutions <sup>135</sup>
<b>Zoledronic acid</b> 4 mg/5 mL (Marcan) (RT) no preservative <sup>136</sup>	N/A	0.8 mg/mL <sup>136</sup>	discard unused portion <sup>136</sup>	100 mL <b>NS</b> , D5W <sup>136</sup>	complete infusion within 24 h of preparation <sup>136</sup>  <b>Refrigerate</b> diluted product if not used immediately after preparation; bring to RT prior to administration <sup>136</sup>	- do <b>NOT</b> mix with calcium containing solutions (e.g., Lactated Ringer's) <sup>136</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Zoledronic acid</b> 4 mg/5 mL (MDA) (RT) no preservative <sup>137</sup>	N/A	0.8 mg/mL <sup>137</sup>	discard unused portion <sup>137</sup>	100 mL <b>NS</b> , D5W <sup>137</sup>	complete infusion within 24 h of preparation <sup>137</sup>  <b>Refrigerate</b> diluted product if not used immediately after preparation; bring to RT prior to administration <sup>137</sup>	- do <b>NOT</b> mix with calcium containing solutions <sup>137</sup>
<b>Zoledronic acid</b> (ZOMETA) 4 mg/ 5 mL (Novartis) (RT) no preservative <sup>138</sup>	N/A	0.8 mg/mL <sup>138</sup>	discard unused portion <sup>39</sup>	100 mL <b>NS</b> , D5W <sup>138</sup>	complete infusion within 24 h of preparation <sup>138</sup>  <b>Refrigerate</b> diluted product if not used immediately after preparation; bring to RT prior to administration <sup>138</sup>	- do NOT mix with calcium containing solutions <sup>138</sup>

**BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART**

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product	Product Stability	Special Precautions/Notes
<b>Zoledronic acid</b> 4 mg/5 mL (Sandoz) (RT) no preservative <sup>139</sup>	N/A	0.8 mg/mL <sup>139</sup>	discard unused portion <sup>139</sup>	100 ml <b>NS</b> , D5W <sup>139</sup>	complete infusion within 24 h of preparation <sup>139</sup>  Refrigerate diluted product if not used immediately after preparation; bring to RT prior to administration <sup>139</sup>	- do NOT mix with calcium- or other divalent cation- containing infusion solutions (e.g., Lactated Ringer's) <sup>139</sup>

\* Suggested volume based on usual dose range and any concentration range of stability data

\*\* Protect from light means minimizing exposure to direct sunlight over a *storage* period. More specific information on protection from light (eg, protecting container and tubing during *administration*) will be indicated in the Under the Special Precautions/Notes column.

\*\*\* Contains DMA (N,N dimethylacetamide). Product may be incompatible with closed system transfer devices such as ChemoLock.

**Centres are not to change the content locally but should forward suggestions to the Cancer Drug Manual staff.**

**Explanatory Notes:**

- *Stability data* assumes products prepared using standard aseptic technique in biological safety cabinet at low risk for contamination according to the classification outlined in USP 797.<sup>140,141</sup>
- *Vial stability*: Stability of solution after first puncture or reconstituted solution.
- *Storage temperature*: If information states same stability with refrigerator and room temperature storage, then fridge stability is bolded as preferred (ie, to minimize growth of micro-organisms).
- *Discard unused portion*: Unused portion from single use vials should be discarded at the end of the day.
- “*overflow known*” is stated if the manufacturer states overflow that is present is within acceptable limits.
- “*Complete administration within \_\_\_*” is stated if the manufacturer specifies that the infusion must be completed in a specific time frame following preparation, usually including entire time required for preparation (from first puncture), storage, and administration of infusion.

## Abbreviations:

BWI = bacteriostatic water for injection  
CIVI: ambulatory pump = Continuous Intravenous Infusion (e.g., elastomeric infusor)  
D5W = dextrose 5% in water  
DMA = N,N dimethylacetamide  
F = refrigerate  
Non-DEHP = not containing Di(2-ethylhexyl) phthalate (DEHP)  
NS = normal saline  
PFL = protect from light  
RT = room temperature  
SWI = sterile water for injection

## References:

1. Generic Medical Partners Inc. Leucovorin calcium injection product monograph. Toronto, Ontario; 13 August 2018.
2. BC Cancer. Provincial Pharmacy Directive Number II-20: Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer; 5 December 2018.
3. Pfizer Canada Inc. Leucovorin calcium injection product monograph. Kirkland, Quebec; 21 June 2018.
4. Teva Canada Limited. Leucovorin calcium injection® product monograph. Toronto, Ontario; 5 May 2014.
5. Hospira Healthcare Corporation. LEUCOVORIN CALCIUM INJECTION® product monograph. Saint-Laurent, Quebec; 7 June 2007.
6. Novopharm Limited (Teva). LEUCOVORIN CALCIUM® Injection product information package. Toronto, Ontario; undated.
7. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 6 January 2006.
8. Jenny Yeung. Personal communication. Medical Information Specialist, Teva Canada; 12 April 2017.
9. GlaxoSmithKline Inc. Alkeran Package Insert. Mississauga, Ontario; Montreal, Quebec; 2004.
10. Trissel LA. Handbook on Injectable Drugs. 13th ed. Bethesda, MD: American Society of Health-System Pharmacists, Inc.; 2005.
11. Baxter Corporation. UROMITEXAN® product monograph. Mississauga, Ontario; 6 August 2013.
12. Mona Ghobros BPharm MSc. Personal communication. Medical Information, Baxter Corporation; 29 November 2018.
13. Trissel's® 2 Clinical Pharmaceutics Database (database on the Internet). Mesna. Lexi-Comp Inc.; created by Lawrence A. Trissel, Available at: <http://online.lexi.com>. Accessed 29 November 2018.
14. Fresenius Kabi Canada Ltd. Mesna for injection product monograph. Richmond Hill, Ontario; 21 December 2017.
15. BC Cancer. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer; 19 September 2007.
16. Fresenius Kabi Canada Ltd. Mesna for injection product monograph. Richmond Hill, Ontario; 30 March 2015.
17. Accord Healthcare Inc. Methotrexate injection product monograph. Kirkland, Quebec; 6 April 2018.
18. BC Cancer Agency Miscellaneous Origins Tumour Group. (MOHDMTX) BCCA Protocol Summary for Treatment of Meningeal Disease (Miscellaneous Tumour Origins) using High Dose Methotrexate with Leucovorin Rescue. Vancouver, British Columbia: BC Cancer Agency; 1 Jan 2013.
19. BC Cancer Agency Sarcoma Tumour Group. (SAHDMTX) BCCA Protocol Summary for Treatment of Osteosarcoma Using High Dose Methotrexate with Leucovorin Rescue. Vancouver, British Columbia: BC Cancer Agency; 1 Nov 2012.
20. BC Cancer Agency Lymphoma Tumour Group. (LYHDMRP) BCCA Protocol Summary for Treatment of Primary Intracerebral Lymphoma with High Dose Methotrexate and ritUXimab. Vancouver, British Columbia: BC Cancer Agency; 1 Jun 2014.
21. BC Cancer Agency Lymphoma Tumour Group. (LYHDMTXP) BCCA Protocol Summary for Treatment of Primary Intracerebral Lymphoma with High Dose Methotrexate. Vancouver, British Columbia: BC Cancer Agency; 1 Jun 2014.

22. BC Cancer Agency Lymphoma Tumour Group. (LYHDMTXR) BCCA Protocol Summary for Treatment of Leptomeningeal Lymphoma or Recurrent Intracerebral Lymphoma with High Dose Methotrexate. Vancouver, British Columbia: BC Cancer Agency; 1 Jun 2014.
23. Mayne Pharma Canada. Methotrexate Product Monograph. Montreal, Quebec; December 2003.
24. BC Cancer Lymphoma Tumour Group. (LYIT) BC Cancer Protocol Summary for Treatment of Lymphoma using Intrathecal Methotrexate and Cytarabine. Vancouver, British Columbia: BC Cancer; 1 June 2014.
25. BC Cancer Miscellaneous Origin Tumour Group. (MOIT) BC Cancer Protocol Summary for Solid Tumours using Intrathecal Methotrexate and/or Thiotepa and/or Cytarabine. Vancouver, British Columbia: BC Cancer; 1 October 2018.
26. BC Cancer. Systemic Therapy Policy and Procedure III-50: Administration of High Alert Medications by the Intrathecal Route via Lumbar Puncture or Ommaya Reservoir. Vancouver, British Columbia; 1 May 2019.
27. Pfizer Canada Inc. Methotrexate injection product monograph. Kirkland, Quebec; 13 October 2017.
28. Accord Healthcare Inc. Mitomycin product monograph. Kirkland, Quebec; 7 June 2017.
29. Au JLS, Badalament RA, Wientjes MG, et al. Methods to improve efficacy of intravesical mitomycin C: results of a randomized phase III trial. J Natl Cancer Inst April 18, 2001;93(8):597-604.
30. Jessie LS Au PharmD PhD. Personal communication. Distinguished University Professor, The Ohio State University; 14 May 2007.
31. Myers AL, Zhang Y, Kawedia JD, et al. Solubilization and stability of mitomycin C solutions prepared for intravesical administration. Drugs R D 2017;17:297-304.
32. Teva Canada Limited. Mitomycin for injection® product monograph. Toronto, Ontario; 30 June 2017.
33. Fresenius Kabi Canada Ltd. Mitoxantrone injection® product monograph. Richmond Hill, Ontario; 28 September 2016.
34. Hospira Healthcare Corporation. Mitoxantrone Injection, USP Product Monograph. Saint-Laurent, Quebec; 13 June 2007.
35. Novopharm Limited. Mitoxantrone Injection Product Monograph. Toronto, Ontario; 19 May 2005.
36. Angie Chan. Personal communication. Drug Information Pharmacist, Novopharm; 7 Mar 2008.
37. Bristol-Myers Squibb Canada. OPDIVO® product monograph. Montreal, Quebec; 16 July 2018.
38. Hoffmann-La Roche Ltd. GAZYVA® product monograph. Mississauga, Ontario; 21 December 2015.
39. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 19 September 2007.
40. Anna Sivojelezova MSc. Personal communication. Drug Information Associate; Hoffmann-La Roche Ltd Drug Information; 24 April 2015.
41. Omega Laboratories Ltd. Octreotide Acetate Injection product monograph. Montreal, Quebec; 23 July 2010.
42. Novopharm Limited. Octreotide Injection Product Monograph. Scarborough, Ontario; 15 March 2007.
43. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 19 September 2007.
44. Novartis Pharmaceuticals Canada Inc. SANDOSTATIN® Product Monograph. Dorval, Quebec; 9 January 2001.
45. Repchinsky C, editor. Sandostatin LAR monograph, Compendium of Pharmaceuticals and Specialties. Ottawa, Ontario: Canadian Pharmacists Association; 2005. p. 1912-1916.
46. Repchinsky C editor. Compendium of Pharmaceuticals and Specialties. 12th ed. Ottawa, Ontario: Canadian Pharmacists Association; 2004.
47. Vancouver Hospital and Health Sciences Centre Pharmacy Department. Octreotide. Parenteral drug therapy manual. Vancouver, BC; February 2002.
48. Eli Lilly Canada Inc. LARTRUVO® product monograph. Toronto, Ontario; 23 November 2017.
49. Pfizer Canada Inc. Oxaliplatin injection product monograph. Kirkland, Quebec; 31 May 2017.
50. Sandoz Canada Inc. Oxaliplatin injection product monograph. Boucherville, Quebec; 12 August 2015.
51. Medical Information. Personal communication. Pfizer Canada Inc.; 6 June 2017.
52. Katryn Vosburg. Personal communication. Drug Information & Pharmacovigilance Specialist, Sandoz Canada Inc.; 26 February 2016.
53. Teva Canada Limited. Teva-Oxaliplatin injection® product monograph. Toronto, Ontario; 11 September 2015.
54. Accord Healthcare Inc. Paclitaxel injection product monograph. Markham, Ontario; 13 August 2012.
55. Claude Mercure. Personal communication. Production Manager Biolyse Pharma Corporation; 21 December 2009.
56. Biolyse. PACLITAXEL FOR INJECTION® product monograph. St. Catharines, Ontario; 2 December 2005.
57. Claude Mercure. Personal communication. Manager, Biolyse Pharma Corporation; 24 June 2014.
58. Zeng Z, Lazakovitch E. Study IR 120: Physical and Chemical Stability Study of Paclitaxel for Injection in 0.9 % Sodium Chloride in concentration range 0.012-0.12 mg/mL. Biolyse Pharma March 2010.
59. Mercure C. Stability of 0.1 mg/mL of paclitaxel for injection in sodium chloride (0.9%) solution. St Catharines, Ontario: Biolyse Pharma; 2 February 2007.

60. Xu Q, Trissel LA, Martinez JF. Stability of paclitaxel in 5% dextrose injection or 0.9% sodium chloride injection at 4, 22, or 32 degrees C. *Am J Hosp Pharm* 1994;51(24):3058-60.
61. Lisa Tavano. Personal communication. Biolyse Pharma Corporation; 14 May 2012.
62. Robyn MacKenzie. Personal communication. Area Manager, Hospira Healthcare Corporation; 4 April 2012.
63. Hospira Healthcare Corporation. PACLITAXEL FOR INJECTION® product monograph. Saint-Laurent, Quebec; 1 September 2009.
64. Rose Toussaint. Personal communication. Hospira Canada Healthcare Corporation; 4 April 2012.
65. Celgene Inc. ABRAXANE® product monograph. Mississauga, Ontario; 24 July 2014.
66. Aisling Cahill. Drug Safety and Medical Information Specialist. Personal communication. Celgene Inc.; 23 April 2015.
67. Celgene Europe Limited. ABRAXANE® product monograph. Uxbridge, UK; 11 January 2013.
68. Celgene Inc. ABRAXANE® product monograph. Mississauga, Ontario; 18 January 2016.
69. Pharmaceutical Partners of Canada. Pamidronate Disodium For Injection product monograph. Richmond Hill, Ontario; 18 January 2010.
70. Mayne Pharma (Canada) Inc. Pamidronate Package Insert. Montreal, Quebec; 2002.
71. Omega Laboratories Ltd. Pamidronate Disodium product monograph. Montreal, Quebec; 06 June 2005.
72. Pfizer Canada ULC. Pamidronate disodium for injection product monograph. Kirkland, Quebec; 11 December 2018.
73. Sandoz Canada Inc. Pamidronate injection product monograph. Boucherville, Quebec; 28 February 2006.
74. Amgen Canada. VECTIBIX® product monograph. Mississauga, Ontario; 5 March 2009.
75. Diane Lord. Personal communication. Amgen Canada Inc Medical Information Department; 19 June 2009.
76. Shire Pharma Canada ULC. ONCASPAR® product monograph. Toronto, Ontario; 20 April 2018.
77. Merck Canada Inc. KEYTRUDA® product monograph. Kirkland, Quebec; 21 February 2018.
78. Merck Canada Inc. KEYTRUDA® product monograph. Kirkland, Quebec; 5 February 2016.
79. Accord Healthcare Inc. Pemetrexed disodium for injection product monograph. Kirkland, Quebec; 12 March 2015.
80. Eli Lilly Canada. Pemetrexed product information. Toronto, Ontario; 2008.
81. Eli Lilly Canada Inc. ALIMTA® Product Monograph. Toronto, Ontario; 21 May 2004.
82. Hoffmann-La Roche Limited. PERJETA® product monograph. Mississauga, Ontario; 12 April 2013.
83. sanofi-aventis Canada Inc. MOZOBIL® product monograph. Laval, Quebec; 8 October 2014.
84. Maureen Coughlin BSc Pharm. Personal communication. Solutions in Health Inc. acting as an authorized agent of sanofi-aventis; 24 May 2017.
85. Axcan Pharma Inc. Photofrin Package Insert. Mont-Saint-Hilaire, Quebec; 2004.
86. Marie-Helene Doyon B.Pharm. M.Sc. Personal communication. Medical Information Specialist, Porfimer, Axcan Pharma. 3 March 2005.
87. Pfizer Canada Inc. TOMUDEX® product monograph. Kirkland, Quebec; 8 August 2017.
88. Eli Lilly Canada Inc. CYRAMZA® product monograph. Toronto, Ontario; 16 July 2015.
89. Marilyn Bain BScN. Personal communication. Senior Medical Information Associate, Eli Lilly Canada, Inc.; 16 January 2017.
90. Hoffmann-La Roche Ltd. RITUXAN® product monograph. Mississauga, Ontario; 29 March 2012.
91. Hoffmann-La Roche Ltd. RITUXAN® product monograph. Mississauga, Ontario; 29 May 2014.
92. Diana Fung pharmacist. Personal communication. Hoffmann-La Roche Medical Information; 16 July 2014.
93. Hoffmann-La Roche Ltd. RITUXAN® SC product monograph. Mississauga, Ontario; 21 March 2018.
94. Celgene Inc. ISTODAX® product monograph. Mississauga, Ontario; 13 December 2016.
95. Celgene Inc. INFO Rx ISTODAX® (romidepsin) for Injection. Mississauga, Ontario; 10 July 2017.
96. Aisling Cahill, Drug Safety and Medical Information Specialist. Personal communication. Celgene Inc.; 17 July 2015.
97. Janssen Inc. SYLVANT® product monograph. Toronto, Ontario; 6 January 2016.
98. Pharmacia Canada Inc. Zanosar Package Insert. Mississauga, Ontario; March 2003.
99. Wyeth Canada. TORISEL® product monograph. Montreal, Canada; 16 October 2008.
100. McEvoy GK, editor. AHFS 2008 Drug Information. Bethesda, Maryland: American Society of Health-System Pharmacists, Inc. p. 1226-1228.
101. Anna Sivojelezova M.Sc. Personal communication. Medical Information Associate, Wyeth. 6 January 2010.
102. Bristol-Myers Squibb Canada. VUMON® product monograph. St. Laurent, Quebec; 26 October 2004.
103. Trissel's®2 IV Compatibility (database on the Internet). Teniposide. Thomson Reuters MICROMEDEX® 2.0, updated periodically. Available at: <http://www.micromedex.com>. Accessed 27 April 2011.
104. Adienne SA. TEPADINA® product monograph. Lugano, Switzerland; 28 March 2017.



105. AHFS Drug Information® (database on the Internet). Thiotepa. Lexi-Comp Inc., 27 February 2018. Available at: <http://online.lexi.com>. Accessed 21 August 2018.
106. Hematology/Oncology Pharmacy Association. HOPA News Clinical Pearls: Intrathecal Chemotherapy: Focus on Drugs, Dosing, and Preparation. 13(4) ed. Chicago, Illinois, USA: Hematology/Oncology Pharmacy Association; 2016.
107. Genzyme Canada. Thyrogen Product Monograph. Mississauga, Ontario; 2004.
108. Hoffmann-La Roche Limited. ACTEMRA® product monograph. Mississauga, Ontario; 27 October 2017.
109. Accord Healthcare Inc. Topotecan hydrochloride for injection product monograph. Kirkland, Quebec; 9 May 2019.
110. Actavis Pharma Company. Topotecan for injection product monograph. Mississauga, Ontario; 3 January 2018.
111. Pfizer Canada Inc. Topotecan hydrochloride for injection product monograph. Kirkland, Quebec; 5 January 2018.
112. Sandoz Canada Inc. Topotecan injection product monograph. Boucherville, Quebec; 5 September 2014.
113. Hoffman-La Roche Limited. HERCEPTIN® product monograph. Mississauga, Ontario; 16 November 2012.
114. Hoffmann-La Roche Limited. KADCYLA® product monograph. Mississauga, Ontario; 11 September 2013.
115. Tracon Pharmaceuticals Inc. TRC105 (carotuximab) investigational brochure. San Diego, California; 14 February 2017 (version 10.0).
116. Tracon Pharmaceuticals Inc. Clincial Protocol: A Phase 2A Study of TRC105 (with Option to Add Bevacizumab) in Patients with Refractory Gestastional Trophoblastic Neoplasia (GTN). San Diego, California; 20 September 2016 Amendment #3.
117. medac UK. TREOSULFAN injection® product monograph. Hamburg, Germany; 24 June 2008.
118. Henrik Fenger. Personal communication. Management Associate, International Division medac; 03 March 2010.
119. Mayne Pharma (Canada) Inc. Vinblastine product monograph. Kirkland, Quebec; 10 August 2003.
120. Lexi-Drugs® (database on the Internet). VinBLASTine. Lexi-Comp Inc., 3 November 2014. Available at: <http://online.lexi.com>. Accessed 20 November 2014.
121. Jan Barrow. Personal communication. Supervisor, Hospira Canada Clinical Support. Saint-Laurent, Quebec; 03 December 2007.
122. Tanya Leduc. Personal communication. Acting editor, BC Cancer Agency Cancer Drug Manual; 18 Dec 2007.
123. World Health Organization. Information Exchange System: Alert No. 115 (QSM/MC/IEA.115). Geneva, Switzerland: World Health Organization; 18 July 2007.
124. BCCA Provincial Systemic Therapy Program. Labeling of vinca alkaloid syringes. Policy # V-40. Vancouver, British Columbia: BC Cancer Agency; 27 May 1999.
125. Teva Canada Limited. Vinblastine sulfate injection® product monograph. Toronto, Ontario; 1 February 2013.
126. Trissel's® 2 IV Compatibility (database on the Internet). Vinblastine sulfate. Thomson Reuters MICROMEDEX® 2.0, updated periodically. Available at: <http://www.micromedex.com>. Accessed 8 June 2015.
127. Mayne Pharma (Canada) Inc. Vincristine Package Insert. Montreal, QC; Undated.
128. Teva Canada Limited. Vincristine sulfate injection® product monograph. Scarborough, Ontario; 27 March 2014.
129. Lexi-Drugs® (database on the Internet). VinCRISTine. Lexi-Comp Inc., 3 June 2015. Available at: <http://online.lexi.com>. Accessed 9 June 2015.
130. Pharmaceutical Partners of Canada. Vinorelbine Injection product monograph. Richmond Hill, Ontario; 15 January 2008.
131. BC Cancer Provincial Systemic Therapy Program. Policy V-40: Dispensing and Labelling of Vinca Alkaloid Preparations. Vancouver, British Columbia: BC Cancer; 1 April 2015.
132. Generic Medical Partners Inc. Vinorelbine Injection product monograph. Toronto, Ontario; 3 September 2014.
133. Pfizer Canada Inc. Vinorelbine Tartrate for Injection product monograph. Kirkland, Quebec; 21 July 2017.
134. Teva Canada Limited. Vinorelbine tartrate for Injection product monograph. Toronto, Ontario; 20 March 2014.
135. Innomar Strategies Inc. (for Dr. Reddy's Laboratories Limited). Zoledronic acid for injection concentrate® product monograph. Oakville, Ontario; 11 March 2015.
136. Marcan Pharmaceuticals Inc. Zoledronic acid concentrate for injection product monograph. Ottawa, Ontario; 5 February 2018.
137. MDA Inc. Zoledronic acid for injection product monograph. Mississauga, Ontario; 11 August 2015.
138. Novartis Pharmaceuticals Canada Inc. ZOMETA® product monograph. Dorval, Quebec; 26 July 2013.
139. Sandoz Canada Inc. Zoledronic Acid - Z® product monograph. Boucherville, Quebec; 02 December 2016.
140. The United States Pharmacopeia (USP). General Chapter 797: Pharmaceutical compounding - sterile preparations. USP 27-NF 22. Rockville, Maryland: The United States Pharmacopeial Convention, Inc.; 2004.
141. Kastango ES. The ASHP discussion guide for compounding sterile preparations. Bethesda (MD): American Society of Health-System Pharmacists, Inc.; 2004. p. 5.