

DRUG NAME: Brentuximab vedotin

SYNONYM(S): SGN-35¹

COMMON TRADE NAME(S): ADCETRIS®

CLASSIFICATION: miscellaneous

Special pediatric considerations are noted when applicable, otherwise adult provisions apply.

MECHANISM OF ACTION:

Brentuximab vedotin is an antibody-drug conjugate (ADC) composed of a chimeric monoclonal antibody (cAC10) linked to an anti-tubulin agent, monomethylauristatin E (MMAE). The monoclonal antibody targets CD-30 expressing cells and MMAE is released intracellularly to bind to tubulin. The binding of MMAE to tubulin disrupts the microtubule network, causing cell cycle arrest and apoptosis.² Brentuximab vedotin is cell cycle phase-specific (G2/M phase).³

PHARMACOKINETICS:

Absorption	monomethylauristatin E (MMAE): time to peak 1-3 days	
Distribution	antibody-drug conjugate (ADC) is primarily limited to vascular space; organ distribution of MMAE in humans is unknown	
	cross blood brain barrier?	no information found
	volume of distribution	6-10 L (ADC); 44 L (MMAE)
	plasma protein binding	68-82% (MMAE)
Metabolism	MMAE: minimal; primarily via oxidation by CYP 3A4/5	
	active metabolite(s)	yes; at concentrations less than 10% of MMAE
	inactive metabolite(s)	no information found
Excretion	primarily as unchanged MMAE; limited by the rate of release from ADC	
	urine	MMAE: 72%
	feces	MMAE: 28%
	terminal half life	4-6 days
	clearance	no information found
Sex	ADC volume of distribution is 14% lower in females	

Adapted from standard reference^{3,4} unless specified otherwise.

USES:

Primary uses:

- *Lymphoma, Hodgkin's
- *Lymphoma, non-Hodgkin's
- *Lymphoma, cutaneous T-cell

*Health Canada approved indication

Other uses:

SPECIAL PRECAUTIONS:

Contraindications:

- history of hypersensitivity reaction to brentuximab vedotin or Chinese hamster ovary cell proteins³
- concurrent therapy with bleomycin; due to increased risk of pulmonary toxicity, bleomycin must be discontinued prior to starting brentuximab vedotin treatment²
- patients with a diagnosis of progressive multifocal leukoencephalopathy (PML) or a history of PML³

Caution:

- patients with rapidly proliferating tumour and high tumour burden are at risk of **tumour lysis syndrome**²
- concurrent medications should be carefully reviewed for potential **drug interactions**, particularly in regard to CYP 3A4 and CYP 3A5²

Carcinogenicity: no information found

Mutagenicity: Not mutagenic in Ames test or mammalian *in vitro* mutation test. Brentuximab vedotin is clastogenic in mammalian *in vivo* chromosome tests.³

Fertility: In rats, brentuximab vedotin causes partially reversible, dose-dependent testicular toxicity (e.g., seminiferous tubule degeneration, Sertoli cell vacuolation, reduced spermatogenesis, and aspermia).³

Pregnancy: FDA Pregnancy Category D.² There is positive evidence of human fetal risk, but the benefits from use in pregnant women may be acceptable despite the risk (e.g., if the drug is needed in a life-threatening situation or for a serious disease for which safer drugs cannot be used or are ineffective).

In rats, brentuximab vedotin has been shown to cross the placenta, causing embryo-fetal toxicities, including early resorption, post-implantation loss, decreased numbers of live fetuses, and external malformations (i.e., umbilical hernias and malrotated hindlimbs). For women of childbearing potential, two contraceptive methods are recommended during treatment.³

Breastfeeding is not recommended due to the potential secretion into breast milk.

SIDE EFFECTS:

The table includes adverse events that presented during drug treatment but may not necessarily have a causal relationship with the drug. Because clinical trials are conducted under very specific conditions, the adverse event rates observed may not reflect the rates observed in clinical practice. Adverse events are generally included if they were reported in more than 1% of patients in the product monograph or pivotal trials.⁵ When placebo-controlled trials are available, adverse events will generally be included if the incidence is $\geq 5\%$ higher in the treatment group.

ORGAN SITE	SIDE EFFECT
Clinically important side effects are in bold, italics	
allergy/immunology	infusion-related reactions (12%); see paragraph following Side Effects table
blood and lymphatic system/ febrile neutropenia	anemia (33-52%, severe 2-10%)
	lymphadenopathy (10-11%)
	neutropenia (54-55%, severe 21%); can be prolonged (≥ 1 week), may require dose modification, delays, or growth factor support ³
cardiac	thrombocytopenia (16-28%, severe 10%); can be prolonged (≥ 1 week), may require dose delays or platelet transfusions ³
	myocardial infarction (<1%) ³
	QT shortening ³ ; clinical significance is unknown

ORGAN SITE	SIDE EFFECT
Clinically important side effects are in <i>bold, italics</i>	
	supraventricular arrhythmia (3%)
gastrointestinal	<i>emetogenic potential: low</i> ⁶
	abdominal pain (9-25%, severe 1-2%)
	constipation (16-19%, severe 2%)
	diarrhea (29-36%, severe 1-3%)
	nausea (38-42%, severe 2%)
	pancreatitis ⁷ ; sometimes fatal
	vomiting (17-22%, severe 3%)
general disorders and administration site conditions	<i>extravasation hazard: none</i> ⁸
	chills (12-13%)
	fatigue (41-49%, severe 2-3%)
	fever (29-38%, severe 2%)
	night sweats (9-12%)
	peripheral edema (4-16%)
infections and infestations	septic shock (3%)
	upper respiratory tract infection (12-47%)
	urinary tract infection (3%)
metabolism and nutrition	anorexia (11-16%, severe 2%)
	tumour lysis syndrome ³ (<1%); see paragraph following Side Effects table
	weight loss (1-12%, severe 3%)
musculoskeletal and connective tissue	arthralgia (9-19%)
	back pain (10-14%, severe 2%)
	muscle spasms (9-10%, severe 2%)
	myalgia (16-17%, severe 2%)
	pain (7-28%, severe 5%)
	pain in extremity (10%, severe 4%)
nervous system	anxiety (7-11%, severe 2%)
	dizziness (11-16%)
	headache (16-19%, severe 2%)
	insomnia (14-16%)
	peripheral motor neuropathy (7-16%, severe 3-4%); see paragraph following Side Effects table
	<i>peripheral sensory neuropathy</i> (52-53%, severe 8-10%); see paragraph following Side Effects table
	<i>progressive multifocal leukoencephalopathy</i> (< 1%); see paragraph following Side Effects table
renal and urinary	pyelonephritis (2%)
respiratory, thoracic and mediastinal	cough (17-25%)
	dyspnea (13-19%, severe 1-2%)
	oropharyngeal pain (9-11%)

ORGAN SITE	SIDE EFFECT
Clinically important side effects are in <i>bold, italics</i>	
	pneumonitis (2%)
	pneumothorax (2%)
	pulmonary embolism (2%)
skin and subcutaneous tissue	alopecia (13-14%)
	dry skin (4-10%)
	pruritus (17-19%)
	rash (27-31%)
	Stevens-Johnson syndrome

Adapted from standard reference² unless specified otherwise.

Peripheral neuropathy is usually sensory in nature, but motor neuropathy has also been reported. Peripheral neuropathy is dose-cumulative² and usually occurs several months into therapy.⁹ Improvement or resolution of symptoms usually takes an average of seven weeks.⁹ About half of patients who experience neuropathy may have residual symptoms of neuropathy. Symptoms include hypoesthesia, hyperesthesia, paresthesia, discomfort, a burning sensation, neuropathic pain or weakness. Dose modifications or discontinuation may be needed.²

Infusion-related reactions, including anaphylaxis, have been reported. Reactions may be immediate or delayed, with symptoms occurring within 2 days after administration.³ Symptoms include chills, nausea, dyspnea, pruritus, pyrexia and cough.² Severe reactions present as wheezing, difficulty breathing, hives, itching, and swelling.³ If a reaction occurs, interrupt the infusion and institute appropriate medical management. Patients with a prior reaction should be premedicated with acetaminophen, antihistamine, and corticosteroid for subsequent infusions. If anaphylaxis occurs, the infusion should be immediately and permanently discontinued, and appropriate medical management instituted.^{2,3} For management of infusion-related reactions, see BC Cancer Protocol SCDRUGRX [Management of Infusion-Related Reactions to Systemic Therapy Agents](#).

Progressive multifocal leukoencephalopathy (PML) and death have been reported following active John Cunningham (JC) virus infection in patients treated with brentuximab vedotin. Other risk factors include prior immunosuppressive therapies or underlying immunosuppressive disease. PML should be considered in any patient treated with brentuximab vedotin who presents with new onset of signs and symptoms of central nervous system abnormalities. Symptoms include changes in mood or usual behaviour, confusion, thinking problems, loss of memory, changes in vision, speech or walking, and decreased strength or weakness on one side of the body. Brentuximab vedotin should be held if PML is suspected, and discontinued permanently if diagnosis is confirmed.²

Tumour lysis syndrome has been reported; patients with rapidly proliferating tumour or high tumour burden may be at increased risk. Monitor patient closely; symptoms include nausea, vomiting, edema, shortness of breath, heart rhythm disturbances, and acute renal failure.³

INTERACTIONS:

AGENT	EFFECT	MECHANISM	MANAGEMENT
grapefruit juice ³	may increase plasma level of monomethylauristatin E (MMAE)	may inhibit CYP 3A4 metabolism of MMAE in the intestinal wall	monitor for brentuximab vedotin toxicity

AGENT	EFFECT	MECHANISM	MANAGEMENT
ketoconazole ³	increased AUC of MMAE by 34%	potent inhibition of CYP 3A4 and P-gp by ketoconazole	monitor for brentuximab vedotin toxicity
rifampin ³	decreased AUC of MMAE by 46%	potent induction of CYP 3A4 by rifampin	avoid concurrent therapy if possible

Monomethylauristatin E (MMAE) is a substrate of CYP 3A4/5. Co-administration with potent inhibitors or inducers of CYP enzymes may affect plasma levels of MMAE. Monitor for brentuximab vedotin toxicity in patients receiving strong CYP 3A4/5 inhibitors.³

MMAE inhibited CYP 3A4/5 in certain *in vitro* studies, however it is not expected to alter the exposure of other drugs metabolized by CYP 3A4 enzymes.³

Brentuximab vedotin is a substrate of p-glycoprotein. Monitor for brentuximab vedotin toxicity in patients receiving strong inhibitors of p-glycoprotein.³

SUPPLY AND STORAGE:

Injection: GMD Distribution Inc. (for Seattle Genetics Inc.) supplies brentuximab vedotin as a lyophilized cake or powder in 50 mg single-use (preservative-free) vials. Refrigerate. Do NOT freeze. Protect from light.²

For basic information on the current brand used at the BC Cancer Agency, see [Chemotherapy Preparation and Stability Chart](#) in Appendix.

SOLUTION PREPARATION AND COMPATIBILITY:

For basic information on the current brand used at the BC Cancer Agency, see [Chemotherapy Preparation and Stability Chart](#) in Appendix.

Additional information:

Compatibility: consult detailed reference

PARENTERAL ADMINISTRATION:

BCCA administration guideline noted in **bold, italics**

Subcutaneous	no information found
Intramuscular	no information found
Direct intravenous	do NOT use ^{2,3}
Intermittent infusion	over 30 minutes^{1-3,10}
Continuous infusion	no information found
Intraperitoneal	no information found
Intrapleural	no information found
Intrathecal	no information found
Intra-arterial	no information found
Intravesical	no information found

DOSAGE GUIDELINES:

Refer to protocol by which patient is being treated. Numerous dosing schedules exist and depend on disease, response and concomitant therapy. Guidelines for dosing also include consideration of absolute neutrophil count (ANC). Dosage may be reduced, delayed or discontinued in patients with bone marrow depression due to cytotoxic/radiation therapy or with other toxicities.

Adults:

BCCA usual dose noted in ***bold, italics***

<i>Intravenous:</i>	Cycle Length:	
	3 weeks^{2,3,9-11:}	<i>1.8 mg/kg IV for one dose on day 1</i> (total dose per cycle 1.8 mg/kg) (for weight > 100 kg: calculate dose based on a weight of 100 kg)
	4 weeks^{12-14:}	<i>1.2 mg/kg IV for one dose on days 1 and 15</i> (total dose per cycle 2.4 mg/kg) (for weight > 100 kg: calculate dose based on a weight of 100 kg)

<i>Dosage in myelosuppression</i>	modify according to protocol by which patient is being treated ¹⁰ ; if no guidelines are available, the following have been suggested ^{2,3} : <ul style="list-style-type: none"> • for grade 3 or 4 neutropenia: hold dose until resolution to baseline or grade 2 or lower; consider filgrastim support for subsequent cycles • for recurrent grade 4 neutropenia despite filgrastim support: discontinue or reduce dose to 1.2 mg/kg • for grade 3 or 4 thrombocytopenia: consider platelet transfusion or dose delay
<i>Dosage in severe peripheral neuropathy^{2,3:}</i>	<ul style="list-style-type: none"> • for new or worsening grade 2 or 3 neuropathy: hold dose until neuropathy improves to grade 1 or baseline; then restart at 1.2 mg/kg • for grade 4 neuropathy: discontinue treatment

Dosage in renal failure: no information found

Dosage in hepatic failure: no information found

Children: no information found

REFERENCES:

1. Seattle Genetics. Pharmacy instructions: Product information and administration - Brentuximab (SGN-35). Bothell, WA; 05 May 2011.
2. Seattle Genetics Inc. ADCETRIS® full prescribing information. Bothell, Washington; January 2012.
3. GMD Distribution Inc. for Seattle Genetics Inc. ADCETRIS® product monograph. Oakville, Ontario; 1 February 2013.
4. Lexi-Drugs® (database on the Internet). Brentuximab vedotin. Lexi-Comp Inc., 10 July 2013. Available at: <http://online.lexi.com>. Accessed 16 July 2013.
5. Laurie Sehn MD. BC Cancer Agency Lymphoma Tumour Group. Personal communication. 30 September 2013.
6. BC Cancer Agency. (SCNAUSEA) Guidelines for Prevention and Treatment of Chemotherapy-induced Nausea and Vomiting in Adults. Vancouver, British Columbia: BC Cancer Agency; 1 Mar 2012.

7. Seattle Genetics Inc. ADCETRIS® full prescribing information. Bothell, Washington; September 2013.
8. BC Cancer Agency Provincial Systemic Therapy Program. Provincial Systemic Therapy Program Policy III-20: Prevention and Management of Extravasation of Chemotherapy. Vancouver, British Columbia: BC Cancer Agency; 1 February 2012.
9. Bradley AM, Devine M, DeRemer D. Brentuximab vedotin: an anti-CD30 antibody-drug conjugate. *Am J Health-Syst Pharm* 2013;70:589-597.
10. BC Cancer Agency Lymphoma Tumour Group. (ULYBRENTUX) BCCA Protocol Summary for Treatment of Hodgkin Lymphoma and Anaplastic Large Cell Lymphoma with Brentuximab. Vancouver, British Columbia: BC Cancer Agency; 1 June 2014.
11. BC Cancer Agency Lymphoma Tumour Group. (LYCTCLBV) BC Cancer Protocol Summary for Treatment of Cutaneous T-Cell Lymphoma (CTCL) with Brentuximab Vedotin. Vancouver, British Columbia: BC Cancer; 1 August 2021.
12. Connors JM, Jurczak W, Straus DJ, et al. Brentuximab vedotin with chemotherapy for stage III or IV Hodgkin's Lymphoma. *N Engl J Med* 2018;378(4):331-344.
13. Straus DJ, Długosz-Danecka M, Alekseev S, et al. Brentuximab vedotin with chemotherapy for stage III/IV classical Hodgkin lymphoma: 3-year update of the ECHELON-1 study. *Blood* 2020;135(10):735-742.
14. BC Cancer Agency Lymphoma Tumour Group. (LYAVDBV) BC Cancer Protocol Summary for Treatment of Previously Untreated, Stage IV Hodgkin Lymphoma with DOXOrubicin, vinBLAStine, Dacarbazine, and Brentuximab Vedotin. Vancouver, British Columbia: BC Cancer; 1 August 2021.