



Provincial Health Services Authority

For the Patient: Daratumumab

Other names: DARZALEX®

- **Daratumumab** (dar" a toom' ue mab) is a drug that is used to treat some types of cancer. It is a monoclonal antibody, a type of protein designed to target and interfere with the growth of cancer cells. It is a clear liquid that is injected into a vein.
- Tell your doctor if you have ever had an unusual or **allergic reaction** to daratumumab before receiving daratumumab.
- A **blood test** may be taken before each treatment. The timing of your treatment may be changed based on the test results and/or other side effects.
- The **drinking of alcohol** (in small amounts) does not appear to affect the safety or usefulness of daratumumab.
- Daratumumab may damage sperm and may harm the baby if used during pregnancy. It is best to use **birth control** while being treated with daratumumab and for at least three months after treatment has ended. Tell your doctor right away if you or your partner becomes pregnant. Do not breastfeed during treatment.
- **Tell** doctors, dentists, and other health professionals that you are being treated with daratumumab before you receive any treatment from them.

Changes in blood counts

Daratumumab may cause temporary changes in your blood counts. Your doctor will be following these changes carefully by performing blood tests. Adjustment of your treatment may be needed in certain circumstances.

BLOOD COUNTS	MANAGEMENT
<p>Normal white blood cells protect your body by fighting bacteria (germs) that cause infection. When they are low, you are at greater risk of having an infection.</p>	<p>To help prevent infection:</p> <ul style="list-style-type: none"> • Wash your hands often and always after using the bathroom. • Avoid crowds and people who are sick. • Call your doctor immediately at the first sign of an infection such as fever (over 100°F or 38°C by an oral thermometer), chills, cough, or burning when you pass urine.

BLOOD COUNTS	MANAGEMENT
<p>Normal platelets help your blood to clot normally after an injury (e.g., cut). When the platelet count is low, you may be more likely to bruise or bleed.</p>	<p>To help prevent bleeding problems:</p> <ul style="list-style-type: none"> • Try not to bruise, cut, or burn yourself. • Clean your nose by blowing gently. Do not pick your nose. • Avoid constipation. • Brush your teeth gently with a soft toothbrush as your gums may bleed more easily. <p>Some medications such as ASA (e.g., ASPIRIN®) or ibuprofen (e.g., ADVIL®) may increase your risk of bleeding.</p> <ul style="list-style-type: none"> • Do not stop taking any medication that has been prescribed by your doctor (e.g., ASA for your heart). • For minor pain, try acetaminophen (e.g., TYLENOL®) first, but occasional use of ibuprofen may be acceptable.

Side effects are listed in the following table in the order in which they may occur. Tips to help manage the side effects are included.

SIDE EFFECTS	MANAGEMENT
<p>Allergic reactions may sometimes occur. Signs of an allergic reaction may include runny or stuffy nose, throat irritation, cough, rash, itching, dizziness, and swelling or breathing problems including wheezing.</p> <p>This can occur immediately or several hours after receiving daratumumab.</p>	<p>Tell your nurse if this happens while you are receiving daratumumab or contact your oncologist immediately if this happens after you leave the clinic.</p>
<p>Pain or tenderness may occur where the needle was placed.</p>	<p>Apply cool compresses or soak in cool water for 15-20 minutes several times a day.</p>
<p>Nausea and vomiting may occur after your treatment. Most people have little or no nausea.</p>	<p>You may be given a prescription for anti-nausea drug(s) to take before your treatment and/or at home. It is easier to prevent nausea than treat it once it has occurred, so follow directions closely.</p> <ul style="list-style-type: none"> • Drink plenty of fluids. • Eat and drink often in small amounts. <p>Try the ideas in <i>Practical Tips to Manage Nausea</i>.*</p>

SIDE EFFECTS	MANAGEMENT
Fever and chills may sometimes occur shortly after treatment with daratumumab. Fever should last no longer than 24 hours.	<ul style="list-style-type: none"> • Take acetaminophen (e.g., TYLENOL®) every 4-6 hours, to a maximum of 4 g (4000 mg) per day. • Fever (over 100°F or 38°C by an oral thermometer) which occurs more than a few days after treatment may be a sign of an infection. Report this to your doctor immediately.
Diarrhea may sometimes occur.	<p>If diarrhea is a problem:</p> <ul style="list-style-type: none"> • Drink plenty of fluids. • Eat and drink often in small amounts. • Avoid high fibre foods as outlined in <i>Food Choices to Help Manage Diarrhea</i>.*
Constipation may sometimes occur.	<ul style="list-style-type: none"> • Exercise if you can. • Drink plenty of fluids. • Try ideas in <i>Food Choices to Manage Constipation</i>.*
Headache may sometimes occur.	Take acetaminophen (e.g., TYLENOL®) every 4-6 hours if needed, to a maximum of 4 g (4000 mg) per day.
Muscle, joint, or back pain may sometimes occur.	You may take acetaminophen (e.g., TYLENOL®) every 4-6 hours to a maximum of 4 g (4000 mg) per day for mild to moderate pain. Tell your doctor if the pain interferes with your activity.
Loss of appetite may sometimes occur.	Try the ideas in <i>Food Ideas to Help with Decreased Appetite</i> .*
Tiredness and lack of energy may sometimes occur.	<ul style="list-style-type: none"> • Do not drive a car or operate machinery if you are feeling tired. • Try the ideas in <i>Fatigue/Tiredness – Patient Handout</i> or <i>Your Bank to Energy Savings: Helping People with Cancer Handle Fatigue</i>.*
Hair loss does not occur with daratumumab.	
Increased calcium in the blood may sometimes occur.	<p>To help prevent high calcium:</p> <ul style="list-style-type: none"> • Keep active. • Drink plenty of fluids.

***Please ask your chemotherapy nurse or pharmacist for a copy.**

