

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Aldesleukin 22 million units (1.3 mg) (SteriMax) (F)(PFL) no preservative ¹	1.2 mL SWI ¹ direct diluent against side of vial during reconstitution ¹ do NOT shake ¹	18 million unit/mL (1.1 mg/mL) ¹	12 h F, RT ^{1,2}	30-70 mcg/mL ¹ 50 mL D5W ¹ <30 mcg/mL: dilute in D5W containing human albumin 0.1% ³	48 h F, RT ¹ bring to RT prior to use ¹	- do NOT use in- line filter ¹ - avoid bacteriostatic water for injection or NS due to increased aggregation ¹
				SC syringe ^{4,5}	10 d F ^{2,5} **(PFL)	
Aldesleukin intralesional 22 million units (1.3 mg) (SteriMax) (F)(PFL) no preservative ¹	1.2 mL SWI ¹ direct diluent against side of vial during reconstitution ¹ do NOT shake ¹	18 million unit/mL (1.1 mg/mL) ¹	12 h F, RT ^{1,2}	add 3.2 mL D5W to reconstituted vial to give 5 million units/mL ^{6,7} withdraw entire contents of vial into syringes for administration ^{6,7}	syringe: 48 h F ⁶ (discard any remaining unused syringes following procedure)	- avoid bacteriostatic water for injection or NS due to increased aggregation ¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Aldesleukin 22 million units (1.3 mg) (Iovance USA) (F)(PFL) no preservative ^{1,8}	1.2 mL SWI ¹ direct diluent against side of vial during reconstitution ¹ do NOT shake ¹	18 million unit/mL (1.1 mg/mL) ¹	12 h F, RT ^{1,2}	30-70 mcg/mL ¹ 50 mL D5W ¹ <30 mcg/mL: dilute in D5W containing human albumin 0.1% ³	48 h F, RT ¹ bring to RT prior to use ¹	- do NOT use in- line filter ¹ - avoid bacteriostatic water for injection or NS due to increased aggregation ¹
				SC syringe ^{4,5}	10 d F ^{2,5} **(PFL)	
Aldesleukin intralesional 22 million units (1.3 mg) (Iovance USA) (F)(PFL) no preservative ^{1,8}	1.2 mL SWI ¹ direct diluent against side of vial during reconstitution ¹ do NOT shake ¹	18 million unit/mL (1.1 mg/mL) ¹	12 h F, RT ^{1,2}	add 3.2 mL D5W to reconstituted vial to give 5 million units/mL ^{6,7} withdraw entire contents of vial into syringes for administration ^{6,7}	syringe: 48 h F ⁶ (discard any remaining unused syringes following procedure)	- avoid bacteriostatic water for injection or NS due to increased aggregation ¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Alemtuzumab 30 mg/mL (Genzyme/Bayer) ⁹ (F)(PFL) do not shake no preservative ¹⁰	N/A	filter NOT required ¹⁰ 30 mg/mL ¹⁰	discard unused portion ¹⁰	SC syringe ¹¹	discard at the end of the day F , RT	- do NOT shake ¹²
				100 mL NS , D5W ¹⁰	8 h F , RT ¹⁰ **(PFL) ¹²	
Amivantamab (JNJ-61186372) ^{13,14} 350 mg (Janssen) (F)(PFL) no preservative ¹⁵ (SAP)	N/A	50 mg/mL	discard unused portion ¹⁵	250 mL NS , D5W ¹⁵ dilute to final volume by withdrawing volume from bag equal to volume of drug to be added ¹⁵ mix by gentle inversion ¹⁵	complete administration within 10 h RT ¹⁵	- do not shake ¹⁵ - discard if discolouration or visible particles are present ¹⁵ - administer with 0.2 micron in-line filter ¹⁵

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Amivantamab 350 mg (Janssen) (F)(PFL) no preservative ¹⁶	N/A	50 mg/mL ¹⁶	discard unused portion ¹⁶	250 mL NS , D5W ¹⁶ dilute to final volume by withdrawing volume from bag equal to volume of drug to be added ¹⁶ mix by gentle inversion; do not shake ¹⁶	complete administration within 10 h RT ¹⁶	- each vial contains 0.5 mL overfill ¹⁶ - discard if discolouration or visible particles are present ¹⁶ - administer with 0.2 micron in-line filter ¹⁶
Amsacrine 75 mg/1.5 mL (Erfa Canada) (RT) no preservative ¹⁷	glass syringes preferred for reconstitution; MAX time in plastic syringe ¹⁷ : 15 min 13.5 mL supplied diluent (L-lactic acid) ¹ to reconstitute: transfer 1.5 mL from ampoule into the diluent vial ¹⁷	5 mg/mL ¹⁷	12 h RT ^{2,17} **(PFL) ¹⁷	500 mL D5W ¹⁷ (plastic or glass container) ¹⁷	7 d F , 4 d RT ^{2,17}	- contains DMA***

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Arsenic trioxide 10 mg/10 mL (Phebra/ICON) (RT) no preservative ¹⁸	N/A	1 mg/mL ¹⁸	discard unused portion ¹⁸	100-250 mL NS , D5W ¹⁸	48 h F, 24 h RT ¹⁸	
Arsenic trioxide 10 mg/10 mL (Sandoz) (RT) no preservative ¹⁹	N/A	1 mg/mL ¹⁹	discard unused portion ¹⁹	100-250 mL NS , D5W ¹⁹	48 h F, 24 h RT ¹⁹	
Arsenic trioxide 10 mg/10 mL (SteriMax) (RT) no preservative ²⁰	N/A	1 mg/mL ²⁰	discard unused portion ²⁰	100-250 mL NS , D5W ²⁰	48 h F, 24 h RT ²⁰	
Asparaginase E. Coli pegylated - see Calaspargase pegol - see Pegaspargase in L-Z chart						
Asparaginase-erwinia (asparaginase <i>Erwinia chrysanthemi</i>) - see Crisantaspase recombinant						

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
PEG-asparaginase (pegylated asparaginase <i>E. coli</i>) - see Calaspargase pegol - see Pegaspargase in L-Z chart						
Atezolizumab 840 mg/14 mL 1200 mg/20 mL (Hoffman-La Roche) (F)(PFL) do not shake no preservative ²¹	N/A	60 mg/mL ²¹	discard unused portion ²¹	250 mL NS ²¹ mix by gentle inversion ²¹	24 h F, 8 h RT ²¹	- do NOT shake ²¹
Atezolizumab subcutaneous (TECENTRIQ® SC) 1875 mg/15 mL Hoffman-La Roche) (F)(PFL) do not shake no preservative ²²	N/A	125 mg/mL ²²	discard unused portion ²	SC syringe ²²	10 d F, 8 h RT ^{2,22}	- contains hyaluronidase ²² - formulations are NOT interchangeable ²²

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Avelumab 200 mg/10 mL (EMD) (F)(PFL) no preservative ²³	N/A	20 mg/mL ²³	discard unused portion ²⁴	250 mL NS , ½-NS ²³ mix by gentle inversion ²³	complete administration within 24 h F, 8 h RT ²³ if refrigerated, bring bag to RT prior to administration ²³	- do NOT shake ²³ - administer with 0.2 micron in-line filter ²³

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
azaCITIDine 100 mg (Celgene) (RT) no preservative ²⁵	4 mL SWI ²⁵ shake vigorously ²⁵ record time of reconstitution	25 mg/mL ²⁵	use within 45 min RT or 8 h F ²⁵	SC syringe ²⁵	45 min RT (including preparation time) or 8 h F ²⁵ refrigerate syringe immediately after preparation if not to be used within 45 min of reconstitution ²⁵ Refrigerated syringes ²⁵: - allow up to 30 min prior to administration to reach temperature of ~20-25°C - discard syringe if time elapsed at RT is greater than 30 min	- discard if contains large particles ²⁵ - re-suspend syringe contents before injection by vigorously rolling syringe between palms ²⁵ - if cold diluent reconstitution is used to extend stability, minimize exposure to RT; ensure proper refrigeration of diluent, reconstituted vial and final product ^{26,27}
	cold diluent reconstitution: 4 mL SWI at 2-8°C ^{26,27}	25 mg/mL ²⁵	12 h F ^{2,26,27}		22 h F ^{26,27}	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
azaCITIDine 100 mg (Dr. Reddy's) (RT) no preservative ²⁸	4 mL SWI ²⁸ shake vigorously ²⁸	25 mg/mL ²⁸	use within 45 min RT or 8 h F ²⁸	SC syringe ²⁸	45 min RT (including preparation time) or 8 h F ²⁸ refrigerate syringe immediately after preparation if not to be used within 45 min of reconstitution ²⁸ Refrigerated syringes ²⁸: • allow up to 30 min prior to administration to reach temperature of ~20-25°C • discard syringe if time elapsed at RT is greater than 30 min	- do not filter ²⁸ - discard if contains large particles ²⁸ - re-suspend syringe contents before injection by vigorously rolling syringe between palms ²⁸

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
azaCITIDine 100 mg (Hikma) (RT) no preservative ²⁹	4 mL SWI ²⁹ shake vigorously ²⁹	25 mg/mL ²⁹	use within 45 min RT or 8 h F ²⁹	SC syringe ²⁹	45 min RT (including preparation time) or 8 h F ²⁹ refrigerate syringe immediately after preparation if not to be used within 45 min of reconstitution ²⁹ Refrigerated syringes ²⁹: • allow up to 30 min prior to administration to reach temperature of ~20-25°C • discard syringe if time elapsed at RT is greater than 30 min	- do not filter ²⁹ - discard if contains large particles ²⁹ - re-suspend syringe contents before injection by vigorously rolling syringe between palms ²⁹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
BCG <i>(Tice strain)</i> (OncoTICE®) <u>intravesical</u> 50 mg (1 to 8 x 10 ⁸ CFU) (Merck Canada) (F)(PFL) no preservative ³⁰	1 mL preservative-free NS ³⁰ allow to stand for a few min; gently swirl to suspend ³⁰ do NOT shake ³⁰ record time of reconstitution	1 to 8×10 ⁸ CFU/vial ³⁰	2 h F ³⁰ **(PFL) ³⁰	transfer contents from vial to 50 mL syringe, rinse vial with 1 mL NS and transfer rinse solution to the 50 mL syringe, then qs up to 45 mL with NS ³⁰ if a CSTD is used: transfer contents from vial to 50 mL syringe and qs up to 45 mL with NS; do NOT rinse vial ³⁰	use within 2 h F of reconstitution ^{30,31} **(PFL) ³⁰	- auxiliary info: biohazard ³¹ - do NOT filter ³⁰ - do NOT shake ³⁰

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
BCG (Russian strain) (VERITY-BCG®) <u>intravesical</u> 40 mg (1 to 8 x 10 ⁸ CFU) (Verity) (F)(PFL) no preservative ³²	1 mL preservative-free NS ³² allow to stand for a few min; gently swirl to suspend ³² do NOT shake ³² record time of reconstitution	1 to 8×10 ⁸ CFU/vial ³²	2 h F ³² **(PFL) ³²	transfer contents from 1 st vial to 50 mL syringe, rinse vial with 1 mL NS and transfer rinse solution to the 50 mL syringe; then, repeat steps for 2 nd vial and qs up to 45 mL with NS ³²	use within 2 h F of reconstitution ^{31,32} **(PFL) ³²	- auxiliary info: biohazard ³¹ - TWO vials must be used to achieve the recommended full dose ³² - do NOT shake ³²
Belantamab mafodotin 30 mg/1.5 mL (GSK) (frozen)(PFL) do not shake no preservative ³³ (SAP)	n/a	20 mg/mL ³³	thaw up to 4 h RT, F before use ³³ once thawed: unpunctured vial: 10 d F ³³ once thawed: punctured vial: discard unused portion ^{31,33} **(PFL) ³³ do NOT shake ³³	0.2-2 mg/mL NS ³³ 250 mL* NS ³³	8 h RT ³³	- supplied as frozen liquid ³³ - recommended freezer temp ³³ is (-50°C to -15°C) - thawed drug cannot be refrozen ³³

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Belantamab mafodotin 100 mg (GSK) (F)(PFL) no preservative ³⁴ (SAP)	allow vial to stand at RT for 10 min before reconstitution ³⁵ 2 mL SWI ³⁴ swirl gently to mix; do NOT shake ³⁵	50 mg/mL ³⁴	use immediately after reconstitution ³⁴ discard unused portion ³⁴	0.2-2 mg/mL NS ³⁴ 250 mL * NS ³⁴ mix by gentle inversion; do NOT shake ³⁵	complete administration within 8 h RT ³⁴	- discard if particulate matter is present ³⁴
Belinostat 500 mg (Spectrum) (RT) no preservative ³⁶ (SAP)	9 mL SWI ³⁶	50 mg/mL ³⁶	12 h RT ³⁶	250 mL NS ³⁶	complete administration within 36 h RT ³⁶	- administer with 0.2 micron in-line filter ³⁶
Bendamustine 25 mg 100 mg (Natco) (RT)(PFL) no preservative ³⁷	25 mg: 5 mL SWI ³⁷ 100 mg: 20 mL SWI ³⁷ shake well; dissolves completely in 5 min ³⁷	5 mg/mL ³⁷	30 min ³⁷	0.2-0.6 mg/mL NS, D2.5-1½NS ³⁷ 100-500 mL†	complete administration within 24 h F, 3 h RT ³⁷	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Bendamustine 25 mg 100 mg (Eugia) (RT)(PFL) no preservative ³⁸	25 mg vial: 5 mL SWI ³⁸ 100 mg vial: 20 mL SW ³⁸ shake well; dissolves completely in 5 minutes ³⁸	5 mg/mL ³⁸	30 minutes ³⁸	0.2-0.6 mg/mL NS , D2.5-½NS ³⁸ 100-500 mL†	complete administration within 24 h F, 3 h RT ³⁸	
Bendamustine 25 mg 100 mg (Teva) (RT,F)(PFL) no preservative ³⁹	25 mg: 5 mL SWI ³⁹ 100 mg: 20 mL SW ³⁹ shake well; dissolves completely in 5 min ³⁹	5 mg/mL ³⁹	30 min ³⁹	0.2-0.6 mg/mL NS , D2.5-½NS ³⁹ 100-500 mL†	complete administration within 24 h F, 3 h RT ⁴⁰	
Bevacizumab (AVASTIN®) 100 mg/4 mL 400 mg/16 mL (Roche) (F)(PFL) do not shake no preservative ⁴¹	N/A	25 mg/mL ⁴¹	discard unused portion ⁴¹	1.4-16.5 mg/mL NS only ⁴¹ 100-250 mL†	48 h F, RT ⁴¹	- do NOT shake ⁴¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Bevacizumab (AYBINTIO®) 100 mg/4 mL 400 mg/16 mL (Organon/Samsung) (F)(PFL) do not shake no preservative ⁴²	N/A	25 mg/mL ⁴²	discard unused portion ⁴²	1.4-16.5 mg/mL NS only ⁴² 100-250 mL†	10 d F, 72 h RT ^{2,42}	- do NOT shake ⁴²
Bevacizumab (MVASI®) 100 mg/4 mL 400 mg/16 mL (Amgen) (F)(PFL) do not shake no preservative ⁴³	N/A	25 mg/mL ⁴³	discard unused portion ⁴³	1.4-16.5 mg/mL NS only ⁴³ 100-250 mL†	48 h F, RT ⁴³	- do NOT shake ⁴³
Bevacizumab (ZIRABEV®) 100 mg/4 mL 400 mg/16 mL (Pfizer) (F)(PFL) do not shake no preservative ⁴⁴	N/A	25 mg/mL ⁴⁴	discard unused portion ⁴⁴	1.4-16.5 mg/mL NS only ⁴⁴ 100-250 mL†	10 d F, 48 h RT ^{2,44}	- do NOT shake ⁴⁴

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Bleomycin 15 units (NB: dose in units only) (Fresenius Kabi) (F)(PFL) no preservative ⁴⁵	6 mL* NS ⁴⁵	2.5 units/mL	12 h F ^{2,45}	50 mL* NS ⁴⁵	24 h RT ⁴⁵	
Bleomycin 15 units (NB: dose in units only) (Pfizer/Hospira) (F)(PFL) no preservative ⁴⁶	6 mL* NS , SWI ⁴⁶	2.5 units/mL	12 h F, RT ^{2,46}	50 mL* NS ⁴⁶	4 h RT ^{2,31,46}	
Blinatumomab 38.5 mcg (Amgen) (F)(PFL) do not shake no preservative ⁴⁷	3 mL SWI ⁴⁷ do NOT use supplied IV solution stabilizer to reconstitute vials ⁴⁷ direct diluent against side of vial during reconstitution ⁴⁷ gently swirl to avoid excess foaming ⁴⁷	12.5 mcg/mL ⁴⁷	12 h F ^{2,48} , 4 h RT ⁴⁸	250 mL NS ⁴⁷ add supplied IV solution stabilizer to NS bag and gently mix to avoid foaming ⁴⁷ add reconstituted drug to bag following addition of IV solution stabilizer ⁴⁷	complete administration within 10 d F, 96 h RT ⁴⁸	- use non-DEHP bag and IV administration set ⁴⁷ - administer with 0.2 micron in-line filter ⁴⁷ - prime lines with blinatumomab solution; do NOT use NS

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Bortezomib 3.5 mg (Actavis) (RT)(PFL) no preservative ⁴⁹	3.5 mL NS ⁴⁹	1 mg/mL ⁴⁹	12 h F, RT ^{2,50}	IV syringe ⁴⁹	10 d F, 4 d RT ^{2,50}	- auxiliary info: WARNING: INTRAVENOUS use only. Fatal if given by other routes.
Bortezomib SC injection 3.5 mg (Actavis) (RT)(PFL) no preservative ⁴⁹	1.4 mL NS ⁴⁹	2.5 mg/mL ⁴⁹	12 h F, RT ^{2,50}	SC syringe ⁴⁹	10 d F, 4 d RT ^{2,50}	- auxiliary info: WARNING: SUBCUTANEOUS use only. Fatal if given by other routes.
Bortezomib 3.5 mg (Apotex) (RT)(PFL) no preservative ⁵¹	3.5 mL NS ⁵¹	1 mg/mL ⁵¹	12 h F, RT ^{2,52}	IV syringe ⁵¹	10 d F, 4 d RT ^{2,52}	- auxiliary info: WARNING: INTRAVENOUS use only. Fatal if given by other routes.

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Bortezomib SC injection 3.5 mg (Apotex) (RT)(PFL) no preservative ⁵¹	1.4 mL NS ⁵¹	2.5 mg/mL ⁵¹	12 h F, RT ^{2,52}	SC syringe ⁵¹	10 d F, 4 d RT ^{2,52}	- auxiliary info: WARNING: SUBCUTANEOUS use only. Fatal if given by other routes.
Bortezomib 3.5 mg (Janssen) (RT)(PFL) no preservative ⁵³	3.5 mL NS ⁵³	1 mg/mL ⁵³	12 h F, RT ^{2,50}	IV syringe ⁵³	10 d F, 4 d RT ^{2,50}	- auxiliary info: WARNING: INTRAVENOUS use only. Fatal if given by other routes.
Bortezomib SC injection 3.5 mg (Janssen) (RT)(PFL) no preservative ⁵³	1.4 mL NS ⁵³	2.5 mg/mL ⁵³	12 h F, RT ^{2,50}	SC syringe ⁵³	10 d F, 4 d RT ^{2,50}	- auxiliary info: WARNING: SUBCUTANEOUS use only. Fatal if given by other routes.

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Bortezomib 1 mg 2.5 mg 3.5 mg (Juno/MDA) (RT)(PFL) no preservative ⁵⁴	1 mg: 1 mL NS ⁵⁴ 2.5 mg: 2.5 mL NS ⁵⁴ 3.5 mg: 3.5 mL NS ⁵⁴	1 mg/mL ⁵⁴	12 h F, RT ^{2,55}	IV syringe ⁵⁴	10 d F, 4 d RT ^{2,55}	- auxiliary info: WARNING: INTRAVENOUS use only. Fatal if given by other routes.
Bortezomib SC injection 2.5 mg 3.5 mg (Juno/MDA) (RT)(PFL) no preservative ⁵⁴	2.5 mg: 1 mL NS ⁵⁴ 3.5 mg: 1.4 mL NS ⁵⁴	2.5 mg/mL ⁵⁴	12 h F, RT ^{2,55}	SC syringe ⁵⁴	10 d F, 4 d RT ^{2,55}	- auxiliary info: WARNING: SUBCUTANEOUS use only. Fatal if given by other routes.
Bortezomib 3.5 mg (Marcan) (RT)(PFL) no preservative ⁵⁶	3.5 mL NS ⁵⁶	1 mg/mL ⁵⁶	12 h F, RT ^{2,57,58}	IV syringe ⁵⁶	10 d F, 2 d RT ^{2,57,58}	- auxiliary info: WARNING: INTRAVENOUS use only. Fatal if given by other routes.

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Bortezomib SC injection 3.5 mg (Marcan) (RT)(PFL) no preservative ⁵⁶	1.4 mL NS ⁵⁶	2.5 mg/mL ⁵⁶	12 h F, RT ^{2,57,58}	SC syringe ⁵⁶	10 d F, 2 d RT ^{2,57,58}	- auxiliary info: WARNING: SUBCUTANEOUS use only. Fatal if given by other routes.
Bortezomib 3.5 mg (PMS) (RT)(PFL) no preservative ⁵⁹	3.5 mL NS ⁵⁹	1 mg/mL ⁵⁹	8 h RT ⁵⁹	IV syringe ⁵⁹	8 h RT ⁵⁹	- auxiliary info: WARNING: INTRAVENOUS use only. Fatal if given by other routes.
Bortezomib SC injection 3.5 mg (PMS) (RT)(PFL) no preservative ⁵⁹	1.4 mL NS ⁵⁹	2.5 mg/mL ⁵⁹	8 h RT ⁵⁹	SC syringe ⁵⁹	8 h RT ⁵⁹	- auxiliary info: WARNING: SUBCUTANEOUS use only. Fatal if given by other routes.

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Bortezomib 1 mg 2.5 mg 3.5 mg (Taro/Sun Pharma) (RT)(PFL) no preservative ⁶⁰	1 mg: 1 mL NS ⁶⁰ 2.5 mg: 2.5 mL NS ⁶⁰ 3.5 mg: 3.5 mL NS ⁶⁰	1 mg/mL ⁶⁰	8 h RT ⁶⁰	IV syringe ⁶⁰	8 h RT ⁶⁰	- auxiliary info: WARNING: INTRAVENOUS use only. Fatal if given by other routes.
Bortezomib SC injection 1 mg 2.5 mg 3.5 mg (Taro/Sun Pharma) (RT)(PFL) no preservative ⁶⁰	1 mg: 0.4 mL NS ⁶⁰ 2.5 mg: 1 mL NS ⁶⁰ 3.5 mg: 1.4 mL NS ⁶⁰	2.5 mg/mL ⁶⁰	8 h RT ⁶⁰	SC syringe ⁶⁰	8 h RT ⁶⁰	- auxiliary info: WARNING: SUBCUTANEOUS use only. Fatal if given by other routes.
Bortezomib 3.5 mg (Teva) (RT)(PFL) no preservative ⁶¹	3.5 mL NS ⁶¹	1 mg/mL ⁶¹	12 h F, RT ^{2,50}	IV syringe ⁶¹	10 d F, 4 d RT ^{2,50}	- auxiliary info: WARNING: INTRAVENOUS use only. Fatal if given by other routes.

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Bortezomib SC injection 3.5 mg (Teva) (RT)(PFL) no preservative ⁶¹	1.4 mL NS ⁶¹	2.5 mg/mL ⁶¹	12 h F, RT ^{2,50}	SC syringe ⁶¹	10 d F, 4 d RT ^{2,50}	- auxiliary info: WARNING: SUBCUTANEOUS use only. Fatal if given by other routes.
Brentuximab vedotin 50 mg (Seagen/Pfizer) (F)(PFL) no preservative ⁶²	10.5 mL SWI ⁶² direct diluent against side of vial during reconstitution ⁶² do NOT shake ⁶²	5 mg/mL ⁶²	12 h F ^{2,62}	0.4-1.8 mg/mL NS , D5W, LR ⁶² 50-100 mL† gently invert to mix ⁶²	24 h F ^{2,62}	- solution should be colorless, clear to slightly opalescent, and free of visible particulates ⁶²
Busulfan 60 mg/10 mL (PMS) (F) no preservative ⁶³	N/A	6 mg/mL ⁶³	discard unused portion ^{31,63}	dilute to volume 10 times drug volume to achieve final concentration of ~0.5 mg/mL NS , D5W ⁶³ 250-1000 mL†	complete administration within 12 h F, 8 h RT ⁶³	- contains DMA*** - always add busulfan to diluent to mix; do NOT add diluent to busulfan ⁶³

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Busulfan 60 mg/10 mL (SteriMax) (F) no preservative ⁶⁴	N/A	6 mg/mL ⁶⁴	discard unused portion ^{24,64}	dilute to volume 10 times drug volume to achieve final concentration of ~0.5 mg/mL NS , D5W ⁶⁴ 250-1000 mL†	in NS : complete administration within 12 h F, 8 h RT ⁶⁴ in D5W : complete administration within 8 h RT ⁶⁴	- contains DMA*** - always add busulfan to diluent to mix; do NOT add diluent to busulfan ⁶⁴
Cabazitaxel 60 mg/1.5 mL (Dr. Reddy's) (RT) no preservative ⁶⁵	supplied diluent: withdraw entire contents of diluent vial and inject into the concentrate vial ⁶⁵ slowly direct diluent against inside of vial to limit foaming ⁶⁵ mix by repeated inversions for 45 sec ⁶⁵ do NOT shake ⁶⁵ let sit for 5 min ⁶⁵	10 mg/mL ⁶⁵	1 h RT ⁶⁵	0.10-0.26 mg/mL NS , D5W ⁶⁵ 100-250 mL†	complete administration within 48 h F, 8 h RT ⁶⁵	- use non-DEHP bag and tubing ⁶⁵ - administer with 0.2 micron in-line filter ⁶⁵ - concentrate and diluent vials contain overfill ⁶⁵ - diluent contains 13% (w/w) ethanol in water ⁶⁵ - discard if crystallization occurs ⁶⁵

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Cabazitaxel 45 mg/4.5 mL 60 mg/6 mL (Formative) (RT) preservative ⁶⁶	N/A	10 mg/mL ⁶⁶	10 d F , RT ⁶⁶	0.10-0.26 mg/mL NS , D5W ⁶⁶ 100-250 mL†	complete administration within 48 F, 8 h RT ⁶⁶	- use non-DEHP bag and tubing ⁶⁶ - administer with 0.2 micron in-line filter ⁶⁶ - vials contains overflow ⁶⁶ - discard if crystallization occurs ⁶⁶
Cabazitaxel 45 mg/4.5 mL 60 mg/6 mL (Sandoz) (RT) preservative ⁶⁷	N/A	10 mg/mL ⁶⁷	10 d F , RT ⁶⁷	0.10-0.26 mg/mL NS , D5W ⁶⁷ 100-250 mL†	complete administration within 48 h F, 8 h RT ⁶⁷	- use non-DEHP bag and tubing ⁶⁷ - administer with 0.2 micron in-line filter ⁶⁷ - vials contain overflow ⁶⁷ - discard if crystallization occurs ⁶⁷

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Cabazitaxel 60 mg/1.5 mL (sanofi-aventis) (RT) no preservative ⁶⁸	supplied diluent: withdraw entire contents of diluent vial and inject into the concentrate vial ⁶⁸ slowly direct diluent against inside of vial to limit foaming ⁶⁸ mix by repeated inversions for 45 sec ⁶⁸ do NOT shake ⁶⁸ let sit for 5 min ⁶⁸	10 mg/mL ⁶⁸	1 h RT ⁶⁸	0.10-0.26 mg/mL NS, D5W ⁶⁸ 100-250 mL†	complete administration within 48 h F, 8 h RT ⁶⁸	- use non-DEHP bag and tubing ⁶⁸ - administer with 0.2 micron in-line filter ⁶⁸ - concentrate and diluent vials contain overfill ⁶⁸ - diluent contains 13% (w/w) ethanol in water ⁶⁸ - discard if crystallization occurs ⁶⁸

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Calaspargase pegol (pegylated asparaginase <i>E. coli</i>) 3750 units/5 mL (Servier) (F)(PFL) do not shake no preservative ⁶⁹	N/A	750 units/mL ⁶⁹	discard unused portion ⁶⁹	100 mL NS , D5W ⁶⁹	24 h F , 4 h RT ⁶⁹ **(PFL) ⁷⁰	- discard if discolouration, cloudiness, or visible particles are present ⁶⁹ - unopened vials may be stored at RT for 48 h ⁶⁹ - protect IV bag and tubing from light during administration ⁷⁰
CARBOplatin 50 mg/5 mL 150 mg/15 mL 450 mg/45 mL 600 mg/60 mL (Accord) (RT)(PFL) no preservative ⁷¹	N/A	10 mg/mL ⁷¹	discard unused portion ⁷¹	0.5-10 mg/mL NS , D5W ⁷¹ 50-250 mL†	24 h F , 8 h RT ⁷¹	- do NOT use aluminum- containing needle, syringe, or tubing ⁷¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
CARBOplatin 50 mg/5 mL 150 mg/15 mL 450 mg/45 mL 600 mg/60 mL (Omega) (RT)(PFL) no preservative ⁷²	N/A	10 mg/mL ⁷²	discard unused portion ⁷²	0.3-10 mg/mL NS , D5W ⁷² 50-250 mL†	48 h F ⁷² , 24 h RT ⁷³	- do NOT use aluminum- containing needle, syringe or tubing ⁷²
CARBOplatin 50 mg/5 mL 150 mg/15 mL 450 mg/45 mL 600 mg/60 mL (Pfizer/Hospira) (RT)(PFL) no preservative ⁷⁴	N/A	10 mg/mL ⁷⁴	discard unused portion ⁷⁴	0.3-10 mg/mL NS , D5W ⁷⁴ 50-250 mL†	48 h F ⁷⁴	- do NOT use aluminum- containing needle, syringe, or tubing ⁷⁴
CARBOplatin 50 mg/5 mL 150 mg/15 mL 450 mg/45 mL (Teva) (RT)(PFL) no preservative ⁷⁵	N/A	10 mg/mL ⁷⁵	discard unused portion RT ⁷⁵	0.5-10 mg/mL ⁷⁶ NS , D5W ^{75,77,78} 50-250 mL†	8 h F ⁷⁹ , RT ⁷⁵	- do NOT use aluminum- containing needle, syringe, or tubing ⁷⁵

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Carfilzomib 10 mg 30 mg 60 mg (Amgen) (F)(PFL) no preservative ⁸⁰	10 mg: 5 mL SWI ⁸⁰ 30 mg: 15 mL SWI ⁸⁰ 60 mg: 29 mL SWI ⁸⁰ direct diluent against side of vial during reconstitution ⁸⁰ swirl gently; do NOT shake ⁸⁰ if foaming occurs, allow to settle until clear (~5 min) ⁸⁰	2 mg/mL ⁸⁰	12 h F, 4 h RT ^{2,80}	50-100 mL* D5W only ⁸⁰ do NOT dilute in NS ⁸⁰	24 h F, 4 h RT ^{2,80}	- if a CSTD is not used during compounding, a 21 gauge (or larger gauge) needle is recommended to prevent coring of the stopper ⁸¹⁻⁸³ - do not use NS for reconstitution or dilution ⁸⁰ - discard if contains particulates ⁸⁰

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Carmustine (note concentration) 100 mg/1 mL 300 mg/3 mL (Marcan) (F)(PFL) no preservative ⁸⁴	N/A	100 mg/mL ⁸⁴	discard unused portion ^{2,84} crystals may be re-dissolved by warming the vial to RT with shaking ⁸⁴ (PFL) ⁸⁴	0.2-3.06 mg/mL NS, D5W ^{84,85} 500-1000 mL† in glass or polypropylene containers ONLY ⁸⁴	10 h RT ⁸⁶ or 48 h F plus an additional 8 h RT ⁸⁶ **(PFL) ^{84,86}	- CAUTION: alternative concentration - unstable in PVC containers and tubing ⁸⁴ - administer with PVC-free infusion set ⁸⁴ - protect from light for administration ⁸⁴
Carmustine (note concentration) 100 mg (Marcan) (F) no preservative ⁸⁴	3 mL supplied diluent ⁸⁴ bring drug and diluent vials to RT prior to mixing ⁸⁴ completely dissolve drug in diluent; solution should be clear when fully dissolved ⁸⁴	33.3 mg/mL ⁸⁴	20 d F, 24 h RT ⁸⁴ crystals may be re-dissolved by warming the vial to RT with shaking ⁸⁴ (PFL) ⁸⁴	0.2-3.06 mg/mL NS, D5W ^{84,87} 500 - 1000 mL† in glass or polypropylene containers ONLY ⁸⁴	8 h RT ⁸⁸ or 48 h F plus an additional 6 h RT ⁸⁸ **(PFL) ⁸⁸	- CAUTION: alternative concentration after reconstitution - supplied diluent is propylene glycol ⁸⁴ - do not use vial if oily film is present - unstable in PVC containers and tubing ⁸⁴ - administer with PVC-free infusion set ⁸⁴ - protect from light for administration ⁸⁴

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Carmustine 100 mg (SteriMax) (F) no preservative ⁸⁹	3mL supplied diluent ⁸⁹ bring drug and diluent vials to RT prior to mixing ⁸⁹ completely dissolve drug in diluent, then add 27 mL SWI ⁸⁹	3.3 mg/mL in ethanol 10% ⁸⁹	48 h F ⁸⁹ precipitates can be re-dissolved by warming the vial to RT with gentle shaking ⁸⁹	500 mL NS, D5W ⁸⁹ in glass or polypropylene containers ONLY ⁸⁹	8 h RT ⁸⁹ or 48 h F plus an additional 6 h RT ⁸⁹ **(PFL) ⁸⁹	- supplied diluent is dehydrated alcohol ⁸⁹ - do not use vial if oily film is present ⁸⁹ - to remix bag contents prior to administration, gently shake final product for ~10 sec ⁸⁹ - administer with PVC-free infusion set ⁸⁹ - protect from light for administration ⁸⁹
Cemiplimab 350 mg/7 mL (Regeneron) (F)(PFL) do not shake no preservative ⁹⁰	N/A	50 mg/mL ⁹⁰	discard unused portion ⁹⁰	1-20 mg/mL NS, D5W ⁹⁰ 50 mL† mix by gentle inversion ⁹⁰	complete administration within 10 d F, 8 h RT ⁹⁰ bring to RT prior to use ⁹⁰	- administer with 0.2-5 micron filter ^{2,90} - solution may contain white particulates which do not affect product quality ⁹⁰

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Cemiplimab 250 mg/5 mL 350 mg/7 mL (sanofi) (F)(PFL) do not shake no preservative ⁹¹	N/A	50 mg/mL ⁹¹	discard unused portion ^{31,91}	1-20 mg/mL NS , D5W ⁹¹ 50 mL† mix by gentle inversion	complete administration within 24 h F, 8 h RT ⁹¹	- administer with 0.2-5 micron filter ⁹¹ - solution may contain white particulates which do not affect product quality ⁹¹
Cetuximab 100 mg/50 mL 200 mg/100 mL (Imclone/Lilly) (F) do not shake no preservative ⁹²	N/A	2 mg/mL ⁹²	12 h F, 8 h RT ⁹²	syringe ⁹²	12 h F, 8 h RT ⁹²	- administer with 0.2 micron filter ⁹² - solution may contain white particulates which do not affect product quality ⁹²
				evacuated container or bag ⁹²		

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
CISplatin 10 mg/10 mL 50 mg/50 mL 100 mg/100mL (Accord) (RT)(PFL) no preservative ⁹³	N/A	1 mg/mL ⁹³	discard unused portion ³¹	NS ⁹³ 100-500 mL† or 2 L D5-½NS or D5-⅓NS containing 37.5 g of mannitol ⁹³	24 h RT ⁹³	- do NOT use aluminum- containing needle, syringe or tubing ⁹³ - suggested dose limits relate to the physical limitations of the bag size and added drug volume; it is not a concentration- dependent property of the drug - for ULY0 D-PACE protocol, see entry for DPACE (3-in-1 solution containing etoposide, CISplatin, cyclophosphamide)

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
CISplatin 50 mg/50 mL 100 mg/100mL (Pfizer/Hospira) (RT)(PFL) no preservative ⁹⁴	N/A	1 mg/mL ⁹⁴	discard unused portion ³¹	NS ⁹⁴ 100-500 mL† or 2 L D5-½NS or D5-⅓NS containing 37.5 g of mannitol ⁹⁴	24 h RT ⁹⁴	- do NOT use aluminum- containing needle, syringe or tubing ⁹⁴ - suggested dose limits relate to the physical limitations of the bag size and added drug volume; it is not a concentration- dependent property of the drug - for ULY0 D-PACE protocol, see entry for DPACE (3-in-1 solution containing etoposide, CISplatin, cyclophosphamide)

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
CISplatin 10 mg/10 mL 50 mg/50 mL 100 mg/100mL (Sandoz) (RT)(PFL) no preservative ⁹⁵	N/A	1 mg/mL ⁹⁵	12 h RT ^{2,96}	NS ⁹⁵ 100-500 mL† or 2 L D5-½NS or D5-⅓NS containing 37.5 g of mannitol ⁹⁵	24 h RT ⁹⁶	- do NOT use aluminum- containing needle, syringe or tubing ⁹⁵ - suggested dose limits relate to the physical limitations of the bag size and added drug volume; it is not a concentration- dependent property of the drug - for ULY0 D-PACE protocol, see entry for DPACE (3-in-1 solution containing etoposide, CISplatin, cyclophosphamide)

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
CISplatin 10 mg/10 mL 50 mg/50 mL 100 mg/100mL (Teva) (RT)(PFL) no preservative ⁹⁷	N/A	1 mg/mL ⁹⁷	discard unused portion ²⁴	NS ⁹⁷ 100-500 mL† or 2 L D5-½NS or D5-⅓NS containing 37.5 g of mannitol ⁹⁷	24 h RT ⁹⁷	- do NOT use aluminum- containing needle, syringe or tubing ⁹⁷ - suggested dose limits relate to the physical limitations of the bag size and added drug volume; it is not a concentration- dependent property of the drug - for ULY0 D-PACE protocol, see entry for DPACE (3-in-1 solution containing etoposide, CISplatin, cyclophosphamide)

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Cladribine 10 mg/10 mL (Fresenius Kabi) (F)(PFL) no preservative ⁹⁸	N/A	1 mg/mL ⁹⁸	discard unused potion ⁹⁸	SC syringe ⁹⁹	48 h F, discard end of day RT ^{31,100,101}	
				500 mL NS only ⁹⁸ do NOT use D5W ⁹⁸	24 h RT ⁹⁸	
				Cassette: qs to 100 mL with bacteriostatic NS only via SIMS DELTEC INC. MEDICATION CASSETTES® ⁹⁸ filter drug and diluent through 0.22 micron filter as each solution is being introduced into the cassette ⁹⁸	at least 7 days ⁹⁸	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Cladribine 10 mg/10 mL (GMP) (F)(PFL) no preservative ¹⁰²	N/A	1 mg/mL ¹⁰²	discard unused portion ^{31,102}	SC syringe ⁹⁹	48 h F, discard end of day RT ^{31,100,101}	
				500 mL NS only ¹⁰² do NOT use D5W ¹⁰²	24 h RT ¹⁰²	
				Cassette: qs to 100 mL with bacteriostatic NS only via SIMS DELTEC INC. MEDICATION CASSETTES® ¹⁰² filter drug and diluent through 0.22 micron filter as each solution is being introduced into the cassette ¹⁰²	at least 7 days ¹⁰²	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Crisantaspase recombinant (asparaginase <i>Erwinia chrysanthemum</i>) 10 mg/0.5 mL (Jazz) (F)(PFL) do not shake preservative free ¹⁰³	N/A	20 mg/mL ¹⁰³	discard unused portion ¹⁰³	IM syringe ¹⁰³ max volume: 2 mL if volume >2 mL, divide volume into separate syringes for administration ¹⁰³	use within 4 h RT ¹⁰³	- discard if cloudy, discoloured, or contains particulates ¹⁰³ - do NOT shake ¹⁰³

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Cyclophosphamide 200 mg 500 mg 1000 mg 2000 mg (Baxter) (RT)(PFL) no preservative ¹⁰⁴	200 mg ¹⁰⁴ : 10 mL NS 500 mg ¹⁰⁴ : 25 mL NS 1000 mg ¹⁰⁴ : 50 mL NS 2000 mg ¹⁰⁴ : 100 mL NS	20 mg/mL ¹⁰⁴	12 h F, RT ^{2,104}	NS, D5W, D5NS ¹⁰⁴ 100-250 mL† high dose in BMT: may need 500 mL*	36 h F, 24 h RT ¹⁰⁵⁻¹⁰⁷	- suggested dose limits relate to the physical limitations of the bag size and added drug volume; it is not a concentration-dependent property of the drug - for ULY0 D-PACE protocol, see entry for DPACE (3-in-1 solution containing etoposide, CISplatin, cyclophosphamide) - do NOT use aluminum-containing needle, syringe, or tubing ¹⁰⁴

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Cyclophosphamide 500 mg 1000 mg 2000 mg (SteriMax) (RT)(PFL) no preservative ¹⁰⁸	500 mg ¹⁰⁸ : 25 mL NS 1000 mg ¹⁰⁸ : 50 mL NS 2000 mg ¹⁰⁸ : 100 mL NS	20 mg/mL ¹⁰⁸	12 h F, RT ¹⁰⁸	NS, D5W, D5NS ¹⁰⁸ 100-250 mL† high dose in BMT: may need 500 mL*	8 h F, RT ¹⁰⁹	- suggested dose limits relate to the physical limitations of the bag size and added drug volume; it is not a concentration-dependent property of the drug - for ULY0 D-PACE protocol, see entry for DPACE (3-in-1 solution containing etoposide, CISplatin, cyclophosphamide) - do NOT use aluminum-containing needle, syringe, or tubing ¹⁰⁸
Cytarabine 1000 mg/10mL 2000 mg/20mL (Pfizer/Hospira) (RT)(PFL) no preservative ¹¹⁰	N/A	100 mg/mL ¹¹⁰	12 h RT ^{2,110}	0.1-37.5 mg/mL NS, D5W, SWI ¹¹⁰ 100 mL†	in NS: 4 d RT ^{2,110} other solutions: 72 h F, 24 h RT ¹¹⁰ **(PFL) ¹¹⁰	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Cytarabine IT injection 1000 mg/10mL 2000 mg/20mL (Pfizer/Hospira) (RT)(PFL) no preservative ¹¹⁰	N/A record time of puncture	100 mg/mL ¹¹⁰	use within 4 h of initial puncture ²	IT syringe qs to 6 mL with preservative free NS ¹¹¹⁻¹¹³ diluent containing preservatives should NOT be used for intrathecal administration ¹¹⁴	use within 4 h of initial puncture ² **(PFL) ¹¹⁰	- auxiliary info ² : IT - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag ¹¹³
Cytarabine SC injection 1000 mg/10mL 2000 mg/20mL (Pfizer/Hospira) (RT)(PFL) no preservative ¹¹⁰	N/A	100 mg/mL ¹¹⁰	12 h RT ^{2,110}	SC syringe	10 d F, 4 d RT ^{2,115-117} **(PFL) ¹¹⁰	
Cytarabine 1000 mg/10mL 2000 mg/20mL (PMS) (RT)(PFL) no preservative ¹¹⁸	N/A	100 mg/mL ¹¹⁸	discard unused portion ^{31,118}	0.1-37.5 mg/mL NS, D5W, SWI ¹¹⁸ 100 mL†	10 d F, 48 h RT ¹¹⁸ **(PFL)	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Cytarabine IT injection 1000 mg/10mL 2000 mg/20mL (PMS) (RT)(PFL) no preservative ¹¹⁸	N/A record time of puncture	100 mg/mL ¹¹⁸	use within 4 h of initial puncture ³¹	IT syringe qs to 6 mL with preservative free NS ^{111,112} diluent containing preservatives should NOT be used for intrathecal administration ¹¹⁴	use within 4 h of initial puncture ³¹ **(PFL)	- auxiliary info: IT ³¹ - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag ¹¹³
Cytarabine SC injection 1000 mg/10mL 2000 mg/20mL (PMS) (RT)(PFL) no preservative ¹¹⁸	N/A	100 mg/mL ¹¹⁸	discard unused portion ^{31,118}	SC syringe	10 d F, 48 h RT ¹¹⁸ **(PFL)	
Cytarabine 2000 mg/20mL (SteriMax) (RT)(PFL) no preservative ¹¹⁹	N/A	100 mg/mL ¹¹⁹	12 h RT ^{2,119}	0.1-37.5 mg/mL NS , D5W, SWI, LR ¹¹⁹ 100 mL*	in NS: 4 d RT ^{2,119} other solutions: 72 h F, 24 h RT ¹¹⁹ **(PFL) ¹¹⁹	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Cytarabine IT injection 2000 mg/20mL (SteriMax) (RT)(PFL) no preservative ¹¹⁹	N/A record time of puncture	100 mg/mL ¹¹⁹	use within 4 h of initial puncture ²	IT syringe qs to 6 mL with preservative free NS ¹¹¹⁻¹¹³ diluent containing preservatives should NOT be used for intrathecal administration ¹¹⁴	use within 4 h of initial puncture ² **(PFL) ¹¹⁹	- auxiliary info: IT ² - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag ¹¹³
Cytarabine SC injection 2000 mg/20mL (SteriMax) (RT)(PFL) no preservative ¹¹⁹	N/A	100 mg/mL ¹¹⁹	12 h RT ^{2,119}	SC syringe	10 d F, 4 d RT ^{2,115-117} **(PFL) ¹¹⁹	
Dacarbazine 600 mg (Pfizer) (F)(PFL) no preservative ¹²⁰	59.1 mL SWI ¹²⁰	10 mg/mL ¹²⁰	12 h F, 8 h RT ^{2,120}	0.19-3.0 mg/mL NS , D5W ¹²⁰ 500-1000 mL†	24 h F ¹²⁰ **(PFL) ¹²¹	- protect container from light during administration ¹²¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
DACTINomycin 0.5 mg (GMD Pharma for Recordati) (RT)(PFL) no preservative ¹²² (SAP)	1.1 mL SWI ¹²² do NOT use SWI with preservative (may form precipitate) ¹²²	0.5 mg/mL (500 mcg/mL) ¹²²	discard unused portion ¹²³	syringe ¹²²	use within 4 h of initial vial puncture ¹²³	- drug loss reported with some cellulose ester membrane in- line filters ¹²²
				10 mcg/mL or greater ¹²² NS, D5W ^{122,124}		
Daratumumab 100 mg/5mL 400 mg/20mL (Janssen) (F)(PFL) do not shake no preservative ¹²⁵	N/A	20 mg/mL ¹²⁵	discard unused portion ¹²⁵	500-1000 mL NS dilute to final volume by withdrawing volume from bag equal to volume of drug to be added ¹²⁵ mix by gentle inversion ¹²⁵	24 h F , followed by 15 h infusion (total 39 h) ¹²⁵ allow bag to come to RT, then use immediately ¹²⁵ **(PFL)	- administer with 0.2 micron in-line filter ¹²⁵ - discard if visible particles are observed ¹²⁵ - complete infusion within 15 h ¹²⁵

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Daratumumab subcutaneous (DARZALEX SC®) 1800 mg/15 mL (Janssen) (F)(PFL) do not shake no preservative ¹²⁶	N/A	120 mg/mL ¹²⁶ allow vial to come to RT prior to use ¹²⁶	discard unused portion ^{2,126}	SC syringe ¹²⁶	24 h F, plus an additional 12 h RT ¹²⁶ bring to RT prior to use ¹²⁶	- contains hyaluronidase ¹²⁶ - formulations are NOT interchangeable ¹²⁶ - discard if opaque particles or discolouration are present ¹²⁶ - unpunctured vial may be stored up to 24 h at RT ¹²⁶
Datopotamab deruxtecan 100 mg (Daiichi) (F)(PFL) no preservative ¹²⁷	5 mL SWI ¹²⁷ gently swirl to mix do not shake ¹²⁷ record time of reconstitution ¹²⁷	20 mg/mL ¹²⁷	discard unused portion ¹²⁷	0.1-6.7 mg/mL D5W ONLY ¹²⁷ 100 mL* gently invert to mix do not shake ¹²⁷ do NOT use sodium chloride solution ¹²⁷	24 h F from initial vial puncture ¹²⁷ **(PFL) ¹²⁷ bring to RT prior to use ¹²⁷ MAX time at RT = 4.5 h from initial vial puncture (including preparation, storage at RT, and administration time) ¹²⁷	- NOT compatible with saline ¹²⁷ - administer with 0.2 micron in-line filter ¹²⁷ - protect infusion bag from light for administration ¹²⁷

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
DAUNOrubicin 20 mg (Erfa) (RT)(PFL) no preservative ¹²⁸	4 mL SWI ¹²⁸	5 mg/mL ¹²⁸	12 h F, RT ^{2,128} **(PFL) ¹²⁸	100-250 mL NS, D5W ¹²⁸	48 h F, 24 h RT ¹²⁹ **(PFL) ¹²⁸	
DAUNOrubicin 20 mg/4 mL (Hikma) (F)(PFL) no preservative ¹³⁰	N/A	5 mg/mL ¹³⁰	discard unused portion ²	100-250 mL ¹³⁰ NS, D5W	7 d F, 30 h RT ¹³¹ **(PFL) ¹³⁰	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Daunorubicin- cytarabine liposome 44 mg-100 mg (Jazz) (F)(PFL) no preservative ¹³²	19 mL SWI ¹³² allow vial to come to RT for 30 min prior to use ¹³² swirl gently for 5 min, inverting the vial every 30 sec; do NOT shake ¹³² allow vial to rest for 15 min after reconstitution ¹³² gently invert each vial 5 times prior to withdrawing concentrate for dilution ¹³² record time of reconstitution	2.2 mg/mL daunorubicin- 5 mg/mL cytarabine ¹³²	4 h F ¹³² max <i>combined</i> storage time for reconstituted vial and diluted product is 4 h F ¹³² (NOT 4 h F each)	500 mL NS, D5W ¹³² mix by gentle inversion ¹³²	use within 4h F ¹³² of initial vial puncture max <i>combined</i> storage time for reconstituted vial and diluted product is 4 h F ¹³² (NOT 4 h F each)	- reconstituted product is an opaque, purple, homogenous dispersion ¹³² - before administration, final product should be gently inverted to remix solution after refrigeration ¹³²

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Degarelix 80 mg 120 mg (Ferring) (RT) do not shake ¹³³ no preservative ¹³⁴	80 mg: 4.2 mL SWI (supplied diluent) ¹³³	20 mg/mL ¹³³	2 h RT ¹³³	SC syringe ¹³³	2 h RT ¹³³	
	120 mg: 3 mL SWI (supplied diluent) ¹³³	40 mg/mL ¹³³				
	swirl gently; avoid shaking to prevent foam formation ¹³³ reconstitution may take up to 15 min ¹³³					

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Denosumab (WYOST®) 120 mg/1.7 mL (Sandoz) (F)(PFL) do not shake no preservative ¹³⁵	N/A	71 mg/mL ¹³⁵	discard unused portion ¹³⁵	SC syringe ¹³⁵	use within 4 h F, RT of initial puncture ¹²³ bring to RT 15-30 min prior to use ¹³⁵	- not interchangeable with PROLIA® or JUBBONTI® - do not use if solution is cloudy or contains visible particles ¹³⁵ - avoid vigorous shaking ¹³⁵ - use a 27 gauge needle to withdraw drug from vial ¹³⁵

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Denosumab (XGEVA®) 120 mg/1.7 mL (Amgen) (F)(PFL) do not shake no preservative ¹³⁶	N/A	71 mg/mL ¹³⁶	discard unused portion ^{123,136}	SC syringe ¹³⁶	use within 4 h F, RT of initial puncture ¹²³ bring to RT 15-30 min prior to use ¹³⁶	- not interchangeable with PROLIA® ¹³⁶ or JUBBONTI® - do not use if solution is cloudy ¹³⁶ - trace amounts of translucent to white proteinaceous particles are acceptable ¹³⁶ - avoid vigorous shaking ¹³⁶ - use a 27 gauge needle to withdraw drug from vial ¹³⁷
Dexrazoxane 250 mg 500 mg (Juno) (RT) no preservative ¹³⁸	250 mg ¹³⁸ : 25 mL SWI 500 mg ¹³⁸ : 50 mL SWI	10 mg/mL ¹³⁸	3 h F, 30 min RT ¹³⁸	dilute with LR in empty infusion bag to final concentration of 1.3-3.0 mg/mL ¹³⁸ qs to 110-600 mL†	4 h F, 1 h RT ¹³⁸	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Dexrazoxane 250 mg 500 mg (Pfizer) (RT) no preservative ¹³⁹	250 mg: 25 mL SWI ¹³⁹ 500 mg: 50 mL SWI ¹³⁹	10 mg/mL ¹³⁹	3 h F, 30 min RT ¹³⁹	dilute with LR in empty infusion bag to final concentration of 1.3-3.0 mg/mL ¹³⁹ qs to 110-600 mL†	4 h F, 1 h RT ¹³⁹	
Dinutuximab 17.5 mg/5 mL (Unither/United Therapies) (F)(PFL) do not shake no preservative ¹⁴⁰	N/A	3.5 mg/mL ¹⁴⁰	discard unused portion ³¹	100 mL NS ¹⁴⁰ mix by gentle inversion ¹⁴⁰	initiate infusion within 4 h of dilution; refrigerate bag if not hung immediately ¹⁴⁰ complete administration within 24 h of dilution ¹⁴⁰	- do NOT shake ¹⁴⁰

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
DOCetaxel 20 mg/2 mL 80 mg/8 mL 160 mg/16 mL (Pfizer/Hospira) (F, RT)(PFL) preservative ¹⁴¹	N/A	10 mg/mL ¹⁴¹	20mg: discard unused portion ^{2,141} 80 mg or 160 mg: 28 d F ^{2,141} **(PFL) ¹⁴¹ (max number of punctures: up to 3 doses can be removed when a filtered venting needle [e.g., Chemo- Vent®] is also inserted, i.e., 6 punctures total) ¹⁴²	0.3-0.74 mg/mL NS, D5W ¹⁴¹ 25-500 mL†	10 d F, 4 d RT ^{2,143} **(PFL) ¹⁴³ during F storage	- use non-DEHP bag and IV administration sets ¹⁴¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
DOCEtaxel intravesical 20 mg/2 mL 80 mg/8 mL 160 mg/16 mL (Pfizer/Hospira) (F, RT)(PFL) preservative ¹⁴¹	N/A	10 mg/mL ¹⁴¹	20 mg: discard unused portion ^{2,141} 80 mg or 160 mg: 28 d F ^{2,141} **(PFL) ¹⁴¹ (max number of punctures: up to 3 doses can be removed when a filtered venting needle [e.g., Chemo- Vent®] is also inserted, i.e., 6 punctures total) ¹⁴²	syringe dilute with NS to final volume of 45 mL ^{144,145}	up to 0.9 mg/mL: 10 d F, 4 d RT ^{2,143} **(PFL) ¹⁴³ during F storage	
DOCEtaxel 20 mg/2 mL 80 mg/8 mL 160 mg/16 mL (Sandoz) (F,RT)(PFL) preservative ¹⁴⁶	N/A	10 mg/mL ¹⁴⁶	28 d F, RT ^{2,147}	0.3-0.74 mg/mL NS, D5W ¹⁴⁶ 25-500 mL†	24 h F, 4 h RT ^{2,148}	- use non-DEHP bag and IV administration sets ¹⁴⁶

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
DOCEtaxel intravesical 20 mg/2 mL 80 mg/8 mL 160 mg/16 mL (Sandoz) (F,RT)(PFL) preservative ¹⁴⁶	N/A	10 mg/mL ¹⁴⁶	28 d F, RT ^{2,147}	syringe dilute with NS to final volume of 45 mL ^{144,145}	up to 0.9 mg/mL ^{149,150} ; use immediately after preparation to prevent particle formation ^{2,148}	- particle formation occurs earlier with higher temperature and higher concentrations ¹⁴⁸
Dostarlimab 500 mg/10 mL (Glaxo) (F)(PFL) no preservative ¹⁵¹	N/A	50 mg/mL ¹⁵¹	discard unused portion ¹⁵¹	2-10 mg/mL NS, D5W ¹⁵¹ 100 mL* mix by gentle inversion ¹⁵¹	complete administration within 24 h F, 6 h RT ¹⁵¹ if refrigerated, bring bag to RT prior to administration ¹⁵¹	- do not shake ¹⁵¹ - discard if visible particles are present ¹⁵¹ - administer with 0.2 micron in-line filter ¹⁵¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
DOXOrubicin 10 mg/5 mL 20 mg/10 mL 50 mg/25 mL 200 mg/100 mL (Accord) (F)(PFL) no preservative ¹⁵²	N/A	2 mg/mL ¹⁵²	8 h ¹⁵²	syringe ¹⁵²	24 h F, RT from initial vial puncture ¹⁵²	- for LYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing either etoposide or etoposide phosphate AND DOXOrubicin and vinCRISTine)
				0.01–2 mg/mL NS ^{153,154} 1000 mL ¹⁵⁵⁻¹⁵⁷	24 h RT ^{153,154}	
DOXOrubicin 10 mg/5 mL 20 mg/10 mL 50 mg/25 mL 200 mg/100 mL (Teva) (F)(PFL) no preservative ¹⁵⁸	N/A	2 mg/mL ¹⁵⁸	8 h ¹⁵⁸	syringe ¹⁵⁸	48 h F, 24 h RT ¹⁵⁸ from initial vial puncture	- for LYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing either etoposide or etoposide phosphate AND DOXOrubicin and vinCRISTine)
				0.01–2 mg/mL NS ^{153,154} 1000 mL ¹⁵⁵⁻¹⁵⁷	24 h RT ^{153,154}	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
DOXOrubicin 10 mg/5 mL 50 mg/25 mL 200 mg/100 mL (Pfizer) (F) no preservative ¹⁵⁹	N/A	2 mg/mL ¹⁵⁹	discard unused portion ^{123,159}	syringe ¹⁵⁹	48 h F, 24 h RT ¹⁵⁹	- for LYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing either etoposide or etoposide phosphate AND DOXOrubicin and vinCRISTine)
				0.01–2 mg/mL NS ^{153,154} 1000 mL ¹⁵⁵⁻¹⁵⁷	24 h RT ^{153,154}	
DOXOrubicin Pegylated Liposomal 20 mg/10 mL (Janssen) (F) no preservative ¹⁶⁰	N/A	2 mg/mL ¹⁶⁰	discard unused portion ¹⁶⁰	D5W only ¹⁶⁰ <90 mg ¹⁶⁰ : 250 mL ≥90 mg ¹⁶⁰ : 500mL	24 h F ¹⁶⁰	- do not filter ¹⁶⁰
DOXOrubicin Pegylated Liposomal 20 mg/10 mL 50 mg/25 mL (Taro/Sun Pharma) (F) no preservative ¹⁶¹	N/A	2 mg/mL ¹⁶¹	discard unused portion ¹⁶¹	D5W only ¹⁶¹ <90 mg ¹⁶¹ : 250 mL ≥90 mg ¹⁶¹ : 500mL	24 h F ¹⁶¹	- do not filter ¹⁶¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
DOXOrubicin Pegylated Liposomal 20 mg/10 mL 50 mg/25 mL (Sun Pharma ¹⁶² USA) (F) no preservative ^{161,163}	N/A	2 mg/mL ¹⁶¹	discard unused portion ¹⁶¹	D5W only ¹⁶¹ <90 mg ¹⁶¹ : 250 mL ≥90 mg ¹⁶¹ : 500mL	24 h F ¹⁶¹	- do not filter ¹⁶¹ - discard if discoloured or contains particulates ¹⁶¹
DPACE (ULY0D-PACE protocol) (RT) no preservative ^{2,157,164,165}	see brand specific entries for: cyclophosphamide as applicable	see brand specific entries for: CISplatin, cyclophosphamide, etoposide	see brand specific entries for: CISplatin, cyclophosphamide, etoposide	in 1000 mL NS ^{156,164,165}	≤0.2 mg/mL: 24 h RT ^{2,164,165}	- final product is a 3-in-1 solution containing etoposide, CISplatin, cyclophosphamide (see ULY0D-PACE protocol) - use non-DEHP bag and tubing only - administer with 0.2 micron in-line filter

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Durvalumab 120 mg/2.4 mL 500 mg/10 mL (AstraZeneca) (F)(PFL) do not shake no preservative ¹⁶⁶	N/A	50 mg/mL ¹⁶⁶	discard unused portion ¹⁶⁶	1-15 mg/mL NS, D5W ¹⁶⁶ 100 mL† mix by gentle inversion ¹⁶⁶	10 d F, 12 h RT ^{2,166}	- do NOT shake ¹⁶⁶ - administer with 0.2 micron in-line filter ¹⁶⁶ - discard vial if solution is cloudy, discolored, or visible particles are present ¹⁶⁶ - use filtered venting needle (e.g., Chemo- Vent®) in place of CSTD for compounding ¹⁶⁷
Elranatamab 44 mg/1.1 mL 76 mg/1.9 mL (Pfizer) (F)(PFL) do not shake no preservative ¹⁶⁸	N/A	40 mg/mL ¹⁶⁸ allow vials to reach RT before using ¹⁶⁸	discard unused portion ¹⁶⁸	SC syringe ¹⁶⁸	use within 4 h F, RT ¹⁶⁸	- do not use if contains particulates ¹⁶⁸

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Elranatamab 76 mg/1.9 mL (Pfizer) (F)(PFL) no preservative ¹⁶⁹ (SAP)	N/A	40 mg/mL ¹⁶⁹ allow vials up to 15 min to reach RT before using ¹⁶⁹	discard unused portion ^{2,169}	SC syringe ¹⁶⁹	use immediately after preparation ^{2,169}	- supplied diluent to be used only for doses <8 mg ¹⁶⁹ - solution colour may be colourless to yellow/brown ¹⁶⁹ - unpunctured vials can be kept at RT up to 8 h before returning to F; discard if longer than 8 h RT ¹⁶⁹ - solutions can be prepared in normal room light; avoid direct sunlight ¹⁶⁹ - CSTD cannot be used during storage of prepared doses ^{169,170} - to prepare 76 mg dose ONLY: use filtered venting needle (e.g., Chemo-Vent®) in place of CSTD ¹⁷¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Enfortumab vedotin 20 mg 30 mg (Seagen) (F)(PFL) do not shake no preservative ¹⁷²	20 mg ¹⁷² ; 2.3 mL SWI 30 mg ¹⁷² ; 3.3 mL SWI slowly swirl until completely dissolved; do not shake ¹⁷² allow to settle until bubbles are gone (≥1 min) ¹⁷²	10 mg/mL ¹⁷²	12 h F ^{2,172}	0.3-4 mg/mL NS , D5W, LR ¹⁷² 50 mL* mix by gentle inversion ¹⁷²	16 h F ¹⁷² **(PFL) ¹⁷²	- discard if visible particles are present or solution is discolored ¹⁷² - do not shake ¹⁷²

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Epcoritamab (AbbVie) 4 mg/0.8 mL (F)(PFL) do not shake no preservative ¹⁷³	N/A bring vial to RT prior to use (<1 h) ¹⁷³ gently swirl vial prior to use ¹⁷³ do not invert, vortex, or shake ¹⁷³	5 mg/mL ¹⁷³ For Step-up Dose 1 (0.16 mg) ¹⁷³ To create intermediate vial (0.8 mg/mL): using 4 mg vial: transfer 0.8 mL drug solution into empty vial and add 4.2 mL NS; gently swirl for 30-45 sec	discard unused portion ¹⁷³	SC syringe ¹⁷³ For Step-up Dose 1 (0.16 mg) ¹⁷³ To create dosing vial (0.16 mg/mL): transfer 2.0 mL from intermediate vial into the dosing vial and add 8.0 mL NS; gently swirl for 30-45 sec withdraw 1.0 mL into syringe for administration ¹⁷³ mix gently; do not invert, vortex, or shake ¹⁷³	24 h F, 12 h RT ¹⁷³ (RT storage includes preparation) **(PFL) ¹⁷³	- CAUTION: two concentrations are available - use 4 mg vial for step-up doses only ¹⁷³ - do not use CSTD for volumes less than 1 mL ² ; use filtered venting needle (Chemo- Vent®) for preparation - minimize exposure to daylight ¹⁷³

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Epcoritamab (AbbVie) 4 mg/0.8 mL (F)(PFL) do not shake no preservative ¹⁷³	N/A bring vial to RT prior to use (<1 h) ¹⁷³ gently swirl vial prior to use ¹⁷³ do not invert, vortex, or shake ¹⁷³	5 mg/mL ¹⁷³ For Step-up Dose 2 (0.8 mg) ¹⁷³ To create intermediate vial (0.8 mg/mL): using 4 mg vial: transfer 0.8 mL drug solution into empty vial and add 4.2 mL NS; gently swirl for 30-45 sec	discard unused portion ¹⁷³	SC syringe ¹⁷³ For Step-up Dose 2 (0.8 mg) ¹⁷³ withdraw 1.0 mL from the intermediate vial into syringe for administration mix gently; do not invert, vortex, or shake ¹⁷³	24 h F, 12 h RT ¹⁷³ (RT storage includes preparation) **(PFL) ¹⁷³	- CAUTION: two concentrations are available ¹⁷³ - use 4 mg vial for step-up doses only ¹⁷³ - do not use CSTD for volumes less than 1 mL ² ; use filtered venting needle (Chemo- Vent®) for preparation - minimize exposure to daylight ¹⁷³

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Epcoritamab (AbbVie) 48 mg/0.8 mL (F)(PFL) do not shake no preservative ¹⁷³	N/A bring vial to RT prior to use (<1 h) ¹⁷³ gently swirl vial prior to use ¹⁷³ do not invert, vortex, or shake ¹⁷³	60 mg/mL ¹⁷³	discard unused portion ¹⁷³	SC syringe ¹⁷³ do not invert, vortex, or shake ¹⁷³	24 h F, 12 h RT ¹⁷³ (RT storage includes preparation) **(PFL) ¹⁷³	- CAUTION: two concentrations are available - use 48 mg vial for full doses only ¹⁷³ - do not use CSTD for volumes less than 1 mL ² ; use filtered venting needle (Chemo- Vent®) for preparation - minimize exposure to daylight ¹⁷³
Epirubicin 10 mg/5 mL 20 mg/10 mL 50 mg/25 mL 150 mg/75 mL 200 mg/100 mL (Teva/Novopharm) (F)(PFL) no preservative ¹⁷⁴	N/A	2 mg/mL ¹⁷⁴	8 h F, RT ¹⁷⁴	syringe ¹⁷⁴	48 h F, 24 h RT from initial vial puncture ¹⁷⁴	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Epirubicin 10 mg/5 mL 50 mg/25 mL 200 mg/100 mL (Fresenius Kabi) (F)(PFL) no preservative ¹⁷⁵	N/A record time of puncture	2 mg/mL ¹⁷⁵	8 h ¹⁷⁵	syringe ¹⁷⁵	48 h F , 24 h RT from initial vial puncture ¹⁷⁵	
				100 mL* NS , D5W	48 h F , RT ^{24,175}	
Epirubicin 10 mg/5 mL 50 mg/25 mL 200 mg/100 mL (Pfizer) (F)(PFL) no preservative ¹⁷⁶	N/A record time of puncture	2 mg/mL ¹⁷⁶	8 h ¹⁷⁶	syringe ¹⁷⁶	48 h F , 24 h RT from initial vial puncture ¹⁷⁶	
				100 mL* NS , D5W ⁷⁷	48 h F , RT ¹⁷⁷	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
EPOCHR (LYEPOCHR protocol) (RT) no preservative ^{24,178-181}	see brand specific entries for: DOXOrubicin as applicable	see brand specific entries for: DOXOrubicin, etoposide, vinCRISTine	see brand specific entries for: DOXOrubicin, etoposide, vinCRISTine	etoposide dose ≤125 mg/24 h: in 500 mL NS etoposide dose >125 mg/24 h: in 1000 mL NS	etoposide concentration ≤0.25 mg/mL: complete administration within 72 h RT precipitation occurs at etoposide concentrations >0.25 mg/mL	- final product is a 3-in-1 solution containing etoposide , DOXOrubicin, and vinCRISTine (refer to LYEPOCHR protocol) - use non-DEHP bag and tubing only - administer with 0.2 micron in-line filter
EPOCHR with etoposide phosphate (LYEPOCHR protocol) (RT) no preservative ^{182,183}	see brand specific entries for: DOXOrubicin and etoposide phosphate as applicable	see brand specific entries for: DOXOrubicin, etoposide phosphate, vinCRISTine	see brand specific entries for: DOXOrubicin, etoposide phosphate, vinCRISTine	500 mL NS ¹⁸⁴	4 d RT, 5 d F ^{2,182}	- final product is a 3-in-1 solution containing etoposide phosphate , DOXOrubicin, and vinCRISTine (refer to LYEPOCHR protocol)

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
eriBULin 1 mg/2 mL (Eisai Limited) (RT)(PFL) ¹⁸⁵ preservative ²⁴	N/A	0.5 mg/mL ¹⁸⁵	discard unused portion ^{24,185}	IV syringe ¹⁸⁵	24 h F, 6 h RT ¹⁸⁵	- do not administer through dextrose containing lines ¹⁸⁵ - vials contain dehydrated alcohol USP (5% v/v) ¹⁸⁵
eriBULin 1 mg/2 mL (Natco) (RT)(PFL) ¹⁸⁶ preservative ¹⁸⁶	N/A	0.5 mg/mL ¹⁸⁶	discard unused portion ¹⁸⁶	IV syringe ¹⁸⁶	24 h F, 6 h RT ¹⁸⁶	- do not dilute or administer with dextrose containing solutions ¹⁸⁶ - vials contain dehydrated alcohol USP (5% v/v) ¹⁸⁶

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Etoposide 100 mg/5 mL 200 mg/10 mL 500 mg/25 mL 1000 mg/50 mL (Teva) (RT)(PFL) no preservative ¹⁸⁷	N/A	20 mg/mL ¹⁸⁷	discard unused portion ¹⁸⁷	0.2-0.4 mg/mL NS ¹⁸⁷ 100-1000 mL†	stability is concentration dependent 0.2-0.3 mg/mL: 7 d F, ¹⁸⁸ 2 d RT ^{188,189} 0.4-0.5 mg/mL: 1 d F, ¹⁸⁸ 1d RT ¹⁸⁸ 0.6-9.0 mg/mL: generally unstable 9.5 mg/mL: 2 d F, ¹⁸⁸ 1d RT ¹⁸⁸ 10-12 mg/mL: 7 d F, ¹⁸⁸ 2 d RT ^{188,189}	- use non-DEHP bag and tubing only - administer with 0.2 micron in-line filter ¹⁹⁰ - for LYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing either etoposide or etoposide phosphate AND DOXOrubicin and vinCRISTine) - for ULY0 D-PACE protocol, see entry for DPACE (3-in-1 solution containing etoposide, CISplatin, cyclophosphamide)
				D5W ¹⁸⁷	4 h RT ^{187,191}	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Etoposide phosphate (ETOPOPHOS®) 100 mg (Xediton/Cheplapharm) (F)(PFL) no preservative ¹⁹²⁻¹⁹⁴ (SAP)	5 mL NS , D5W, SWI, BWI ¹⁹⁵	20 mg/mL ¹⁹⁵	in NS , D5W, SWI: 12 h F, RT ^{2,195} in BWI: 7 d F, 48 h RT ¹⁹⁵	500 mL NS , D5W ¹⁹⁵ (do not dilute to less than 0.1 mg/mL) ¹⁹⁵	24 h F, RT ¹⁹⁵	- for LYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing either etoposide or etoposide phosphate AND DOXOrubicin and vinCRISTine)
	10 mL NS , D5W, SWI, BWI ¹⁹⁵	10 mg/mL ¹⁹⁵				
Filgrastim (NEUPOGEN®) 300 mcg/1 mL 480 mcg/1.6 mL (Amgen) (F)(PFL) do not shake no preservative ¹⁹⁶	N/A	300 mcg/mL ¹⁹⁶	discard unused portion ¹⁹⁶	SC syringe ¹⁹⁶	10 d F ^{2,197}	- albumin is added to D5W to prevent filgrastim adsorption to plastic ¹⁹⁶ - incompatible with saline ^{196,198} - do NOT dilute to concentration less than 5 mcg/mL ¹⁹⁶
				50-100 mL D5W only ¹⁹⁸ in PVC, polyolefin, or glass ¹⁹⁶ (for filgrastim concentrations of 5-15 mcg/mL in D5W, add albumin 2 mg/mL) ¹⁹⁶	7 d F ¹⁹⁷	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Filgrastim (NIVESTYM®) 300 mcg/1 mL 480 mcg/1.6 mL (Pfizer) (F)(PFL) do not shake no preservative ¹⁹⁹	N/A	300 mcg/mL ¹⁹⁹	discard unused portion ¹⁹⁹	SC syringe	10 d F, 24 h RT ^{2,200}	- albumin is added to D5W to prevent filgrastim adsorption to plastic ¹⁹⁹ - incompatible with saline ¹⁹⁹ - do NOT dilute to concentration less than 5 mcg/mL ¹⁹⁹
				50-100 mL D5W only ¹⁹⁸ in PVC, polyolefin, or glass ¹⁹⁹ (for filgrastim concentrations of 5-15 mcg/mL in D5W, add albumin 2 mg/mL) ¹⁹⁹	complete administration within 24 h RT ²⁰¹	
Fludarabine 50 mg (Accord) (F) no preservative ²⁰²	N/A	25 mg/mL ²⁰²	discard unused portion ²⁰²	dilute to maximum of 1 mg/mL NS, D5W ²⁰² 100 mL†	72 h F, 24 h RT ²⁰²	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Fludarabine 50 mg (Teva) (F) no preservative ²⁰³	N/A	25 mg/mL ²⁰³	discard unused portion ²⁰³	dilute to maximum of 1 mg/mL NS, D5W ²⁰³ 100 mL†	72 h F, 24 h RT ²⁰³	
Fluorouracil 5000 mg/100 mL (Accord) (RT)(PFL) no preservative ²⁰⁴	N/A	50 mg/mL ²⁰⁴	12 h RT ^{2,205}	syringe ²⁰⁴	4 d RT ²⁰⁵	
				0.5-10 mg/mL D5W ²⁰⁵ 500 mL†	4 d RT ²⁰⁵	
				CIVI: ambulatory pump ²⁰⁶	complete within 8 d ²⁰⁵	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Fluorouracil 500 mg/10 mL 5000 mg/100 mL (Sandoz) (RT)(PFL) no preservative ²⁰⁷	N/A	50 mg/mL ²⁰⁷	12 h RT ^{2,208}	syringe	4 d RT ^{2,208}	
				0.35-15 mg/mL D5W ²⁰⁸ 500 mL†	10 d F, 4 d RT ^{2,208}	
				CIVI: ambulatory pump ²⁰⁶	complete within 8 d ²⁰⁹⁻²¹¹	
Gemcitabine 1000 mg 2000 mg (Accord) (RT) no preservative ²¹²	1000 mg: 25 mL NS ²¹² 2000 mg: 50 mL NS ²¹²	38 mg/mL ²¹²	12 h RT ^{2,212} refrigeration may cause crystallization ²¹²	syringe ²¹²	24 h RT ^{2,212}	
				0.1-38 mg/mL NS ²¹² 50-250 mL†	4 d RT ^{2,213,214}	
Gemcitabine intravesical 1000 mg 2000 mg (Accord) (RT) no preservative ²¹²	1000 mg: 25 mL NS ²¹² 2000 mg: 50 mL NS ²¹²	38 mg/mL ²¹²	12 h RT ^{2,212} refrigeration may cause crystallization ²¹²	syringe dilute with NS to final volume of 45-90 mL ^{144,145,215-217}	up to 38 mg/mL: 24 h RT ^{2,212}	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Gemcitabine 200 mg/5.3 mL 1000 mg/26.3 mL 2000 mg/52.6 mL (Pfizer/Hospira) (F) no preservative ²¹⁷	N/A	38 mg/mL ²¹⁷	discard unused portion ²¹⁷	syringe ²¹⁷	0.1-26 mg/mL: 10 d F, 24 h RT **(PFL) ^{2,218,219}	
				0.1-38 mg/mL NS, D5W ²¹⁷ 50-250 mL†	27-38 mg/mL: 24 h RT ²¹⁹	
Gemcitabine intravesical 200 mg/5.3 mL 1000 mg/26.3 mL 2000 mg/52.6 mL (Pfizer/Hospira) (F) no preservative ²¹⁷	N/A	38 mg/mL ²¹⁷	discard unused portion ²¹⁷	syringe dilute with NS to final volume of 45-90 mL ^{144,145,215-217}	0.1-26 mg/mL: 10 d F, 24 h RT **(PFL) ^{2,218,219} 27-38 mg/mL: 24 h RT ²¹⁹	
Gemcitabine (NOTE: concentration) 200 mg/5 mL 1000 mg/25 mL 2000 mg/50 mL (Sandoz) (F) no preservative ²²⁰	N/A	40 mg/mL ²²⁰	discard unused portion ²²⁰	syringe ²²⁰	1-25 mg/mL: 10 d F, 4 d RT ^{2,220,221}	CAUTION: alternative concentration
				0.1–40 mg/mL NS, D5W ²²⁰ 50-250 mL†	26-40 mg/mL: 24 h RT ²²⁰	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Gemcitabine (NOTE: concentration) intravesical 200 mg/5 mL 1000 mg/25 mL 2000 mg/50 mL (Sandoz) (F) no preservative ²²⁰	N/A	40 mg/mL ²²⁰	discard unused portion ²²⁰	syringe dilute with NS to final volume of 45-90 mL ^{144,145,215-217}	1-25 mg/mL: 10 d F, 4 d RT ^{2,220,221} 26-40 mg/mL: 24 h RT ²²⁰	CAUTION: alternative concentration
Gemtuzumab ozogamicin 4.5 mg (Pfizer) (F)(PFL) no preservative ²²²	5 mL SWI ²²² allow vial to come to RT prior to use (~5 min) ²²² swirl gently to mix; do NOT shake ²²²	1 mg/mL ²²²	6 h F, 3 h RT ²²² protect from light if not used immediately ²²²	0.075-0.234 mg/mL NS ²²² 25-50 mL† mix by gentle inversion; do NOT shake ²²²	complete administration within 12 h F, 6 h RT ²²² (PFL)** if refrigerated, bring bag to RT over 1 h prior to administration ²²²	- administer with 0.2 micron in-line filter ²²² - protect infusion bag from light (including UV) for administration ²²² - protect administration line from light ONLY if hang time will be longer than 2 h ^{222,223} - solution may contain white particulates which do not affect product quality ²²²

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Glofitamab 2.5 mg/2.5 mL (Roche) (F)(PFL) no preservative ²²⁴	N/A	1 mg/mL ²²⁴	discard unused portion ²²⁴	0.1-0.6 mg/mL NS, ½NS ²²⁴ 25 mL† ^{224,225} For Step-up Dose 1 (2.5 mg) withdraw 7 mL from infusion bag prior to adding drug volume ^{224,225} gently invert to mix do NOT shake ²²⁴	64 h F, plus an additional 12 h RT including infusion time ²²⁴ if refrigerated, bring bag to RT prior to administration (max 4 h RT prior to infusion) ²²⁴	- use 2.5 mg vials for 2.5 mg dose only ²²⁴ - do not use if contains visible particulates or is cloudy or discoloured ²²⁴ - in-line filter is not required, but may be used ²²⁴ (e.g., 0.2 micron ²²⁶)
Glofitamab 10 mg/10 mL (Roche) (F)(PFL) no preservative ²²⁴	N/A	1 mg/mL ²²⁴	discard unused portion ²²⁴	0.1-0.6 mg/mL NS, ½NS ²²⁴ 50-100 mL† gently invert to mix do NOT shake ²²⁴	64 h F, plus an additional 12 h RT including infusion time ²²⁴ if refrigerated, bring bag to RT prior to administration (max 4 h RT prior to infusion) ²²⁴	- use 10 mg vials for 10 mg and 30 mg doses only ²²⁴ - do not use if contains visible particulates or is cloudy or discoloured ²²⁴ - in-line filter is not required, but may be used ²²⁴ (e.g., 0.2 micron ²²⁶)

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
IDarubicin PFS 5 mg/5 mL 10 mg/10 mL 20 mg/20 mL (Pfizer) (F)(PFL) no preservative ²²⁷	N/A	1 mg/mL ²²⁷	discard unused portion ²²⁷ **(PFL) ²²⁷	syringe ²²⁷	use within 4 h from initial puncture ^{227,228}	- avoid alkaline solutions ²²⁷
Ifosfamide 1000 mg 3000 mg (Baxter) (RT) no preservative ²²⁹	1000 mg: 20 mL SWI ²²⁹ 3000 mg: 60 mL SWI ²²⁹ shake well	50 mg/mL ²²⁹	12 h F, RT ^{2,230}	0.6-20 mg/mL NS , D5W, LR ²²⁹ 500 mL†	72 h F, 24 h RT ²³⁰ 24 h F, RT when mixed with mesna ⁷⁷	
Ifosfamide 1000 mg 3000 mg (Fresenius Kabi) (RT) no preservative ²³¹	1000 mg: 20 mL SWI ²³¹ 3000 mg: 60 mL SWI ²³¹ shake well	50 mg/mL ²³¹	12 h F, RT ^{2,232}	0.6-20 mg/mL NS , D5W, LR ²³¹ 500 mL†	72 h F, 24 h RT ²³² 24 h F, RT when mixed with mesna ⁷⁷	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Iniparib 100 mg/10 mL (sanofi-aventis) (F) no preservative ²³³ (SAP)	N/A	10 mg/mL ²³³	discard unused portion ²³³	250 mL NS, D5W dilute to 250 mL final volume by withdrawing volume from bag equal to volume of drug to be added ²³³ (OR undiluted in empty infusion bag and qs to final volume of 250 mL with NS, D5W ²³³)	24 h RT ²³³	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Inotuzumab ozogamicin 0.9 mg (Pfizer) (F)(PFL) no preservative ²³⁴	4 mL SWI ²³⁴ gently swirl vial to mix ²³⁴	0.25 mg/mL ²³⁴ record time of reconstitution	4 h F ²³⁴ dilute dose within 4 h of reconstitution ²³⁴ protect from light if not used immediately ²³⁵	0.01-0.1 mg/mL NS ²³⁴ 25-50 mL† mix by gentle inversion ²³⁴	complete administration within 8 h of reconstitution F, RT ²³⁴ (PFL) ²³⁴ if refrigerated, bring bag to RT over 1 h prior to administration ²³⁴	- do NOT shake ²³⁴ - protect container from UV and fluorescent light during storage and administration ^{234,235} - protect administration line from light ONLY if hang time will be longer than 1 h ^{234,235}
Ipilimumab 50 mg/10 mL 200 mg/40 mL (BMS Canada) (F)(PFL) no preservative ²³⁶	N/A	5 mg/mL ²³⁶	12 h F, RT ^{2,237}	1-4 mg/mL NS, D5W ²³⁶ 25-250 mL† OR undiluted in empty infusion bag or glass bottle (allow vials to stand at RT for ~5 min prior to withdrawal of contents) ²³⁶	24 h F, RT ²³⁷	- do NOT shake ²³⁶ - administer with 0.2 micron in-line filter ²³⁶ - vials may contain translucent-to- white amorphous particles ²³⁶ - discard if cloudy or has pronounced colour change (should be clear to pale yellow) ²³⁶

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Irinotecan 40 mg/2 mL 100 mg/5 mL 500 mg/25 mL (Accord) (RT)(PFL) no preservative ²³⁸	N/A	20 mg/mL ²³⁸	discard unused portion ²³⁸	0.12-3.0 mg/mL D5W (preferred), NS ²³⁸ 250-500 mL†	48 h F, 24 h RT **(PFL) ²³⁸	
Irinotecan 40 mg/2 mL 100 mg/5 mL 300 mg/15 mL 500 mg/25 mL (Auro) (RT)(PFL) no preservative ²³⁹	N/A	20 mg/mL ²³⁹	discard unused portion ²³⁹	0.12-3.0 mg/mL D5W (preferred), NS ²³⁹ 250-500 mL†	10 d F, 4 d RT ^{2,239} **(PFL) ²³⁹ if NOT protected from light: 72 h RT ²³⁹	
Irinotecan 40 mg/2 mL 100 mg/5 mL 300 mg/15 mL 500 mg/25 mL (Eugia) (RT)(PFL) no preservative ²⁴⁰	N/A	20 mg/mL ²⁴⁰	discard unused portion ²⁴⁰	0.12-3.0 mg/mL D5W (preferred), NS ²⁴⁰ 250-500 mL†	10 d F, 4 d RT ^{2,240} **(PFL) ²⁴⁰ if NOT protected from light: 72 h RT ²⁴⁰	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART						
DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Irinotecan 40 mg/2 mL 100 mg/5 mL 300 mg/15 mL 500 mg/25 mL (GMP) (RT)(PFL) no preservative ²⁴¹	N/A	20 mg/mL ²⁴¹	discard unused portion ²⁴¹	0.12-3.0 mg/mL D5W (preferred), NS ²⁴¹ 250-500 mL†	10 d F, 4 d RT ²⁴¹ **(PFL) ²⁴¹ if NOT protected from light: 72 h RT ²⁴¹	- due to bung curvature, filtered venting needle (e.g., Chemo- Vent®) must be used in place of CSTD for compounding ²⁴²
Irinotecan 40 mg/2 mL 100 mg/5 mL 300 mg/15 mL 500 mg/25 mL (Pfizer/Hospira) (RT)(PFL) no preservative ²⁴³	N/A	20 mg/mL ²⁴³	discard unused portion ²⁴³	0.12-3.0 mg/mL D5W (preferred), NS ²⁴³ 250-500 mL†	10 d F, 4 d RT ^{2,243} **(PFL) ²⁴³ if NOT protected from light: 72 h RT ²⁴³	
Irinotecan liposomal 43 mg/10 mL (Ipsen) (F)(PFL) no preservative ²⁴⁴	N/A	4.3 mg/mL ²⁴⁴	discard unused portion ²⁴⁴	500 mL NS, D5W ²⁴⁴ mix by gentle inversion ²⁴⁴	24 h F, 4 h RT ²⁴⁴ **(PFL) if refrigerated, bring bag to RT prior to administration ²⁴⁴	- if a CSTD is not used during compounding, use a 21 gauge (or lower gauge) needle to withdraw drug from vial ²⁴⁴ - do not use in-line filter ²⁴⁴

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Isatuximab 100 mg/5 mL 500 mg/25 mL (sanofi-aventis) (F)(PFL) do not shake no preservative ²⁴⁵	N/A	20 mg/mL ²⁴⁵ inspect vial and discard if discolouration or visible particles are present ²⁴⁵	discard unused portion ²⁴⁵	250 mL NS, D5W ²⁴⁵ mix by gentle inversion; do NOT shake ²⁴⁵	48 h F plus an additional 8 h RT including infusion time ²⁴⁵	- administer with a 0.2 micron in-line filter ²⁴⁵
Ixabepilone 15 mg (contains 16 mg) 45 mg (contains 47 mg) (BMS) (F)(PFL) no preservative ²⁴⁶ (SAP)	15 mg: 8 mL diluent (supplied) ²⁴⁶ 45 mg: 23.5 mL diluent (supplied) ²⁴⁶	2 mg/mL ²⁴⁶	1 h RT ²⁴⁶	0.2-0.6 mg/mL LR ²⁴⁶	6 h RT ²⁴⁶	- use non-DEHP bag and administration set ²⁴⁶ - administer with 0.2 micron in-line filter ²⁴⁶

* Suggested volume based on usual dose range and any concentration range of stability data

† see [BC Cancer IV Bag Selection table](#); standardized bag sizes are provided for select Benefit Drugs with concentration-dependent stability or large drug volume

** Protect from light means minimizing exposure to direct sunlight over a *storage* period. More specific information on protection from light (eg, protecting container and tubing *during administration*) will be indicated in the Special Precautions/Notes column.

*** Contains DMA (N,N dimethylacetamide). Product may be incompatible with closed system transfer devices (CSTD) such as ChemoLock.

Centres are not to change content locally. All suggestions for change are to be forwarded to the Cancer Drug Manual editor.

Explanatory Notes:

Stability data assumes products prepared using standard aseptic technique in biological safety cabinet at low risk for contamination according to the classification outlined in USP 797.^{247,248}

Vial stability: Stability of solution after first puncture or reconstituted solution.

Storage temperature: If information states same stability with refrigerator and room temperature storage, then fridge stability is bolded as preferred (ie, to minimize growth of micro-organisms).

Discard unused portion: Unused portion from single use vials should be discarded at the end of the day.

“overfill known” is stated if the manufacturer states overfill that is present is within acceptable limits.

“Complete administration within ___” is stated if the manufacturer specifies that the infusion must be completed in a specific time frame following preparation, usually including entire time required for preparation (from first puncture), storage, and administration of infusion.

Nomenclature for **In-line filters** has been standardized in the chart to 0.2 micron filter size. For more information, refer to CDM drug monograph.

Abbreviations:

BWI = bacteriostatic water for injection

CIVI: ambulatory pump = Continuous Intravenous Infusion (e.g., elastomeric infusor)

CSTD = closed system transfer device

D5W = dextrose 5% in water

DMA = N,N dimethylacetamide

F = refrigerate

LR = lactated ringer's solution

non-DEHP = not containing Di(2-ethylhexyl) phthalate (DEHP)

non-PVC = not containing polyvinyl chloride (PVC)

NS = normal saline

PES = polyethersulfone

PFL = protect from light

RT = room temperature

SAP = drug is approved for use through the Health Canada Special Access Program (SAP)

SWI = sterile water for injection

References:

1. SteriMax Inc. PROLEUKIN® product monograph. Oakville, Ontario; July 6, 2020.
2. BC Cancer. Provincial Pharmacy Directive Number II-20: Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer; May 1, 2022.
3. McEvoy GK, editor. AHFS 2008 Drug Information. Bethesda, Maryland: American Society of Health-System Pharmacists, Inc.; p. 917–925
4. Koreth J, Matsuoka K, Kim HT, et al. Interleukin-2 and regulatory T cells in graft-versus-host disease. *N Engl J Med* ; 2011;365(22):2055–2066
5. Koreth J, Alyea EP, Cutler C, Ho VT, et al. Clinical Study Protocol: A phase I study of ultra-low dose subcutaneous interleukin-2 (IL-2) for treatment of refractory chronic graft versus host disease. Boston, MA, USA: Dana Farber Cancer Institute; Harvard Medical Centre; 14 Dec, 2010.
6. Temple-Oberle CF, Byers BA, Hurdle V, et al. Intra-lesional interleukin-2 for in-transit melanoma. *J Surg Oncol* ; 2014;109(4):327–331
7. BC Cancer Skin and Melanoma Tumour Group. (SMILALD) BC Cancer Protocol Summary for Treatment of In-Transit Melanoma Using Intralesional Aldesleukin (IL-2). Vancouver, British Columbia: BC Cancer; 1 August, 2021.
8. SteriMax Inc. Personal Communication: Importation of US-Labelled PROLEUKIN® for Injection (by Iovance) due to the current shortage of Canadian-Authorized PROLEUKIN®. January 10, 2025.
9. Rui Paiva. Business Unit Director, Transplant and Oncology. Personal communication. 1 June, 2009.
10. Bayer HealthCare Pharmaceuticals. MabCampath® Package Insert. Toronto, Ontario; 1 September, 2007.
11. Lundin J, Porwit-MacDonald A, Rossmann ED, et al. Cellular immune reconstitution after subcutaneous alemtuzumab (anti-CD52 monoclonal antibody, CAMPATH-1H) treatment as first-line therapy for B-cell chronic lymphocytic leukaemia. *Leukemia* ; 22 January, 2004(18):484–490
12. Berlex Canada Inc. Campath Drug Information. San Antonio, Texas; undated,
13. Neijssen J, Cardoso RMF, Chevalier KM, et al. Discovery of amivantamab (JNJ-61186372), a bispecific antibody targeting EGFR and MET. *J Biol Chem* ; January, 2021;296(100641):1–13DOI: <https://doi.org/10.1016/j.jbc.2021.100641>
14. Eric Li, Manager Special Access Program, Janssen Inc. Manager Special Access Program, Janssen Inc. Personal Communication. November 18, 2021.
15. Janssen Biotech Inc. RYBREVANT® full prescribing information. Horsham, PA, USA; May, 2021.
16. Janssen Inc. RYBREVANT® product monograph. Toronto, Ontario; March 30, 2022.
17. Erfa Canada Inc. AMSA PD® injection product monograph. Westmount, Quebec; 16 August, 2005.
18. Phebra Pty Inc (distributed in Canada by ICON Plc). Arsenic trioxide product monograph. Dundas, Ontario; November 7, 2019.
19. Sandoz Canada Inc. Arsenic trioxide for injection product monograph. Boucherville, Quebec; March 8, 2021.
20. SteriMax Inc. Arsenic Trioxide product monograph. Oakville, Ontario; September 27, 2019.
21. Hoffmann-La Roche Limited. TECENTRIQ® product monograph. Mississauga, Ontario; September 19, 2019.
22. Hoffmann-La Roche Limited. TECENTRIQ® SC product monograph. Mississauga, Ontario; March 13, 2024.
23. EMD Serono. BAVENCIO® product monograph. Mississauga, Ontario; 4 May, 2018.
24. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 19 September, 2007.
25. Celgene Inc. VIDAZA® product monograph. Mississauga, Ontario; 22 March, 2018.
26. Tutino A, Lai M. Cold water reconstitution of Vidaza with subsequent refrigerated storage prolongs drug stability. *Eur J Oncol Pharm* ; 2011;5(3-4):24–25, 34
27. Celgene Corporation. VIDAZA® full prescribing information. Summit, New Jersey, USA; May, 2022.
28. Dr. Reddy's Laboratories Limited. Azacitidine for injection product monograph. Mississauga, Ontario; 19 April, 2017.
29. Hikma Canada Inc. Azacitidine for injection product monograph. Caledon, Ontario; November 6, 2020.
30. Merck Canada Inc. OncoTICE® product monograph. Kirkland, Quebec; 29 April, 2019.
31. BC Cancer. Provincial Pharmacy Directive Number II-20: Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer; December 5, 2018.
32. Verity Pharmaceuticals Inc. VERITY-BCG® Product Monograph. Mississauga, Ontario; 5 January, 2021.
33. GlaxoSmithKline. Belantamab mafodotin (GSK2857916) Investigational Product Information for Compassionate Use - Version 5.0. Collegeville, Pennsylvania, USA; 25 September, 2019.
34. GlaxoSmithKline. Belantamab mafodotin (GSK2857916) LYOPHILIZED POWDER Investigational Product Information for Compassionate Use - Version 1. Collegeville, Pennsylvania, USA; 26 October, 2020.
35. GlaxoSmithKline. BLENREP® full prescribing information. Research Triangle Park, North Carolina, USA; August, 2020.

36. Spectrum Pharmaceuticals Inc. BELEODAQ® full prescribing information. Irvine, CA, USA; April, 2012.
37. Natco Pharma Canada Inc. NAT-Bendamustine product monograph. Mississauga, Ontario; March 4, 2020.
38. Eugia Pharma Inc. Bendamustine hydrochloride product monograph. Woodbridge, Ontario; October 4, 2023.
39. Teva Canada Limited. TREANDA® product monograph. Toronto, Ontario; January 10, 2018.
40. Lundbeck Canada Inc. TREANDA® product monograph. Montreal, Quebec; August 22, 2012.
41. Hoffman-La Roche Limited. AVASTIN® product monograph. Mississauga, Ontario; 6 June, 2018.
42. Organon Canada Inc. (for Samsung Bioepis Co. Ltd.). AYBINTIO® product monograph. Kirkland, Quebec; June 27, 2024.
43. Amgen Canada Inc. MVASI® product monograph. Mississauga, Ontario; 5 June, 2019.
44. Pfizer Canada-ULC. ZIRABEV® product monograph. Kirkland, Quebec; June 17, 2021.
45. Fresenius Kabi Canada Ltd. Bleomycin for injection product monograph. Richmond Hill, Ontario; 1 June , 2016.
46. Pfizer Canada Inc. Bleomycin for Injection product monograph. Kirkland, Quebec; 8 August, 2017.
47. Amgen Canada Inc. BLINCYTO® product monograph. Mississauga, Ontario; 12 July , 2016.
48. Amgen Canada Inc. BLINCYTO® product monograph. Mississauga, Ontario; May 6, 2021.
49. Actavis Pharma Company. ACT BORTEZOMIB® Bortezomib for injection product monograph. Mississauga, Ontario; 24 September, 2015.
50. Law S, Charbonneau LF, Iazzeta J, et al. Stability of generic formulations of bortezomib 1.0 and 2.5 mg/mL in vials and syringes stored at 4°C and room temperature (23°C). CJHP Canadian Society of Hospital Pharmacists Professional Practice Conference 2018 Poster Abstracts. Feb 4 - 6, 2018url
51. Apotex Inc. Bortezomib for injection product monograph. Toronto, Ontario; 15 February, 2019.
52. Walker SE, Law S, Ma N. (Abstract) Stability of 1.0 and 2.5 mg/mL bortezomib solution in vials and syringes following reconstitution with 0.9% sodium chloride at 4°C and room temperature (25°C). (Apotex brand). Department of Pharmacy, Sunnybrook Health Sciences Centre and Leslie Dan Faculty of Pharmacy, University of Toronto. Toronto, ON ; 2019
53. Janssen Inc. VELCADE® product monograph. Toronto, Ontario; 20 June, 2013.
54. Juno Pharmaceuticals Corp. Bortezomib for injection product monograph. Mississauga, Ontario; 23 Jun, 2020.
55. Law S, Charbonneau F, Iazzeta J, et al. Stability of generic formulations of bortezomib 1.0 and 2.5 mg/mL in vials and syringes stored at 4°C and room temperature (23°C or 25°C). CJHP ; 2021;74(1):57–69
56. Marcan Pharmaceuticals Inc. Bortezomib product monograph. Ottawa, Ontario; October 17, 2019.
57. MSN Laboratories Private Limited, Formulations Division - Nandigama. Reconstitution Study Report for Bortezomib for Injection 3.5 mg vial (Reference Protocol No SSP-CL011-003-00) . May 2, 2016:1–8
58. Suresh Arvapalli. Director - Technical. Marcan Pharmaceuticals Inc. Personal communication - extended stability study on Bortezomib for Injection 3.5 mg. August 9, 2021.
59. Pharmascience Inc. pms-BORTEZOMIB product monograph. Montréal, Quebec; March 8, 2019.
60. Taro Pharmaceuticals Inc. Taro-Bortezomib product monograph. Brampton, Ontario; October 27, 2021.
61. Teva Canada Limited. Bortezomib for injection® product monograph. Toronto, Ontario; 22 January, 2015.
62. Seagen Canada Inc. ADCETRIS® product monograph. Mississauga, Ontario; June 11, 2021.
63. Pharmascience Inc. Busulfan for injection product monograph. Montreal, Quebec; 14 June, 2018.
64. SteriMax Inc. Busulfan for injection product monograph. Oakville, Ontario; 4 May, 2017.
65. Dr. Reddy's Laboratories Canada Inc. Cabazitaxel for injection product monograph. Mississauga, Ontario; January 14, 2020.
66. Formative Pharma Inc. Cabazitaxel for injection product monograph. Burlington, Ontario; December 17, 2024.
67. Sandoz Canada Inc. Cabazitaxel for injection product monograph. Boucherville, Quebec; 17 December , 2019.
68. sanofi-aventis Canada Inc. JEV TANA® product monograph. Laval, Quebec; 7 September , 2017.
69. Servier Canada Inc. ASPARLAS® product monograph. Laval, Quebec; March 8, 2024.
70. Louis Turcotte, Associate, Medical Affairs. Servier Canada. Personal communication: ASPARLAS® - protection from light. August 8, 2025.
71. Accord Healthcare Inc. Carboplatin injection® product monograph. Kirkland, Quebec; May 15, 2019.
72. Omega Laboratories Ltd. Carboplatin injection product monograph. Montreal, Quebec; 24 March , 2011.
73. Nayla El Zir. Associate, Regulatory Affairs, Omega Laboratories Limited. Personal communication. 12 April, 2017.
74. Pfizer Canada-ULC. Carboplatin injection product monograph. Kirkland, Quebec; 31 December, 2018.
75. Novopharm Limited. Carboplatin Package Insert. Toronto, Canada; Undated,
76. Manjinder S Kang. Regulatory Affairs Drug Information Pharmacist, Novopharm Canada. Personal communication. 14 March, 2005.
77. Trissel LA. Handbook on Injectable Drugs. 13th ed. Bethesda, MD: American Society of Health-System Pharmacists, Inc.; 2005.

78. Repchinsky C editor. Paraplatin-AQ, Compendium of Pharmaceuticals and Specialties. 12th ed. Ottawa, Ontario: Canadian Pharmacists Association; 2004.
79. BC Cancer Provincial Pharmacy Professional Practice Council. BC Cancer Provincial Pharmacy Professional Practice Council meeting minutes . April 13, 2022.
80. Amgen Canada Inc. KYPROLIS® product monograph. Mississauga, Ontario; January 27, 2021.
81. Luis Simao. Area Manager, ICU Medical Canada. Personal communication. 11 May, 2018.
82. Amgen Canada Inc. KYPROLIS® product monograph. Mississauga, Ontario; 6 July, 2017.
83. Diane Lord. Medical Information, Amgen Canada Inc. Personal communication. 8 May, 2018.
84. Marcan Pharmaceuticals Inc. BiCNU® product monograph. Ottawa, Ontario; February 5, 2025.
85. Sindhoor Talasila. Assistant Manager - RA. Marcan Pharmaceuticals Inc. Personal communication - Carmustine for injection. May 20, 2025.
86. Emcure Pharmaceuticals Limited, Research & Development Department. Report for Stability of Admixture Solution: Carmustine Injection 100 mg/mL (1 mL) at Final Concentration of 3.06 mg/mL. Hinjawadi, Pune, India; April, 2025.
87. Pritesh Sen. Pharmacovigilance, Drug Safety & Medical Affairs Specialist. Marcan Pharmaceuticals Inc. Personal communication - Carmustine for injection. April 17, 2024.
88. Emcure Pharmaceuticals Limited, Research & Development Department. Admixture Study Report: Admixture Study of Carmustine for Injection 100 mg/Vial Using Propylene Glycol for Reconstitution (Dose 800 mg/m²). Hinjawadi, Pune, India; August, 2019.
89. SteriMax Inc. Carmustine for Injection product monograph. Oakville, Ontario; October 11, 2022.
90. Regeneron Canada Company. LIBTAYO® product monograph. Mississauga, Ontario; January 30, 2025.
91. sanofi-aventis Canada Inc. LIBTAYO® product monograph. Laval, Quebec; 10 April, 2019.
92. ImClone LLC (distributed by Eli Lilly Canada Inc). ERBITUX® product monograph. Toronto, Ontario; January 10, 2018.
93. Accord Healthcare Inc. Cisplatin injection product monograph. Kirkland, Quebec; February 15, 2019.
94. Pfizer Canada-ULC. Cisplatin injection product monograph. Kirkland, Quebec; December 7, 2018.
95. Sandoz Canada Inc. Cisplatin Injection BP product monograph. Boucherville, Quebec; 13 April , 2011.
96. Sandoz Canada Inc. Cisplatin Injection BP product monograph. Boucherville, Quebec; September 28, 2015.
97. Teva Canada Limited. Cisplatin injection product monograph. Toronto, Ontario; March 6, 2013.
98. Fresenius Kabi Canada Ltd. Cladribine injection® product monograph. Richmond Hill, Ontario; 9 June, 2015.
99. BC Cancer Lymphoma Tumour Group. (LYCDA) BC Cancer Protocol Summary for Treatment of Hairy Cell Leukemia with Cladribine. Vancouver, British Columbia: BC Cancer; 1 January , 2021.
100. Myrna O'Brodovich. Senior Medical Information Associate, Ortho Biotech. Personal communication. 2 April, 2008.
101. de Lemos ML, Hamata L. Stability issues of parenteral chemotherapy drugs. J Oncol Pharm Pract ; Mar, 2007;13(1):27–31
102. Generic Medical Partners Inc. Cladribine injection product monograph. Toronto, Ontario; December 3, 2019.
103. Jazz Pharmaceuticals Canada Inc. RYLAZE® product monograph. Mississauga, Ontario; September 2, 2022.
104. Baxter Corporation. PROCYTOX® product monograph. Mississauga, Ontario; 7 September, 2012.
105. Baxter Corporation Medical Information. Personal communication. 14 May, 2020.
106. Baxter Healthcare Corporation. Cyclophosphamide for injection full prescribing information. Deerfield, Illinois USA; May, 2013.
107. Trissel's® 2 Clinical Pharmaceutics Database - Lexicomp Online (database on the Internet). Cyclophosphamide. Wolters Kluwer Clinical Drug Information Inc.; Accessed 14 May, 2020. Available at: <http://online.lexi.com>
108. SteriMax Inc. Cyclophosphamide for injection product monograph. Ontario, Canada; May 14, 2024.
109. Zuber Patel. Associate Director - Quality Operations. SteriMax Medical Information. Personal communication: cyclophosphamide final product stability. July 31, 2025.
110. Pfizer Canada Inc. Cytarabine Solution for Injection product monograph. Kirkland, Quebec; 3 November, 2015.
111. BC Cancer Lymphoma Tumour Group. (LYIT) BC Cancer Protocol Summary for Treatment of Lymphoma using Intrathecal Methotrexate and Cytarabine. Vancouver, British Columbia: BC Cancer; 1 June , 2014.
112. BC Cancer Miscellaneous Origin Tumour Group. (MOIT) BC Cancer Protocol Summary for Solid Tumours using Intrathecal Methotrexate and/or Thiotepa and/or Cytarabine. Vancouver, British Columbia: BC Cancer; 1 October , 2018.
113. BC Cancer. Systemic Therapy Policy and Procedure III-50: Administration of High Alert Medications by the Intrathecal Route via Lumbar Puncture or Ommaya Reservoir. Vancouver, British Columbia; May 1, 2019.
114. Hematology/Oncology Pharmacy Association. HOPA News Clinical Pearls: Intrathecal Chemotherapy: Focus on Drugs, Dosing, and Preparation. 13(4) ed. Chicago, Illinois, USA: Hematology/Oncology Pharmacy Association; 2016.

115. BC Cancer Agency - Provincial Pharmacy Professional Practice Council. Minutes of The Provincial Pharmacy Professional Practice Council (P4C) Meeting. Vancouver, British Columbia: BC Cancer; Feb 5 , 2014.
116. Astier A, Pinguet F, Vigneron J. The practical stability of anticancer drugs: SFPO and ESOP recommendations. Eur J Oncol Pharm ; 2010;4:4–10
117. Dellamorte Bing C, Nowobilski-Vasilios A. Extended Stability for Parenteral Drugs. 5th ed. Bethesda, Maryland: American Society of Health-System Pharmacists, Inc; 2013.
118. Pharmascience Inc. Cytarabine Solution for Injection product monograph. Montreal, Quebec; 14 February, 2017.
119. SteriMax Inc. Cytarabine injection product monograph. Oakville, Ontario; April 19, 2021.
120. Pfizer Canada Inc. Dacarbazine for Injection product monograph. Kirkland, Quebec; 31 May, 2018.
121. Trissel L. Handbook on injectable drugs. 13th ed. Bethesda, Maryland: American Society of Health-System Pharmacists; 2005. p. 428–431
122. Recordati Rare Diseases Inc. COSMEGEN® product monograph. Lebanon, New Jersey USA; 24 July, 2014.
123. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 19 September , 2007.
124. Andy Harbrow. Global Medical Services Manager, Primevigilance (for Recordati Rare Diseases Inc.). Personal communication: dactinomycin solutions for infusion. 15 July, 2014.
125. Janssen Inc. DARZALEX® product monograph. Toronto, Ontario; 29 June, 2016.
126. Janssen Inc. DARZALEX® SC product monograph. Toronto, Ontario; June 22, 2022.
127. Daiichi Sankyo Inc. Dose Preparation Instructions: DS-1062a for Injection 100 mg (Lyophilized Powder). Version 4. Tokyo, Japan; September 10, 2022.
128. ERFA Canada Inc. CERUBIDINE® product monograph. Montréal, Quebec; August 2, 2017.
129. ERFA Canada Inc. Daunorubicin product monograph. Montreal, Quebec; August 15, 2012.
130. Hikma Canada Limited. Daunorubicin Hydrochloride injection product monograph. Mississauga, Ontario; June 28, 2023.
131. Lisa Sorenson, Associate Director. Hikma Canada Limited Quality Department. Personal Communication: Daunorubicin hydrochloride injection. October 23, 2024.
132. Jazz Pharmaceuticals Canada Inc. VYXEOS® product monograph. Mississauga, Ontario; April 28, 2021.
133. Ferring Pharmaceuticals. FIRMAGON® product monograph. North York, Ontario; 20 March , 2013.
134. Ferring Pharmaceuticals. FIRMAGON® product monograph. North York, Ontario; 06 November , 2009.
135. Sandoz Canada Inc. WYOST® product monograph. Mississauga, Ontario; March 1, 2024.
136. Amgen Canada Inc. XGEVA® product monograph. Mississauga, Ontario; 14 October , 2011.
137. Amgen Canada Inc. XGEVA® product monograph. Mississauga, Ontario; May 30, 2023.
138. Juno Pharmaceuticals Corp. Dexrazoxane for injection product monograph. Mississauga, Ontario; December 20, 2023.
139. Pfizer Canada Inc. ZINECARD® product monograph. Kirkland, Quebec; March 30, 2015.
140. Unither Biotec Inc. for United Therapeutics Corp. UNITUXIN® product monograph. Magog, Quebec; 28 November, 2018.
141. Pfizer Canada ULC. Docetaxel injection product monograph. Kirkland, Quebec; March 26, 2021.
142. Josee Lloyd. Senior Clinical Specialist, Hospira Clinical Support Team, Hospira Healthcare Corporation. Personal communication: multidose vials and venting needles. 13 July, 2011.
143. Hospira Canada Clinical Support Team, Hospira Canada Healthcare Corporation. Personal communication. 21 March, 2011.
144. Steinberg RL, Thomas LJ, Brooks N, et al. Multi-institution evaluation of sequential intravesical gemcitabine and docetaxel as rescue therapy for non-muscle invasive bladder cancer. J Urol ; May, 2020;203(5):902–909
145. BC Cancer Genitourinary Tumour Group. (GUBGEMDOC) BC Cancer Protocol Summary for Intravesical Therapy for Non-Muscle Invasive Bladder Cancer Using Gemcitabine and Docetaxel. Vancouver, British Columbia: BC Cancer; 1 August , 2021.
146. Sandoz Canada Inc. Docetaxel injection product monograph. Boucherville, Quebec; 17 November, 2020.
147. Bazundama Bazuta Feza Sandrine. Medical Information Intern, Sandoz Canada Inc. Personal communication: in-house vial stability for docetaxel injection. 14 August, 2018.
148. Bazundama Bazuta Feza Sandrine. Medical Information Intern, Sandoz Canada Inc. Personal communication: in-house product stability of diluted docetaxel injection. 14 August, 2018.
149. Vigneron J, Astier A, Trittler R, et al. SFPO and ESOP recommendations for the practical stability of anticancer drugs: an update. Ann Pharm Fr ; 2013;71(6):376–389
150. Trissel's® 2 Clinical Pharmaceuticals Database - Lexicomp Online (database on the Internet). Docetaxel. Wolters Kluwer Clinical Drug Information Inc.; Accessed 7 July, 2021. Available at: <http://online.lexi.com>
151. GlaxoSmithKline Inc. JEMPERLI® product monograph. Mississauga, Ontario; April 2, 2024.
152. Accord Healthcare Inc. Doxorubicin injection® product monograph. Montreal, Quebec; 9 April, 2014.

153. Trissel's® 2 Clinical Pharmaceutics Database - Lexicomp Online (database on the Internet). Doxorubicin. Wolters Kluwer Clinical Drug Information Inc.; Accessed March 23, 2023. Available at: <http://online.lexi.com>
154. Trissel's®2 IV Compatibility (database on the Internet). Doxorubicin. Merative MICROMEDEX® 2.0; Accessed March 23, 2023. Available at: <http://www.micromedex.com>
155. Gerrie AS, Mikhael JR, Cheng L, et al. D(T)PACE as salvage therapy for aggressive or refractory multiple myeloma. Br J Haematol ; 2013;161(6):802–810
156. Lee C, Barlogie B, Munshi N, et al. DTPACE: An Effective, Novel Combination Chemotherapy With Thalidomide for Previously Treated Patients With Myeloma. J Clin Oncol ; 2003;21(14):2732–2739
157. BC Cancer Myeloma Tumour Group. (UMY0 [D-PACE]) BC Cancer Interim Treatment Plan for ULY0 CAP 112919 (Treatment of Myeloma with CISplatin, Etoposide, DOXOrubicin, Cyclophosphamide and Dexamethasone). Vancouver, British Columbia: BC Cancer; April 1 , 2023.
158. Novopharm Limited. Doxorubicin Product Monograph. Scarborough, Ontario; 8 November, 1996.
159. Pfizer Canada Inc. ADRIAMYCIN® injection product monograph. Kirkland, Quebec; 28 August , 2007.
160. Janssen Inc. CAELYX® product monograph. Toronto, Ontario; 10 October, 2013.
161. Taro Pharmaceuticals Inc. Taro-DOXOrubicin Liposomal product monograph. Brampton, Ontario; September 24, 2020.
162. HealthPro Canada. Personal Communication: Foreign Imported Product: US Labelled TARO-DOXOrubicin Liposomal 20 mg/10 mL (Distributed by Sun Pharmaceuticals Ltd India). October 17, 2024.
163. Barb Pimentel (VP Canadian Commercial) Taro Pharmaceuticals Inc. Personal Communication (Re: TARO-DOXOrubicin liposomal 20 mg/mL) - Importation of US-labelled doxorubicin hydrochloride liposome injection due to a shortage of Canadian-labelled doxorubicin hydrochloride liposome injection. (undated), 2024.
164. Trissel's® 2 Clinical Pharmaceutics Database - Lexicomp Online (database on the Internet). CISplatin + CycloPHOSphamide - Admixture compatibility/ Study 1 /Note. Wolters Kluwer Clinical Drug Information Inc.; Accessed March 9, 2023. Available at: <http://online.lexi.com>.
165. Trissel's®2 IV Compatibility (database on the Internet). Cisplatin and cyclophosphamide admixture compatibility. Merative MICROMEDEX® 2.0; Accessed March 9, 2023. Available at: <http://www.micromedex.com>.
166. AstraZeneca Canada Inc. IMFINZI® product monograph. Mississauga, Ontario; October 25, 2023.
167. McInerney M, Suess J. SBAR: Preparation of Durvalumab Vials without ChemoLock®. Approved by BC Cancer Provincial Pharmacy Professional Practice Committee. June 28, 2023
168. Pfizer Canada-ULC. ELREXFIO® product monograph. Kirkland, Québec; December 6, 2023.
169. Pfizer Inc and Affiliates Global Product Development. PF-06863135 Solution for Subcutaneous Injection (40 mg/mL) Investigational Product Manual Version 1.0. New York, NY; December 7, 2021.
170. Hicham Gonzalez. Medical Liaison. Pfizer Canada Oncology. Personal Communication. March 29, 2023.
171. Michelle Koberinski. BC Cancer Oncology Certification Pharmacy Technician. Personal Communication - Elranatamab. May 30, 2023.
172. Seagen Canada Inc. PADCEV® product monograph. Mississauga, Ontario; October 29, 2021.
173. AbbVie Corporation. EPKINLY® product monograph. St-Laurent, Quebec; October 13, 2023.
174. Novopharm. Epirubicin for Injection product monograph. Toronto, Ontario; 16 March , 2009.
175. Pharmaceutical Partners of Canada I. Epirubicin Hydrochloride Injection product monograph. Richmond Hill, Ontario; 6 July , 2010.
176. Pharmacia Canada Inc. Pharmorubicin PFS Package Insert. Mississauga, Ontario; May, 2003.
177. Trissel LA. Handbook on Injectable Drugs. 12th ed. Bethesda, MD: American Society of Health-System Pharmacists, Inc.; 2003.
178. BC Cancer Agency Lymphoma Tumour Group. (ULYEPOCHR) Interim BCCA Protocol Summary for Treatment of Lymphoma with Dose-Adjusted Etoposide, DOXOrubicin, VinCRISTine, Cyclophosphamide, PredniSONE, and riTUXimab (LYEPOCHR) with Intrathecal Methotrexate. Vancouver, British Columbia: BC Cancer Agency; 1 July , 2015.
179. Barry Goldspiel. NIH Clinical Centre. Personal communication: EPOCHR. 14 April, 2015.
180. Wolfe JL, Thoma LA, Du C, et al. Compatibility and stability of vincristine sulfate, doxorubicin hydrochloride, and etoposide in 0.9% sodium chloride injection. Am J Health-Syst Pharm ; 1999;56:985–989
181. Dunleavy K, Pittaluga S, Shovlin M, et al. Low-intensity therapy in adults with Burkitt's lymphoma. N Engl J Med ; 2013;369:1915–1925
182. Yuan P, Grimes GJ, Shankman SE, et al. Compatibility and stability of vincristine sulfate, doxorubicin hydrochloride, and etoposide phosphate in 0.9% sodium chloride injection. Am J Health-Syst Pharm ; 2001;58(7):594–598
183. BC Cancer Lymphoma Tumour Group. (LYEPOCHR) BC Cancer Protocol Summary for Treatment of Lymphoma with Dose Adjusted Etoposide, DOXOrubicin, vinCRISTine, Cyclophosphamide, predniSONE and riTUXimab with Intrathecal Methotrexate. Vancouver, British Columbia: BC Cancer; January 1 , 2022.
184. Cancer Institute New South Wales. eviQ: Non-Hodgkin lymphoma DA-R-EPOCH (dose adjusted rituximab etoposide prednisolone vincristine CYCLOPHOSPHamide DOXOrubicin) - Treatment schedule overview . St Leonards, New South Wales, Australia: NSW Government.; September 17 , 2022.

185. Eisai Limited. HALAVEN® product monograph. Mississauga, Ontario; January 17 , 2013.
186. Natco Pharma (Canada) Inc. NAT - eribulin product monograph. Mississauga, Ontario; June 22, 2023.
187. Novopharm Limited. Etoposide Product Monograph. Toronto, Ontario; 2000.
188. Lepage R, Walker S, Godin J. Stability and compatibility of etoposide in normal saline. Canadian Journal of Hospital Pharmacy ; December, 2000;53(5):338–345
189. The United States Pharmacopeial Convention, Inc. General Chapter 797: Pharmaceutical compounding - sterile preparations. USP 27-NF 22. Rockville, Maryland: The United States Pharmacopeial Convention, Inc.; 2003.
190. BC Cancer Agency. Provincial Pharmacy Directive III-50-04: Management of Particulate During Sterile Preparation. Vancouver, British Columbia: BC Cancer Agency; 9 July , 2014.
191. Angie Chan. Drug Information Pharmacist N. Personal communication. 29 September, 2006.
192. Bristol-Myers Squibb Company. ETOPOPHOS® product monograph. Princeton, New Jersey, USA; March, 2011.
193. Joseph Atallah. Interim County Medical Director. Bristol-Myers Squibb Canada. Personal communication re: new Canadian distributor for ETOPOPHOS®. November 15, 2019.
194. George Gafrey. VP Business Development. Xediton Pharmaceuticals Inc. Personal communication re: new Canadian distributor for ETOPOPHOS®. November 27, 2019.
195. Bristol-Myers Squibb Company. ETOPOPHOS® product monograph. Princeton, New Jersey, USA; May, 2019.
196. Amgen Canada Inc. NEUPOGEN® product monograph. Mississauga, Ontario; March 21, 2014.
197. Amgen Medical Information. Amgen Canada Inc. Personal communication. 8 July, 2014.
198. Trissel LA. Handbook on Injectable Drugs. 13th ed. Bethesda, Maryland: American Society of Health-System Pharmacists, Inc; 2005. p. 648–655
199. Pfizer Canada-ULC. NIVESTYM® product monograph. Kirkland, Québec; October 27, 2023.
200. Pfizer Medical Information. Personal Communication - NIVESTYM® (filgrastim) Single-dose Vial (SDV) Stability in Syringe. Updated March 14, 2024. Communication Date: April 2, 2024.
201. Pfizer Medical Information. Personal Communication - NIVESTYM® (filgrastim) Stability in Dextrose 5% in Water +/- Albumin. Updated April 3, 2024. Communication Date: April 2, 2024.
202. Accord Healthcare Inc. Fludarabine phosphate injection product monograph. Kirkland, Quebec; September 29, 2021.
203. Teva Canada Limited. Fludarabine phosphate product monograph. Toronto, Ontario; March 1, 2016.
204. Accord Healthcare Inc. Fluorouracil injection® product monograph. Kirkland, Quebec; 30 September, 2013.
205. Charles Vachon. Quality and Regulatory Affairs, Accord Healthcare Inc. Personal communication. 29 September, 2016.
206. John Korontzis. Regulatory Affairs Associate, Mayne Pharma Canada. Personal communication: Fluorouracil. February 16, 2005.
207. Sandoz Canada Inc. Fluorouracil Injection product monograph. Boucherville, Quebec; 3 April , 2012.
208. Alexandre Dussault. Drug Information & Pharmacovigilance Coordinator, Sandoz Canada Inc. Personal communication. 19 November, 2015.
209. Trissel L. Handbook on Injectable Drugs. 13th ed. Bethesda, Maryland: American Society of Health-System Pharmacists; 2005. p. 613–622
210. BC Cancer Agency Experimental Therapeutics. Physicochemical stability analysis of fluorouracil products in final chemotherapeutic preparations. Vancouver, BC. ; ;Study number 50009:1–43
211. Stiles ML, Allen Jr LV, Tu YH. College of Pharmacy, University of Oklahoma, Health Sciences Center, 1110 N. Stonewall, Oklahoma City, OK 73190; United States. Stability of fluorouracil administered through four portable infusion pumps. American Journal of Hospital Pharmacy ; 1989;46(10):2036–2040
212. Accord Healthcare Inc. Gemcitabine injection product monograph. Kirkland, Quebec; 24 March, 2020.
213. Purvi Agrawal BScPharm. Regulatory Affairs Manager, Accord Healthcare Inc. Personal communication. 07 September, 2012.
214. Astron Research LU. Gemcitabine for Injection (STBRG/ACGEM/01).
215. Skinner EC, Goldman B, Sakr WA, et al. SWOG S0353: Phase II trial of intravesical gemcitabine in patients with nonmuscle invasive bladder cancer and recurrence after 2 prior courses of intravesical Bacillus Calmette-Guerin. J Urol ; 2013;190(4):1200–1204
216. BC Cancer Genitourinary Tumour Group. (GUBGEM) BC Cancer Protocol Summary for Intravesical Therapy for Non-Muscle Invasive Bladder Cancer Using Gemcitabine. Vancouver, British Columbia: BC Cancer; 1 November , 2019.
217. Pfizer Canada Inc. Gemcitabine Injection (ready to use solution) product monograph. Kirkland, Quebec; 25 October, 2018.
218. Vachon, Brigitte. Clinical Specialist, Clinical Support. Hospira Healthcare Corporation. Personal Communication: Stability of Gemcitabine Injection (Ready-to-use solution) Diluted in Solution. June 11, 2015.
219. Pfizer Canada ULC. Gemcitabine injection product monograph. Kirkland, Québec; March 16, 2021.
220. Sandoz Canada Inc. Gemcitabine hydrochloride solution for injection product monograph. Boucherville, Quebec; 14 August, 2014.
221. Bazundama Bazula Faza, Sandrine. Drug Information Coordinator. Sandoz Canada Inc. Personal Communication: Gemcitabine injection. July 22, 2019.

222. Pfizer Canada-ULC. MYLOTARG® product monograph. Kirkland, Quebec; 28 November, 2019.
223. Sharon Keane. Pfizer Canada Medical Information. Personal communication. 7 July, 2020.
224. Hoffmann-La Roche Limited. COLUMVI® product monograph. Mississauga, Ontario; Mar 24, 2023.
225. de Lemos, M. BC Cancer Provincial Pharmacy Professional Practice Leader and Drug Information Coordinator. Glofitamab Preparation SBAR. BC Cancer - Systemic Therapy Program ; September 1, 2024
226. Genentech Inc. COLUMVI® full prescribing information. San Francisco, CA, USA; Jun, 2023.
227. Pfizer Canada ULC. Idarubicin hydrochloride injection product monograph. Kirkland, Quebec; September 14, 2021.
228. BC Cancer Provincial Systemic Therapy Program. Provincial Systemic Therapy Program Policy III-20: Prevention and Management of Extravasation of Chemotherapy. Vancouver, British Columbia: BC Cancer; March 1 , 2021.
229. Baxter Corporation. IFEX® product monograph. Mississauga, Ontario; 5 April, 2012.
230. Baxter Corporation. IFEX® product monograph. Mississauga, Ontario; June 27, 2018.
231. Pharmaceutical Partners of Canada I. Ifosfamide for Injection product monograph. Richmond Hill, Ontario; 17 January , 2008.
232. Fresenius Kabi Canada Ltd. Ifosfamide for Injection product monograph. Toronto, Ontario; October 10, 2018.
233. sanofi-aventis Canada. Iniparib (BSI-201;SAR240550) Special Access Program Guidance for the Physician. Laval, Quebec; 15 December, 2010.
234. Pfizer Canada Inc. BESPONSA® product monograph. Kirkland, Quebec; 15 March, 2018.
235. Pfizer Medical Information. Pfizer Canada Inc. Personal communication. 26 November, 2018.
236. Bristol Myers Squibb Canada. YERVOY® product monograph. Montreal, Quebec; 1 February , 2012.
237. Bristol-Myers Squibb Canada. YERVOY® product monograph. Montreal Canada; December 12, 2017.
238. Accord Healthcare Inc. Irinotecan injection® product monograph. Kirkland, Quebec; 6 May, 2014.
239. Auro Pharma Inc. Irinotecan hydrochloride injection product monograph. Woodbridge, Ontario; June 30, 2020.
240. Eugia Pharma Inc. Irinotecan hydrochloride injection product monograph. Woodbridge, Ontario; July 6, 2022.
241. Generic Medical Partners Inc. Irinotecan hydrochloride trihydrate for injection product monograph. Toronto, Ontario; September 21, 2022.
242. Harris A. Regulated Pharmacy Technician Supervisor, BC Cancer Kelowna. Personal communication - GMP brand irinotecan. April 16, 2025.
243. Pfizer Canada Inc. Irinotecan hydrochloride injection product monograph. Kirkland, Quebec; 8 March, 2019.
244. Ipsen Biopharmaceuticals Canada Inc. ONIVYDE® product monograph. Mississauga, Ontario; 17 August, 2023.
245. sanofi-aventis Canada Inc. SARCLISA® product monograph. Laval, Quebec; October 12, 2022.
246. Bristol-Myers Squibb. IXEMPRA® product monograph. Princeton, New Jersey; 01 October , 2007.
247. The United States Pharmacopeia (. General Chapter 797: Pharmaceutical compounding - sterile preparations. USP 27-NF 22. Rockville, Maryland: The United States Pharmacopeial Convention, Inc.; 2004.
248. Kastango ES. The ASHP discussion guide for compounding sterile preparations. Bethesda (MD): American Society of Health-System Pharmacists, Inc.; 2004. p. 5