

Dry Mouth and Mouth Pain from Cancer Treatments

What is dry mouth?

Dry mouth happens when there is less saliva (spit) in your mouth than is normal for you. Dry mouth is also called “xerostomia” (zeer-uh-stoh-mee-uh).

What is oral mucositis?

Oral mucositis (mew-koh-si-tiss) causes redness, sore spots and ulcerations in your mouth and throat. This can make it hard for you to eat, swallow and talk.

Signs of oral mucositis:

- Pain and sores in and around your mouth and throat.
- Sometimes the pain and sores are in the intestinal tract. This can cause pain when you swallow, nausea, diarrhea and sometimes infections.

What causes dry mouth and oral mucositis?

Dry mouth is a common side effect of radiation therapy and some cancer drugs. It is more common in older people.

These cancer treatments can cause redness and sores in your mouth:

- Radiation treatment for head and neck cancers.
- Stem cell transplant therapy.
- Some chemotherapy drugs.



Will these problems get better?

Dry mouth may get better or it may be permanent (it will never go away), depending on what is causing it.

Oral mucositis may continue up to a few months after cancer treatment.

Permanent dry mouth and oral mucositis are more common in patients who have had radiation therapy for head and neck cancer.

When should I talk to my health care team?

Tell your health care team if you have dry mouth, redness, pain or sores in your mouth or throat.

If your oral mucositis is not treated, it can interrupt your cancer treatment. Oral mucositis may also cause you to worry, become depressed, or stop seeing your friends and family.

What are the treatments?

Before your cancer treatments start, you should visit your dentist. They will examine you and treat any dental problems. Problems like cavities, gum disease, infections, and broken or sharp teeth should be treated or fixed.

If you have dentures your dentist will also check that they fit well and are comfortable.

Your health care team might prescribe medications to help with your mouth symptoms. They may give you a special oral rinse (mouthwash) or tell you to suck on ice chips.

Do not use any other over-the-counter mouth rinses or gels. These may make your pain worse.

If you having trouble eating and you are losing weight, your health care team may refer you to a BC Cancer dietitian.

What can I do for myself?

The most important thing you can do is to take care of your mouth. Start following the directions below two weeks before your treatment starts or as soon as you can.

If you grind or clench your teeth, wear your mouth guard to protect your teeth.

Brush your teeth

- Brush your teeth for at least 2 minutes.
- Brush within 30 minutes of eating.
- Use a soft toothbrush.
- If you have oral mucositis, use an SLS-free toothpaste. Check the ingredients on your toothpaste to make sure it does not contain SLS (Sodium Lauryl Sulphate).
- If your mouth is very sore, do not use mint flavoured toothpaste

If your mouth is too sore to use a toothbrush, talk to your health care team.

Floss your teeth

- If flossing is part of your routine, continue to floss once per day.
- Floss at bedtime, before brushing your teeth.
- Use waxed floss. This will slide between your teeth easier than unwaxed floss.
- If flossing is not part of your routine, do not start flossing unless your dentist tells you to.
- Stop flossing if it is painful or your gums bleed for longer than 2 minutes. Tell your health care team if this happens.
- If you use a Water Pik, make sure you put it on the lowest setting so you do not hurt your gums.

Use an oral rinse

- Oral rinses help keep your mouth moist and clean.
- Use this recipe for mouth rinse:
 - Mix 5 mL (1 teaspoon) of baking soda and 5 mL of salt with 1000 mL (4 cups) of water.
 - Put the mouth rinse in a container with a lid.
 - Keep mouth rinse at room temperature.
 - Throw away the mouth rinse at the end of the day and make a new batch.
- How to use your mouth rinse:
 - Shake well before using.
 - Rinse and gargle with 15 mL (1 tablespoon) and then spit out.
 - Repeat 2 or 3 time at each use.
 - Use mouth rinse every 2 hours during the day.
- Use the oral rinse until your mouth returns to normal. This may take many weeks or months.
- Use mouth rinse or water after eating or drinking.

- Rinse after drinking meal replacement drinks or sports drinks as these have a lot of sugar.
- Use a soft toothbrush and gently brush your teeth or dentures after every meal. Use the toothbrush to clean your tongue after you brush. Remove dentures at night.

If you have pain or infection, your doctor will prescribe medication for you.

Food and drink

- Drink at least 1.5-2 litres (6-8 cups) of fluids each day. Fluids include water, juice (not acidic juices like orange juice as these may hurt your mouth), herbal tea, soup, broths, sports drinks and liquid nutritional drinks like Ensure®.
- Do not drink carbonated drinks (fizzy drinks) like pop. These are very acidic and can cause mucositis, more mouth pain and cavities.
- Do not drink alcohol as this will hurt your mouth.
- Eat bland, soft foods. You need to eat a balanced diet with enough protein and vitamins.
- Talk to a BC Cancer dietitian if you are unsure about what foods to eat and drink.

Take care of your lips

- If your lips are dry, use a lubricant that is water-soluble, wax-based, or oil-based. Lanolin-based lubricants are the best choice.
- Do not use petroleum-based lubricants, like Vaseline.
- Put the lubricant on your lips after cleaning your mouth and teeth and at bedtime and as needed.
- Ask your health care team if you are not sure which lubricant to use.

Taking care of dentures

- Leave dentures out of your mouth when you are sleeping.

- As much as possible during your cancer treatment, leave your dentures out of your mouth during the day.
- Take out your dentures before brushing your teeth.
- Brush and rinse dentures after meals and before bed.
- While you are sleeping, soak your dentures in a cleansing solution for at least 8 hours.
- If you are using an anti-fungal mouth rinse, soak the dentures in the anti-fungal rinse. Or, brush your dentures with your anti-fungal rinse before putting the dentures in your mouth.

Some things you should not do:

- Do not use alcohol-based mouthwashes or over-the-counter rinses. Even alcohol-free rinses may sting due to chemicals in it. It's best to use the bland mouth rinse.
- Do not eat spicy, acidic or coarse foods.
- Do not drink alcohol.
- Do not smoke.

Talk with your health care team if you are having trouble with any of these instructions.

Where can I learn more?

BC Cancer Library Mouth and Jaw Problems pathfinder: bccancer.libguides.com/pathfinder-mouth-jaw

BC Cancer Nutrition Handouts: www.bccancer.bc.ca/health-info/coping-with-cancer/nutrition-support

International Society of Oral Oncology Patient Care resources (multilingual): <http://www.isoo.world/patient-education.html>

BC Cancer Supportive Care Services: www.bccancer.bc.ca/our-services/services/supportive-care

