

# Symptom Management Guidelines: INTIMACY AND SEXUALITY

### Definition(s)

- **Sexuality:** Encompasses sexual behaviors, gender identities and roles, sexual orientation, eroticism, pleasure, intimacy, and reproduction. Is a complex phenomenon which encompasses our biological being, sense of self, and the way in which we relate to others.
- Intimacy: Sharing of identity, mutual acceptance, closeness and reciprocated rapport more closely linked to communication than sexual function. Intimacy, like sexuality, is intrinsic to a sense of self.

Contributing Factors		
Cancer Treatment Related	<ul> <li>Surgery         <ul> <li>Depending on surgical site, may result in infertility, altered sexual function and/or alterations in body image (e.g. prostate surgery, testicular surgery, hysterectomy, oopherectomy, vaginectomy, vulvectomy, abdomnioperineal resection, mastectomy)</li> </ul> </li> <li>Chemotherapy             <ul> <li>Can cause temporary or permanent loss of ovarian and testicular function resulting in infertility for both, and premature menopausal symptoms for women</li> <li>Sensory neuropathy in the genitals</li> </ul> </li> <li>Radiation therapy             <ul> <li>Radiation to the pelvic area can cause vascular or nerve damage which impairs sexual functioning either temporarily or permanently</li> <li>Women: Vaginal atrophy, stenosis, fibrosis, or shortening, amenorrhea, painful intercourse, decreased lubrication, ovarian failure, premature menopause, diminished libido</li> <li>Men: Decreased testosterone secretion, testicular aplasia, diminished semen volume, ejaculatory pain, diminished libido</li> <li>Hormonal Agents</li> <li>May result in weight loss or gain, menopausal symptoms, diminished libido, erectile dysfunction, masculinization in women, emasculation in men</li> </ul> </li> <li>Immunotherapy - Checkpoint Inhibitors</li> <li>Diminished libido, mood changes, weight gain or loss or fatigue due to immune mediated side effects.</li> </ul> <li>Biologic Agents</li> <li>Mucous membrane dryness, diminished libido, erectile dysfunction, pelvic pain</li>	
Symptoms	<ul> <li>Physical symptoms can alter comfort, interest, desire, and ability to engage in sexual activities (e.g. fatigue, pain, dyspnea, nausea)</li> <li>Psychological distress (e.g. anxiety, depression, fear, uncertainty)-can decrease one's energy, ability to concentrate and relax</li> <li>Changes in bowel or bladder functioning e.g. ostomy</li> <li>Changes in mobility and Range of motion- may inhibit ability to comfortably engage in sexual positions</li> </ul>	
Side Effects of Medications	<ul> <li>Many medications used for symptom management (e.g. some anti-emetics, opioids, anxiolytics, antidepressants, antihypertensives, antihistamines) have sedative effects, sense dulling properties, reduce libido and/or interfere with erectile function</li> <li>Vaginal dryness and other menopausal symptoms</li> </ul>	
Body Image Disturbance	<ul> <li>Related to weight changes, hair loss, surgical scarring, changes in elimination patterns (e.g. ostomies, catheters)</li> </ul>	
Partner Concerns	<ul> <li>Emotional distress, fears about hurting patient</li> <li>Role shifting/confusion-intimacy can be difficult when partner has various roles- caregiver, partner, lover.</li> </ul>	

### Consequences

- Altered perceptions of body image, self-esteem, self-confidence
- Psychological distress, feelings of isolation and/or alienation
- · Changes in relationships
- Aversion or reluctance to engage in activities of sexual/intimate expression

The information contained in these documents is a statement of consensus of BC Cancer Agency professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is at your own risk and is subject to BC Cancer Agency's terms of use, available at <a href="https://www.bccancer.bc.ca/legal.htm">www.bccancer.bc.ca/legal.htm</a>.

### **Focused Health Assessment**

- Modify assessment to accommodate variations in patient age, gender, culture, sexual orientation and stage of illness
- Include patient's partner, as appropriate

Contact and General Information	GENERAL ASSESSMENT	SYMPTOM ASSESSMENT	PHYSICAL ASSESSMENT
Allergies  Consider Contributing Factors     Cancer diagnosis and treatment(s) – note type & date of last treatment     Medical history     Medical history     Medication profile     Recent lab or diagnostic reports  Treatment     What aspects of your sexuality and intimacy?  Understanding / Impact on You     What aspects of your sexuality do you believe have been affected by your cancer or treatment (positive or negative)?     Have you talked to your partner about your feelings?  Value     How important is your sexuality to you?     What do you believe is causing your difficulties?     Presence of wounds, surgical scars, degree of healing     Presence of alopecia  Abdominal Assessment     Abdominal Assessment     Abdominal and pelvic pain and tenderness  Range of Motion     With or without discomfort  Genitourinary     Note any changes in continence or structure  Weight     Note any significant changes in weight gain or loss  Value     How important is your sexuality to you?     What do you think is responsible for the changes in your	General Information  Physician name - oncologist, family physician Pharmacy Home health care Other healthcare	<ul> <li>Do you feel that your diagnosis has led to changes or difficulties in your sexual feelings or expression? (if so, in what way(s)?)</li> <li>How do you express your sexuality (before your diagnosis? Now?)</li> </ul> Onset	<ul> <li>Note mood, affect; cognitive status; general physical status</li> <li>Hydration Status</li> <li>Mucous membranes – moist, dry?</li> </ul>
sexual life?  • What are your goals for intimacy and sexuality?	<ul> <li>Allergies</li> <li>Consider         Contributing         Factors         <ul> <li>Cancer diagnosis and treatment(s) – note type &amp; date of last treatment</li> <li>Medical history</li> <li>Medication profile</li> <li>Recent lab or</li> </ul> </li> </ul>	<ul> <li>What do you believe is causing your difficulties?</li> <li>What makes it better? Worse?</li> <li>Quality</li> <li>Can you describe your concerns regarding sexuality and intimacy?</li> <li>Severity</li> <li>How bothersome are your concerns?</li> <li>Treatment</li> <li>Have you tried to do anything to address your concerns regarding sexuality and intimacy? Was this successful?</li> <li>Understanding / Impact on You</li> <li>What aspects of your sexuality do you believe have been affected by your cancer or treatment (positive or negative)?</li> <li>How has this treatment/experience affected your relationship with your partner? Or your partner's sexuality?</li> <li>Have you talked to your partner about your feelings?</li> <li>Value</li> <li>How important is your sexuality to you?</li> <li>What do you think is responsible for the changes in your sexual life?</li> </ul>	<ul> <li>Presence of wounds, surgical scars, degree of healing</li> <li>Presence of alopecia</li> <li>Abdominal Assessment</li> <li>Abdominal distention, Abdominal and pelvic pain and tenderness</li> <li>Range of Motion</li> <li>With or without discomfort</li> <li>Genitourinary</li> <li>Note any changes in continence or structure</li> <li>Weight</li> <li>Note any significant changes in weight gain or</li> </ul>

## Management of Concerns Related to Intimacy and Sexuality

### Goals

- Establish goals for sexuality and intimacy and expected outcomes with patient and if applicable, their intimate partner
- Restore optimal intimacy/sexual practices by reducing or eliminating contributing factor(s); provide symptom management as necessary

Are you interested in expressing your sexuality now?

- Maintain optimal physical health to enjoy and engage in intimacy/sexuality
- Minimize risk of consequences resulting from changes in intimacy or sexuality

Key Consideration	s
Privacy	<ul> <li>Consider appropriate environment to raise sensitive topics</li> <li>Consider person's sense of personal privacy in relation to openly discussing this topic</li> </ul>
Beliefs and Values	Consider cultural beliefs and values that may impose constraints on discussing sexuality and intimacy; respect and/or negotiate these boundaries
Communication	<ul> <li>The nurse's role in intimacy/sexuality support is focused on communication:         <ul> <li>Raising the topic</li> <li>Assessing patient and if appropriate, their partner's concerns</li> <li>Encouraging open communication where patients and their partners can communicate honest feelings, concerns, fears, and preferences</li> <li>Validating their concerns and sharing with other healthcare professionals as appropriate</li> </ul> </li> <li>Think broadly about intimacy and sexuality and the meaning for patient and their partner (e.g. desire for closeness, affection, reproduction)</li> </ul>
Immunotherapy	Decreased libido, weight gain or loss, mood changes, fatigue, irritability or headaches can be a sign of endocrinopathy or an adrenal crisis brought on by immune medicated side effects of Immunotherapy. Refer to protocol specific algorithms and consult with physician.
Sexual Orientation	Openly and non-judgmentally explore the sexual orientation of the patient
Capacity for Sexual Activity or Desire	<ul> <li>A person's capacity and desire for sexual activity is very personal and individualized</li> <li>Explore goals and help the patient and their partner achieve their desires/mutual pleasure to the best of their ability.</li> <li>Patients and their partners may need assistance in accepting changes in patterns of desire and ability to engage in sexual activities. Do not minimize or overestimate these concerns.</li> </ul>
Patient Education: Birth Control and Fluid Protection	<ul> <li>Discuss the common sexual side effects from cancer treatments</li> <li>Prevent pregnancy while receiving chemotherapy or radiation treatment</li> <li>Speak with your healthcare provider to determine best option of birth control for you</li> <li>Use a barrier (e.g. condoms, dental dam) for first 48 hours after chemotherapy to avoid exposure to partner</li> </ul>
Reproduction	Discuss sperm banking and freezing embryos as necessary prior to cancer-treatment delivery

P-LI-SS-IT MODEL	
	des the health professional through assessment and intervention where difficulties arise in intimacy of intervention should be matched with the intensity of the issue.
Giving Permission	<ul> <li>Invite patient to discuss his/her concerns and ask questions in a safe environment</li> <li>Validate concerns and dispel any myths and misconceptions (reinforce that cancer is not passed from person to person)</li> <li>May start conversation by normalizing the topic (e.g. many people with cancer often have questions about sexuality and sexual functioning. Do you have any concerns or questions that you would like to discuss?)</li> <li>Normalize broad definition of sexuality and intimacy (see definitions above)</li> <li>Encourage patient to communicate with partner about cancer and sexuality</li> <li>Give permission for patient and partner to engage in intimate and sexual activities</li> </ul>
Limited Information	<ul> <li>Provide basic information related to sexual functioning and the impact of illness and associated treatments on intimacy and sexuality</li> <li>Provide information regarding altering sexual practices in response to changes in physiology and function</li> <li>Provide relevant patient education and community resources regarding sexuality and cancer See Resources and Referrals Section below</li> </ul>

Specific	Assessment of the relationship may reveal concerns (emotional or psychological) and a referral	
Suggestions	to a counselor may assist.	
	<ul> <li>Ask what activities they have enjoyed doing together in the past. Encourage couple to sit on the couch or nap together, hold hands, hug, play games, or go on dates to build intimacy</li> <li>Sensate focus exercises can assist couples to rebuild intimacy without the pressure to have</li> </ul>	
	intercourse Appendix A: Sensate Focus Exercises	
	<ul> <li>Encourage foreplay, mutual pleasuring, self-stimulation, use of sexual aids, alternatives to penetrative intercourse, and positioning to reduce discomfort and associated symptoms.</li> <li>Describe options for psychosocial and medical interventions to enhance sexual functioning</li> </ul>	
	<ul> <li>Assessment for body image disturbances as a result of treatment or disease (i.e. breast enlargement, surgery, scars)</li> </ul>	
	Women:	
	<ul> <li>Vaginal dryness-water-based or silicone lubricants that are glycerin free. Vaginal estrogen cream as appropriate</li> </ul>	
	<ul> <li>To help prevent shortening of the vagina after radiation therapy, encourage women to continue sexual intercourse during therapy as tolerated. Alternatively, women may use a vaginal dilator on a daily basis or as directed by radiation-oncologist. A water-soluble lubricant should be used for intercourse and dilator use.</li> </ul>	
	Men:	
	- Use of medications (e.g. Sildenafil citrate)	
	Erectile aids (e.g. vasoactive agents, vacuum devices, penile prostheses)	
Intensive Therapy	<ul> <li>When guidance requires expertise beyond the nurse's scope, referral should be made (for individual and/or couple) See Resources &amp; Referrals Section below</li> </ul>	

	RESOURCES & REFERRALS
Referrals	<ul> <li>Counselor</li> <li>Mental health professional</li> <li>Sydney Thomson (physician specialized in supporting sexual dysfunction, available in lower mainland through referral)</li> <li>Sex therapist</li> <li>Reproductive endocrinologist</li> <li>Look Good Feel Better program <a href="http://www.lgfb.ca/en/">http://www.lgfb.ca/en/</a></li> </ul>
Resources in BC	<ul> <li>BC Association For Marriage and Family Therapy: <a href="http://www.bcamft.bc.ca/">http://www.bcamft.bc.ca/</a></li> <li>BC Psychological Association: <a href="http://www.psychologists.bc.ca/">http://www.psychologists.bc.ca/</a></li> <li>BC Association of Registered Clinical Counsellors: <a href="http://www.bc-counsellors.org/">http://www.bc-counsellors.org/</a></li> <li>BC Centre for Disease Control Smart Sex Resource: <a href="http://smartsexresource.com/">http://smartsexresource.com/</a></li> <li>Canadian Fertility and Andrology Society (Fertility Services): <a href="http://www.cfas.ca/index.php?option=com_content&amp;view=article&amp;id=259&amp;Itemid=274">http://www.cfas.ca/index.php?option=com_content&amp;view=article&amp;id=259&amp;Itemid=274</a></li> </ul>
Patient Education	<ul> <li>Canadian Cancer Society:         <ul> <li>Sexuality and Cancer Booklet:</li></ul></li></ul>

	<ul> <li>Sexual Side Effects in Men         <ul> <li>http://www.cancer.org/treatment/treatmentsandsideeffects/physicalsideeffects/sexualsideeffectsimmen/index</li> </ul> </li> <li>Sexual Side Effects in Women         <ul> <li>http://www.cancer.org/treatment/treatmentsandsideeffects/physicalsideeffects/sexualsideeffectsinwomen/index</li> </ul> </li> <li>Mayo Clinic Website:         <ul> <li>Sexual Side Effects For Men: <a href="http://www.mayoclinic.com/health/cancer-treatment/SA00070">http://www.mayoclinic.com/health/cancer-treatment/SA00070</a> </li> </ul> </li> <li>Sexual Side Effects for Women:         <ul> <li>http://www.mayoclinic.com/health/cancer-treatment/SA00071</li> </ul> </li> <li>Library Pathfinder: Sexuality and Partner Support         <ul> <li>http://www.bccancer.bc.ca/library-site/Documents/pathfinder_Sexuality-General.pdf</li> </ul> </li> <li>Resources about managing deep breathing, progressive muscle relaxation, positive thinking, etc. Located in Patient Handouts         <ul> <li>http://www.bccancer.bc.ca/health-info/coping-with-cancer/emotional-support/managing-stress</li> </ul> </li> </ul>
Immunotherapy	Immunotherapy Alert Card     Please refer to protocol specific algorithms to guide management of immune mediated side effects.
Health Professional Info	Sexuality and Prostate Cancer – Dr. Anne Katz <a href="https://www.youtube.com/watch?v=fZLOR7bLHa4">https://www.youtube.com/watch?v=fZLOR7bLHa4</a>
Related Online Resources	E.g. Fair Pharmacare; BC Palliative Benefits. Can be found in "Other Sources of Drug Funding Section" <a href="http://www.bccancer.bc.ca/health-professionals/professional-resources/pharmacy/drug-funding">http://www.bccancer.bc.ca/health-professionals/professional-resources/pharmacy/drug-funding</a>
Bibliography List	http://www.bccancer.bc.ca/health-professionals/professional-resources/nursing/symptom- management

### Date of Print:

Revised: October, 2014 Original: January, 2010

#### **Contributing Authors:**

Revised: Judy Oliver, BScN MEd; Lindsay Schwartz, RN MSc(A); Jagbir Kaur, RN MN Created by: Vanessa Buduhan, RN MN; Rosemary Cashman, RN MSc(A) MA (ACNP); Elizabeth Cooper, RN BScN CON(c); Karen Levy, RN MSN; Ann Syme, RN MSN

Reviewed by: Theressa Zapach, RN, BScN CONc

### Appendix A: Sensate Focus Exercises (Sexual Health Counseling & Cancer – IPODE April 8, 2010)

Definition		
	<b>Sensate Focus Exercises</b> : Series of exercises for couples that encourage each partner to give and receive pleasurable touch while paying attention to their sensual experience of receiving and giving touch	
Stage I	<ul> <li>Each partner takes turns touching each other's body (touching breasts, genitals and intercourse are off limits to start)</li> <li>Encouraged to suspend judgments and expectations</li> <li>Directed to focus their awareness on sensations</li> <li>Partner who is touching is to do so based on what interests them</li> <li>Partner who is the receiver lies on back and receives touch for 15 minutes then receives touch on back of body for 15 minutes; after 30 minutes partners switch roles</li> <li>Encourage couples to be as silent as possible however to let their partner know verbally or non-verbally if any touch is uncomfortable</li> <li>At end of experience, partners give each other specific feedback (3 things enjoyed and 1 thing they would change for next time)</li> </ul>	
Stage II	<ul> <li>Touching now includes breasts and genitals (intercourse off limits)</li> <li>Start with general body touching first</li> <li>As in Stage I, emphasis on awareness of sensations and partners take turn touching and receiving for 15 minutes on front and back</li> <li>Non-verbal communication (receiver places hand on partners hand to indicate if they want a change in pressure, pace, or area)</li> <li>At end of experience, partners give each other specific feedback (3 things enjoyed and 1 thing they would change for next time)</li> </ul>	
Stage III	Couples take turns touching and receiving but can now progress to helping each other experience orgasm through oral or manual stimulation	
Subsequent Stages	Couples slowly progress towards intercourse	