



Provincial Health Services Authority

Summary of BC Cancer’s Pharmacy Practice Standards for Hazardous Drugs

Table of Contents

- Pharmacy Oncology Certification5**
- Module 1 – Safe Handling of Hazardous Drugs.....5**
- Section A - Safe Handling Principles.....5**
 - A.1 Potential Hazards of Handling Hazardous Drugs..... 5
 - A.2 Hazardous Drug List 5
 - A.4 Personal Exposure Records 5
 - A.5 Work Re-Assignment..... 6
- Section B - Decontaminating, Cleaning, Deactivating, Disinfecting, and Sporocidals.....6**
 - B.1 General 6**
- Section C - Controlled Area.....6**
 - C.2 Room Specifications..... 6
 - C.2.1 HEPA Filter 6
 - C.2.2 Signage 6**
 - C.2.3 Hazardous Drug Clean Room..... 7
 - C.2.4 Hazardous Drug Anteroom..... 7
 - C.2.5 Hazardous Drug Storage Room..... 7**
 - C.2.7 Specifications of the Controlled Area..... 7**
- Summary of Engineering Control Requirements for Hazardous Drug Sterile Product Preparation and Storage Areas 8**
- C.3 Equipment and Furniture 9**
 - C.3.1 Containment Primary Engineering Control 9**
 - C.3.1.1 Class I BSC..... 9
 - C.3.1.2 Class II BSC 9
 - C.3.1.3 HEPA Filter..... 9
 - C.3.1.4 Airflow..... 9
 - C.3.1.5 Ultraviolet Lights..... 9

C.3.1.6 Viewing Window	10
C.3.1.7 Location	10
C.3.1.8 Monitoring	10
C.3.1.9 Testing and Certifying Biological Safety Cabinets	10
C.3.1.10 Replacing HEPA Filters	10
C.3.1.11 Turning off a Containment Primary Engineering Control (C-PEC)	10
C.3.1.12 Decontaminating, Cleaning, and Disinfecting Interior Surfaces of the Containment Primary Engineering Control (C-PEC)	11
C.3.2 Communication System	12
C.3.4 Chairs	12
C.4 Decontaminating, Cleaning, Deactivating, and Disinfecting Surfaces in the Controlled Area	12
Section D - Protective Measures	12
D.1 General	12
D.2 Personal Protective Equipment (PPE) and Clean Room Garb	12
D.2.1 Scrubs	13
D.2.2 Footwear	13
D.2.3 Shoe Covers	13
D.2.4 Hair Covers	13
D.2.5 Medical Masks	14
D.2.6 Respirators	14
D.2.6.2 Chemical Cartridge Respirator	14
D.2.7 Chemotherapy Gowns	14
D.2.8 Isolation Gowns	15
D.2.9 Chemotherapy Gloves	15
D.2.10 Eye Protection	16
D.3 Hand Hygiene	16
D.3.1.1 Alcohol-Based Hand Rub	16
D.3.2 Hand Washing After Handling Hazardous Drugs	16
D.3.2.1 Compounding personnel	16
D.3.3 Nails and Nail Polish	16
D.4 Safety Stations	17
D.4.1.2 Hand Held Portable	17
D.4.2 Emergency Showers	17
D.4.3 Safety Stations Maintenance	17
Section E - Supplies and Devices	17
E.1 Supplies	17
E.1.3 Sterile Alcohol Swabs	17
E.2 Devices	17
E.2.1 Syringes	18
E.2.2 Syringe Tip Caps	18
E.2.3 Needles	18
E.2.4 Needle Caps	18
E.2.7 Filters	18
E.2.7.1 Filter Needles	18
E.2.7.2 Filter Discs	18
E.2.8 Filter Venting Devices	18
E.2.8.1 Chemotherapy Dispensing Pins	18
E.2.8.2 Chemotherapy Vents	19

E.2.9 Syringe Fluid Dispensing Connectors/Syringe Tip Connectors	19
E.2.14 Closed System Drug Transfer Devices.....	19
E.3 Containers	19
E.3.2 Vials.....	19
E.3.4 Non-Di(2-ethylhexyl)phthalate (Non-DEHP) Bags.....	19
E.3.5 Empty Sterile Infusion Bags	19
E.4 Ambulatory Drug Delivery Infusion Devices.....	20
E.4.1.2 Infusor™ Flow Rates.....	20
E.4.1.4 Hazardous Drug Medication Infusion Device Labels	20
E.4.2 Computerized Ambulatory Drug Delivery (CADD®) Pump and Medication Cassette Reservoir	20
Section F - Aseptic and Protective Processes	20
F.1 Operational Standards for Sterile Hazardous Drug Preparation	20
F.1.1 Personnel Hygiene in the Controlled Area.....	20
F.1.2 Personal Protective Equipment and Clean Room Garb in the Controlled Area	20
F.1.3 Containment Primary Engineering Control (C-PEC)	21
F.1.4 General Procedures.....	22
F.1.5 Removing Products from the Containment Primary Engineering Control (C-PEC)	23
F.1.6 Warning Labels	23
F.1.7 Exiting the Clean Room	23
F.2 Aseptic/Protective Routines.....	24
F.2.1 Critical Sites.....	24
F.2.2 First Air	24
F.2.3 Disinfecting Critical Sites	24
F.2.4 Coring	24
F.2.5 Safely Capping Needles Used With Hazardous Drug.....	24
F.3 Safe Handling Aseptic Techniques	25
F.3.1 Transfer of Hazardous Drug Solution from a Syringe.....	25
F.3.2 Removal of Bubbles/Air from a Syringe	25
F.3.3 Attaching and Priming Solution / Secondary Administration Sets.....	25
Section G - Clean Up and Waste Disposal.....	25
G.1 Containment Primary Engineering Control (C-PEC) Waste Cleanup	25
G.2 Hazardous Waste Disposal.....	25
Section H - Safe Handling of Oral, Topical and Pre-Packaged Hazardous Drug Dosage Forms	26
H.1 Oral Preparations	26
H.2 Topical Preparations.....	26
Section I - Hazardous Drug Spills	26
I.1 Hazardous Drug Spills.....	26
I.1.1 Recommended Spill Kit Contents	26
Section J - Accidental Exposure to Hazardous Drugs.....	27
J.1 Accidental Exposure to Hazardous Drugs	27

J.1.2 Ingestion	27
Section K - Receipt, Storage, and Transportation of HD	27
K.1 Receipt and Unpacking	27
K.1.1 Receipt	27
K.1.2 Receipt of a Damaged Shipment	27
K.2 Storage	27
K.3 Packaging and Transportation	28
Module 2 – Pharmacy Medication Checks	28
Section A.....	28
A.1 Clinical Medication Order Check.....	28
Section B.....	29
B.1 Documentation of Pharmacy Medication Checks.....	29
Module 3 – Improving Patient Safety and Compliance	29
Section A.....	29
A.1 Intrathecal (IT) Doses.....	29
Section B.....	30
B.1 Vinca Alkaloids.....	30
Section D.....	30
D.1 Outpatient Hazardous Drug Prescription Labelling.....	30
Section E.....	30
E.1 Latex Allergies.....	30
References	31

Pharmacy Oncology Certification

All pharmacy staff involved in the preparation and delivery of hazardous drugs must demonstrate knowledge and competency for the duties that they are required to undertake. Re-evaluation of competency and knowledge must take place on a regular basis with documentation of results.¹

Module 1 – Safe Handling of Hazardous Drugs

Section A - Safe Handling Principles

A.1 Potential Hazards of Handling Hazardous Drugs

Hazardous Drug (HD) safe handling policies and procedures must be developed to address the following:^{2,3}

- Receipt^{3,4}
- Storage^{1,3}
- Preparation^{2,3}
- Labelling¹
- Safety equipment⁵
- Use of Personal Protective Equipment (PPE)^{4,6}
- Emergency procedures for treating accidental contact and spills^{3,4}
- Packaging¹
- Transport¹
- Drug administration³
- Disposal^{3,4}

All pharmacy staff must be informed of HD policies and procedures, and receive training for handling hazardous drugs safely, cleaning up spills, and using all equipment and PPE properly.^{3,4} There must be established work practices related to both drug manipulation techniques and to general hygiene practices.⁴ Workplace procedures must be developed for using and maintaining all equipment that functions to reduce hazardous drug exposure.³

A.2 Hazardous Drug List

Each facility must develop and maintain a hazardous drug list to ensure that healthcare staff working in the facility is made aware of which drugs are hazardous.^{3,4}

The facility's hazardous drug list must be posted in all areas where these drugs are received,^{7,8} stored,^{3,8} prepared³ and administered.^{7,8}

A.4 Personal Exposure Records

WorkSafe BC Occupational Health and Safety (OH&S) [Regulation 6.52](#) states:

1. *“The employer must maintain a record of all workers who prepare or administer cytotoxic (hazardous) drugs, including the name of the drugs handled, and when practicable, the number of preparations or administrations per week.*
2. *Exposure records must be maintained for the duration of employment plus 10 years, and training records for 3 years from the date that the training occurred.”*

A.5 Work Re-Assignment

WorkSafe BC Occupational Health and Safety (OH&S) Regulation 6.49 Reproductive toxins states:

1. “At any worksite where a worker is occupationally exposed to a cytotoxic drug that is a reproductive toxin, the employer must develop policy and procedures appropriate to the risk, which may include protective reassignment.”
2. “The policy and procedures must inform workers about the reproductive toxin and identify ways to minimize exposure to the reproductive toxin for a worker who has advised the employer of pregnancy or intent to conceive a child.”

Section B - Decontaminating, Cleaning, Deactivating, Disinfecting, and Sporicidals

B.1 General

To minimize hazardous drug and environmental (e.g., microbial and particulate) contamination, surfaces must be appropriately decontaminated, cleaned, and disinfected by personnel trained and qualified to safely carry out their responsibilities.¹ All personnel performing these activities must wear appropriate personal protective equipment resistant to the solutions used.⁴ Solutions selected must be appropriate for the type of hazardous drug contaminants, location, and surface materials.⁴

When selecting decontaminating, cleaning, deactivating and disinfecting solutions, consideration must be given to compatibility, effectiveness, and inappropriate or toxic residues.¹ Diluted solutions must be prepared and stored according to the manufacturer's directions¹ and kept in previously cleaned containers.

Partly emptied containers must not be topped up.⁷

Where applicable, the manufacturer's directions regarding the required contact time between the solution and the surface to be disinfected must be followed in order for the disinfectant to be effective (e.g., germicidal disinfecting detergents, sporicidals).¹ When sterile 70% isopropyl alcohol is used, it must be allowed to dry.⁹

Section C - Controlled Area

C.2 Room Specifications

Access to the controlled area must be limited to authorized personnel who are assigned to work there. All personnel entering the controlled area must follow appropriate hand hygiene and garbing procedures as the first major step in preventing microbial contamination of compounded sterile preparations¹ and to minimize healthcare workers' exposure to hazardous drugs.¹⁰

Doors leading into controlled areas must not be left open.¹

C.2.1 HEPA Filter

The air supplied to rooms in the controlled area must pass through at least one HEPA filter to help ensure a very high level of cleanliness. The intake air must come from the ceiling via diffusers, each fitted with a terminal HEPA filter.¹

C.2.2 Signage

Each room in the controlled area must be identified with appropriate informative signs (e.g., restricted access, dress code).¹

Warning signs, which are clearly visible and clearly state the identified hazards, must be posted in all areas where hazardous drugs are received,⁷ stored,³ prepared³ and administered.⁷

C.2.3 Hazardous Drug Clean Room

The hazardous drug clean room must maintain an ISO Class 7 environment and be 2.5 Pa negative pressure to the anteroom. The clean room must also maintain at least 30 air changes per hour (ACPH) of HEPA-filtered air.¹

The return air from the hazardous drug clean room must be exhausted to the exterior of the building.¹

C.2.4 Hazardous Drug Anteroom

There must be a process in place that allows only one door leading into the anteroom to be open at one time.¹

The hazardous drug anteroom must maintain an ISO Class 7 environment and be positive pressure to both the hazardous drug clean room(s) (2.5 Pa) and the rest of the pharmacy (5 Pa). The anteroom must maintain at least 30 air changes per hour (ACPH) of HEPA-filtered air.¹

C.2.5 Hazardous Drug Storage Room

Hazardous drugs requiring manipulation other than counting or repackaging of final dosage forms¹¹ must be stored in a dedicated room that:

- maintains at least 2.5 Pascals negative pressure relative to surrounding areas;¹
- maintains at least 12 Air Changes Per Hour (ACPH), with all air exhausted to the exterior;¹
- contains shelves with lips to help prevent drug containers from falling and breaking;¹ and
- has sufficient ventilation to prevent contamination from spreading to adjoining rooms.¹

The storage bins or shelves must be identified with proper signage to indicate the presence of hazardous drugs.³

Storage conditions specified by the manufacturer must be strictly adhered to (e.g., refrigerator versus room temperature).¹ Alternative storage must be provided if the storage temperature exceeds acceptable variations and when refrigerators and freezers are being cleaned.¹

C.2.7 Specifications of the Controlled Area

- Access to the controlled area must be limited to authorized personnel assigned to work there.^{1,2}
- Doors leading into controlled areas must not be left open.^{1,2}
- The pressure in the anteroom must be positive. The pressure differential between the anteroom and the clean room must be at least 2.5 Pa (equivalent to 0.01 inch water column).¹ The pressure differential between the anteroom and the rest of the pharmacy must be at least 5 Pa (equivalent to 0.02 inch water column).¹
- A pressure indicator must be installed that can readily monitor room pressurization.¹ A notification system must be installed in each pressure monitor to alert pharmacy personnel when pressure differentials deviate from specifications.¹
- Water sources, sinks, and drains must not be located in a clean room but are permitted in the anteroom.¹
- Floors, walls, ceilings and all exposed surfaces in the controlled area must be smooth, impervious, non-friable, free from cracks and crevices, nonporous and resistant to damage from cleaning and disinfecting products.¹
- Cleaning must take place in the clean room at a time when no aseptic operations are in progress.¹²
- Essential furniture in the controlled area must be nonporous, smooth, non-shedding, impermeable, cleanable, and resistant to disinfectants.¹
- Shelves and supplies must be kept to a minimum in the clean room and anteroom to help limit the number of particulates.¹
- Each room in the controlled area must be identified with appropriate informative signs (e.g., pictograms identifying the hazard, the need for special care, restricted access, dress code).¹
- The door opening into the clean room and the door leading to the anteroom must not be opened at the same time in order to maintain pressure differential between the two rooms.¹
- Appropriate personal protective equipment (PPE) and clean room garb must be donned by all personnel prior to entering the controlled area to minimize the spread of skin particles that may shed.¹²
- Lab coats and isolation gowns must not be worn in the controlled area in place of chemotherapy gowns.⁴

- No shipping box or other external cartons may be taken into the controlled area.¹
- Hazardous drugs requiring manipulation other than counting or repackaging of final dosage forms must be stored separately from other inventory in a negative pressure room that maintains at least 12 ACPH.⁴
- Refrigerated hazardous drugs requiring manipulation other than counting or repackaging of final dosage forms must be stored in a dedicated refrigerator in a negative pressure room that maintains at least 12 air changes per hour.^{4,11}
- Refrigerator temperatures must maintain a temperature range of 2°C to 8°C.¹
- An air exhaust must be located behind any refrigerator that is inside the clean room to remove any particles generated by the unit.¹
- Hazardous drug spill kits must be available in or near the hazardous drug storage area.¹

**NAPRA Model Standards for Pharmacy Compounding of
Hazardous Sterile Preparations**

**Summary of Engineering Control Requirements for
Hazardous Drug Sterile Product Preparation and Storage Areas**

	ISO Classification	HVAC (Heating, Ventilation and Air Conditioning)					
		Temperature	Air Changes per Hour (ACPH)	Relative pressurization to adjacent area	Pressure monitoring device installed and monitored	Area exhaust ventilation	HEPA filtered supply air
HD Clean Room	ISO Class 7 (under dynamic conditions)	≤ 20° Celsius	≥ 30 ACPH	Negative ¹² (not less than 2.5 Pa [equivalent to 0.01-inch water column] negative pressure to adjacent anteroom)	Required	Required	Required
Anteroom (adjacent to HD clean room)	ISO Class 7 (under dynamic conditions)		≥ 30 ACPH	Positive ¹² (not less than 2.5 Pa [equivalent to 0.01-inch water column] positive pressure to adjacent clean room; not less than 5.0 Pa [equivalent to 0.02 inch water column] positive pressure to the rest of the pharmacy)	Required (between anteroom and the rest of the pharmacy)		Required
HD Storage Room		≤ 30° Celsius	≥ 12 ACPH	Negative (not less than 2.5 Pa [equivalent to 0.01-inch water column] negative pressure to adjacent area)	Required	Required	

C.3 Equipment and Furniture

All equipment and furniture brought into the controlled area must first be cleaned using a germicidal disinfectant detergent and then disinfected using sterile 70% isopropyl alcohol or a sporicidal before being placed into the controlled area. Equipment removed from corrugated cardboard must be disinfected using a sporicidal, rather than sterile 70% isopropyl alcohol. Equipment must be re-disinfected using sterile 70% isopropyl alcohol before being brought into the clean room.¹

Any equipment removed from the controlled area must be decontaminated first.¹

C.3.1 Containment Primary Engineering Control

WorkSafe BC Occupational Health and Safety (OH&S) [Regulation 6.53\(1\)](#) states:

“All mixing, preparation and priming of administration sets with a cytotoxic (hazardous) drug must be performed in one centralized area in a specially designated Class II Type B biological safety cabinet that:

- *is exhausted to the outside atmosphere in a manner that prevents recirculation into any work area;*
- *has exhaust and ventilation systems that remain in operation for a sufficient period of time to ensure that no contaminants escape from the biological safety cabinet into the workplace; and*
- *is equipped with a continuous monitoring device to permit confirmation of adequate airflow and cabinet performance”.*

C.3.1.1 Class I BSC

Class I BSCs are used when there is a need for containment, but not aseptic product protection and therefore must not be used for sterile hazardous drug preparation.^{7,13} A minimum Class I BSC that is located in a negative pressure room with at least 12 air changes per hour, and that fully exhausts to the outside environment must be used for manipulation of non-sterile hazardous drugs.⁴

C.3.1.2 Class II BSC

Because there is a possibility that HEPA-filtered air recirculated back into the clean room may be contaminated with hazardous drug, Class II Type A cabinets must not be used during preparation of hazardous drugs.³

A minimum Class II Type B BSC that is exhausted to the outside atmosphere with no recirculation into any work area must be used for the preparation of sterile hazardous drugs.³

C.3.1.3 HEPA Filter

HEPA filters must be present in C-PECs used for the preparation of hazardous drug sterile preparations. Air that flows towards the work surface inside the cabinet and air that is expelled out to the environment must first pass through at least one HEPA filter.³

C.3.1.4 Airflow

HEPA-filtered air inside the C-PEC must be supplied at a velocity sufficient to sweep particles away from the critical area and maintain unidirectional airflow during compounding.¹²

Manipulations must be performed in the direct compounding area, at least six inches in from the front opening of a biological safety cabinet (BSC),² behind the air ‘split’.¹⁴ Contaminated air must be able to escape via the rear grill, not via the front opening.¹³

In order for the BSC to help protect the operator, paths of airflow must remain clear.¹⁵

Note:

- Horizontal laminar airflow hoods must not be used for the preparation of hazardous drugs⁴

C.3.1.5 Ultraviolet Lights

The ultraviolet light may cause eye damage and must not be turned on when personnel are working in or near the C-PEC, or in the clean room.^{16,17}

C.3.1.6 Viewing Window

To protect the upper body and face from any splashes or aerosols produced inside a BSC; the viewing window must be kept at the manufacturers' recommended height during hazardous drug preparation.^{16,17}

C.3.1.7 Location

A biological safety cabinet used for hazardous drug compounding must be located away from doorways, traffic corridors, and air conditioning and heating vents.^{1,17}

C.3.1.8 Monitoring

The C-PEC used for hazardous drug sterile compounding must be operated continuously with the blower turned on 24 hours a day, seven days a week^{2,4} unless being serviced.² It must be equipped with a continuous monitoring device to allow confirmation of adequate airflow and cabinet performance.³

For the safety of the patient and the operator, hazardous drug compounding must not take place when a C-PEC alarm is sounding or the lights and/or gauges indicate the cabinet is not functioning within the manufacturer's specifications.¹⁷

Site specific procedures must be created and posted for workers so that when the gauges, lights or alarms indicate that the C-PEC is not working properly or there is a power interruption, the safety of personnel, the environment and the aseptic condition of the product (if possible) will be maintained.⁸

C.3.1.9 Testing and Certifying Biological Safety Cabinets

Testing and certifying the biological safety cabinet must be completed by a qualified person (e.g., a person who has been accredited by the National Sanitation Foundation [NSF] to perform testing of biological safety cabinets) when installed.¹⁸ Certification procedures used must meet the requirements of the NSF Standard 49- Biosafety Cabinetry: Design, Construction, Performance, and Field Certification (current version)^{1,18} The BSC must be re-certified every six months¹ and when the cabinet is altered or repaired or the HEPA filter is changed.^{1,17,18} Testing and certifying the biological safety cabinet must occur during dynamic operating conditions.¹

Prior to servicing a biological safety cabinet, service technicians or maintenance workers must be informed that the BSC may be contaminated with hazardous drugs.¹⁰ Appropriate personal protective equipment must be worn when testing, certifying or servicing the BSC.^{1,10}

After field certification, the BSC must have certification information posted on the front of the cabinet housing in a readily visible location.^{1,18}

Before and after testing and certifying of the BSC, all interior surfaces (including under the work surface) must be decontaminated, cleaned, and disinfected.¹

C.3.1.10 Replacing HEPA Filters

Only NSF certified technicians informed of the hazardous nature of the admixtures prepared in the biological safety cabinet shall replace HEPA and charcoal (if present) filters.¹⁰

Appropriate personal protective equipment must be worn when replacing HEPA filters and the contaminated filters must be handled and disposed of as hazardous waste.^{7,19}

C.3.1.11 Turning off a Containment Primary Engineering Control (C-PEC)

If it is necessary to turn off a C-PEC for testing and certifying or for maintenance, all surfaces inside the cabinet including under the work surface must be decontaminated, cleaned, and disinfected first.¹ If the decontaminating agent chosen contains a germicidal disinfectant detergent, then interior surfaces of the C-PEC may be decontaminated and cleaned in one step.

If the internal blower and external exhaust fan of a BSC are both turned off, the work-access opening and the HEPA exhaust area must be covered with impermeable plastic and sealed with tape to prevent any remaining hazardous drug contamination from inadvertently escaping from the BSC until maintenance work begins. The BSC must be sealed with plastic whenever it is moved or left inoperative for a period of time.¹⁴

C.3.1.12 Decontaminating, Cleaning, and Disinfecting Interior Surfaces of the Containment Primary Engineering Control (C-PEC)

To maintain an aseptic environment and to protect against possible contact with hazardous drug particles, interior surfaces of the C-PEC must be decontaminated, cleaned, and disinfected using a decontaminating agent, a germicidal disinfectant detergent, and sterile 70% isopropyl alcohol.¹

Prior to decontaminating and cleaning a C-PEC, proper hand hygiene procedures must be followed and full personal protective equipment (PPE) must be donned.¹

When cleaning, decontaminating, and disinfecting interior surfaces of the C-PEC (with or without raising the viewing window), additional PPE is required, including a NIOSH-approved elastomeric half face mask respirator (not a N95 respirator), a face-shield, and safety goggles or an elastomeric full face mask respirator. The respirator must be used with an appropriate filter cartridge and be fit-tested for the operator.¹ The compounder's head and upper body must remain outside of the C-PEC at all times; a cleaning tool may be used extend reach if necessary.¹

To protect others from potential exposure to hazardous drugs, pharmacy personnel who must be present in the clean room or in the area of the C-PEC must also wear a NIOSH-approved elastomeric half face mask respirator (with an appropriate filter cartridge) fit-tested for the operator (not a N95 respirator) in addition to all other PPE. If there is a risk of splash, other pharmacy personnel must also wear safety goggles and a face shield or an elastomeric full face mask respirator.¹

C.3.1.12a Decontaminating, Cleaning, and Disinfecting the Work Surface of the Containment Primary Engineering Control (C-PEC)

The work surface of the C-PEC must be decontaminated, cleaned, and disinfected using a decontaminating agent, a germicidal disinfectant detergent, and sterile 70% isopropyl alcohol:¹

- after completing each preparation
- before leaving the C-PEC for an extended period of time (e.g., for a break)
- upon returning to the C-PEC after an extended period of time
- after a minor spill involving the working surface

Alcohol must be allowed to dry before beginning the next sterile preparation.¹²

C.3.1.12b Daily Decontaminating, Cleaning, and Disinfecting Interior Surfaces of the Containment Primary Engineering Control (C-PEC)

Prior to commencing daily compounding, all interior surfaces of the C-PEC (except under the work surface) must be cleaned and disinfected using a germicidal disinfectant detergent followed by sterile 70% isopropyl alcohol.¹² If the viewing window has been raised during cleaning and disinfecting, it must be lowered to the manufacturers recommended operating level. The C-PEC must purge for at least fifteen minutes⁸ afterwards unless otherwise recommended by the manufacturer.

Following hazardous drug compounding, the C-PEC must purge for at least five minutes¹⁷ and then all interior surfaces (except under the work surface) must be decontaminated, cleaned, and disinfected using a decontaminating agent, a germicidal disinfectant detergent, and sterile 70% isopropyl alcohol:^{1,20}

- after preparations within the C-PEC are completed for the day²
- prior to compounding 'latex-free' preparations²
- prior to compounding sterile HD preparations in a C-PEC once it has been used to compound non-sterile HD preparations²
- prior to resuming compounding in a C-PEC that is turned off between aseptic processes for any reason (e.g., power interruption, maintenance)²

C.3.1.12c Weekly Decontaminating and Cleaning of all Interior Surfaces of the Containment Primary Engineering Control (C-PEC)

Decontaminating, cleaning, and disinfecting of all surfaces inside the C-PEC including under the work surface, must occur once a week, after a HD spill in the C-PEC, and before maintenance/ certification/servicing of the C-PEC.⁸ Disinfection of the C-PEC must be augmented with weekly use of a sporicidal agent.²¹

After decontamination and cleaning is complete, the viewing window is lowered to the manufacturers recommended operating level and the C-PEC must purge for at least thirty minutes (unless otherwise recommended by the manufacturer) prior to sterile compounding.⁷

C.3.2 Communication System

Verbal communication between personnel in the clean room and the anteroom or between personnel in the clean room and the general pharmacy must not be through open doors or pass-throughs to minimize the chance of introducing viable and non-viable contaminants into the rooms and the spread of hazardous drug contamination outside the rooms.¹

C.3.4 Chairs

Chairs used in the controlled area must be made of smooth, non-friable, non-porous, washable materials that are resistant to damage from cleaning and disinfecting products.¹

C.4 Decontaminating, Cleaning, Deactivating, and Disinfecting Surfaces in the Controlled Area

Housekeeping activities must not take place in the clean room⁴ or anteroom when compounding is occurring.

A germicidal disinfectant detergent must be used when cleaning surfaces and equipment in the controlled area.¹ A germicidal disinfectant detergent may be used to both clean and then disinfect surfaces.

Sterile 70% isopropyl alcohol must be used to disinfect items just prior to placement into the clean room.¹

Sterile 70% alcohol must be used for disinfecting items just prior to placement into the C-PEC and for disinfecting surfaces of the C-PEC following decontamination or cleaning.¹

Daily, weekly, and monthly decontaminating, cleaning, deactivating, and disinfecting procedures in the controlled area must be performed per Tables 1 to 3 (at a minimum).^{1,12}

The required contact time for the selected agents used must be adhered to.¹

Remove hazardous waste from controlled areas daily¹

- Tie the bag and/or seal the container before removing
- Do not compress contents as it may generate airborne HD particles
- Do not store new garbage bags in the bottom of the HD waste container

All decontaminating, cleaning, and disinfection activities must be recorded and retained in a general maintenance log.¹

Section D - Protective Measures

D.1 General

Personnel must follow all established procedures to minimize the release of particles into the aseptic preparation environment leading to possible contamination of the final product(s) and to decrease the possibility of occupational exposure to hazardous drugs.^{8,12}

There must be policies and procedures that address the safe and aseptic handling of hazardous drugs. There must be strict adherence to safe handling policies and procedures.¹

D.2 Personal Protective Equipment (PPE) and Clean Room Garb

Personal protective equipment and clean room garb must be provided to minimize or prevent healthcare workers exposure to hazardous drugs.³ All personnel entering the controlled areas must follow appropriate hand hygiene and garbing procedures.¹

If disposable PPE and/or clean room garb become contaminated or are suspected of being contaminated with HD, they must be removed and disposed of in a HD waste container¹ or laundered per site-specific policy.²²

D.2.1 Scrubs

Street clothes must be replaced with fresh scrubs daily by all personnel when the work assignment will take place in the controlled area.¹ Scrubs worn into the controlled area must not have been worn outdoors. Scrub bottoms must fully cover the legs including while seated.²

An isolation gown (enclosed at the neck and tied around the waist) or a buttoned lab coat must be worn over scrubs by staff that *will be* (re-)entering the controlled area.¹⁵

Scrubs must not be worn home to ensure that no HD contamination is transported home and to ensure that the process of cleaning the clothing does not introduce lint onto the low-lint scrubs.¹⁵

Scrubs contaminated with hazardous drug (e.g., as a result of a HD spill) must be isolated and placed into a separate laundry bag that is labelled as requiring special handling (e.g., 'Cytotoxic').²²

D.2.2 Footwear

Each facility must be in compliance with WorkSafe BC regulations to help reduce preventable injuries due to inappropriate footwear.⁶

WorkSafe BC Occupational Health and Safety (OH&S) [Regulation 8.22](#) states:

1. "A worker's footwear must be of a design, construction and material appropriate to the protection required and that allows the worker to safely perform the worker's work."
2. "To determine appropriate footwear under subsection (1), the following must be considered: slipping; tripping; uneven terrain; abrasion; ankle protection and foot support; potential for musculoskeletal injury; crushing potential; temperature extremes; corrosive substances; puncture hazards; electrical shock; any other recognizable hazard."

Personnel entering the controlled area must wear socks that are long enough to reach higher than the bottom of the pant legs at all times, including when seated, as well as closed shoes.¹

D.2.3 Shoe Covers

At least one pair of disposable shoe covers must be worn in the controlled area.² When the controlled area comprises of rooms in addition to the anteroom and clean room, this pair of shoe covers is donned in the gowning room.

Two pairs of disposable shoe covers (or a second pair if already wearing one pair of shoe covers) must be donned when stepping from the dirty side of the demarcation line in the anteroom to the clean side. Two pairs of disposable shoe covers must be worn in the clean room.¹ The outer pair of shoe covers must be removed with gloved hands when stepping out of the hazardous drug clean room to the clean side of the anteroom.⁴

The inner pair of shoe covers is removed when stepping from the clean side of the demarcation line in the anteroom to the dirty side.¹ In facilities that have incorporated a gowning room in the controlled area, the inner pair of shoe covers is removed in the gowning room, not in the anteroom.²

Shoe covers must be disposed of in hazardous waste containers and not saved for reuse.⁴

D.2.4 Hair Covers

A disposable hair cover (covering hair and ears completely) and beard cover (if necessary to cover facial hair or stubble) must be worn by all personnel working in the controlled area.¹

The hair cover (and beard cover if necessary) must be donned on the dirty side of the demarcation line in the anteroom.¹ In facilities that have incorporated a gowning room in the controlled area, the hair and beard covers must be donned in the gowning room.²

Hair and beard covers must be removed on the dirty side of the demarcation line in the anteroom.¹ In facilities that have incorporated a gowning room in the controlled area, the hair and beard covers are removed in the gowning room, not in the anteroom.²

Hair and beard covers must be disposed of in hazardous waste containers after each removal and not saved for reuse. Beard covers must be changed after 3.5 hours of continuous use.¹

D.2.5 Medical Masks

At a minimum, medical masks rated at least Level 1 by ASTM F2100²³ must be worn by all personnel present in a clean room, including personnel compounding in a PEC and in a C-PEC, unless decontaminating surfaces inside a C-PEC.^{1,21}

Medical masks do not provide respiratory protection against hazardous drug exposure and therefore must not be worn when respiratory protection from HD exposure is required.¹

Medical masks must be donned on the dirty side of the demarcation line in the anteroom and must cover from the bridge of the nose down to below the chin.¹⁵ Masks must not be saved for reuse. **Masks must be changed after no more than 3.5 hours of use.**¹

Masks worn in the hazardous drug clean room must be disposed of in a hazardous waste container.¹

D.2.6 Respirators

D.2.6.2 Chemical Cartridge Respirator

A chemical cartridge respirator with an appropriate filter cartridge must be worn when cleaning up HD spills, when decontaminating or cleaning the C-PEC with the viewing window raised, or when working in a clean room when a C-PEC is being decontaminated or cleaned.¹

A chemical cartridge respirator with an appropriate filter cartridge must be worn if unpacking HD shipments suspected of being damaged.¹

Personnel wearing a respirator that requires an effective seal with the face for proper functioning must be:⁶

- fit-tested prior to initial use;
- retested at least once a year, when there is a change in the respirator face piece, or when a user's physical condition changes affecting the fit; and
- clean shaven where the respirator seals with the face.

A respirator must not be worn over a medical mask.⁶

Other personal protective equipment that is to be worn at the same time as a respirator and which could interfere with the respirator fit must be worn during a fit test.⁶

WorkSafe BC Occupational Health and Safety (OH&S) [Regulation 8.41](#) states:

“Before each use of a respirator which requires an effective seal with the face for proper functioning, a worker must perform a positive or negative pressure user seal check in accordance with CSA Standard CAN/CSA-Z94.4-02, Selection, Use, and Care of Respirators.”

Disposable respirators must be disposed of in hazardous waste containers after each removal and not saved for reuse. **Disposable respirators must be changed after no more than 3.5 hours of use.**¹

Reusable respirators (including exterior surfaces of the filter cartridge casings) must be cleaned as per the manufacturer's recommendations daily after use. Filter cartridges must be changed per the maximum cumulative use time established by Occupational Health and Safety.¹

Used filters and the cartridge housing they are supplied in must be disposed of in hazardous waste containers when replaced.¹

D.2.7 Chemotherapy Gowns

To decrease particulate levels in the preparation area and to decrease the risk of direct skin contact with hazardous drugs, compounding and cleaning personnel must wear a non-linting, impermeable, disposable chemotherapy gown with long sleeves and fitted cuffs, enclosed at the neck, closing at the back (no open front), and tied around the waist.¹ Chemotherapy gowns must be worn for all activities that may result in the worker's direct exposure to hazardous drugs.¹

Chemotherapy gowns worn when mixing hazardous drugs in the C-PEC must be removed for disposal while still in the clean room to help prevent the spread of hazardous drug contamination to areas outside of the clean room.³

Personnel leaving the hazardous drug clean room to work in another room in the controlled area (e.g., anteroom, set-up room) or leaving the controlled area through a set-up and/or gowning room must wear an isolation gown.²

A chemotherapy gown must be worn if unpacking a suspected damaged hazardous shipment or cleaning up a hazardous drug spill.¹

Used chemotherapy gowns must be discarded into hazardous waste containers after:¹

- no more than 3 hours of continuous compounding
- contamination has occurred or is suspected
- each removal

D.2.8 Isolation Gowns

An isolation or chemotherapy gown must be worn by all personnel working in controlled areas.² Isolation gowns must be low-linting with long sleeves and fitted cuffs, enclosed at the neck, closing at the back (no open front), and tied around the waist.²

Isolation gowns or lab coats must not be worn in the hazardous drug clean room by personnel working in the C-PEC in place of chemotherapy gowns.⁴

An isolation gown (enclosed at the neck and tied around the waist) must be worn:

- by staff compounding non-sterile non-hazardous drugs outside of the controlled area²⁴
- by staff working in the controlled area unless wearing a chemotherapy gown²
- over scrubs by staff that *will be* (re-)entering the controlled area¹⁵

Disposable isolation gowns contaminated with hazardous drugs must be disposed of into a HD waste container. Reusable isolation gowns contaminated with hazardous drugs (e.g., as a result of a HD spill) must be placed into a separate laundry bag that is labelled as requiring special handling (e.g., 'Cytotoxic').²²

D.2.9 Chemotherapy Gloves

Gloves worn when touching surfaces that may be contaminated with hazardous drug must be tested with nine chemotherapy drugs as required in the American Society for Testing and Materials International (ASTM) Standard D6978-05 (*Standard Practice for Assessment of Resistance of Medical Gloves to Permeation by Chemotherapy Gloves*).²⁵

A report of the ASTM D6978-05 Standard test results indicating the minimum breakthrough detection time for each of the nine drugs tested must be provided to the facility by the glove manufacturer for each brand/type of chemotherapy glove to be worn by staff when handling hazardous drugs. The reported breakthrough detection times must be used to determine if the gloves are appropriate and the length of time that each brand and type of chemotherapy glove may be worn while staff handles hazardous drugs.^{4,25}

Two pairs of disposable chemotherapy gloves must be worn for all activities that may result in hazardous drug exposure including handling all hazardous drugs and hazardous drug waste.¹ Two pairs of disposable chemotherapy gloves must be worn at all times by all personnel working in the hazardous drug clean room.¹

Disposable chemotherapy gloves worn when mixing hazardous drugs in the C-PEC must be powder-free, sterile² and long enough to cover the cuff of the chemotherapy gown.¹ Two pairs of chemotherapy gloves must be donned by all personnel on the clean side of the demarcation line in the anteroom immediately after performing hand hygiene.¹ In facilities that have incorporated a gowning room in the controlled area, at least one pair of non-sterile gloves is donned by all personnel in the gowning room after washing hands.²

To minimize the spread of hazardous drug contamination outside of the clean room, outer chemotherapy gloves worn during hazardous drug compounding must not be worn outside of the clean room.¹

The inner pair of chemotherapy gloves worn during hazardous drug compounding must be removed on the clean side of the demarcation line in the anteroom immediately prior to washing hands. In facilities that have incorporated a gowning room in the controlled area, the inner pair of gloves worn during hazardous drug compounding must be removed in the anteroom (immediately prior to washing hands). A new pair of non-sterile chemotherapy gloves must be donned prior to leaving the anteroom. When leaving the controlled area, the new pair of gloves is removed in the gowning room immediately prior to washing hands.²

Hands must be washed with soap and water every time chemotherapy gloves are removed.⁴

Latex-free chemotherapy gloves must be made available to staff.³

Both pairs of disposable chemotherapy gloves worn when handling hazardous drugs must be changed every 30 minutes (unless otherwise indicated by the manufacturer's documentation) or immediately if a tear, puncture or contamination is known or suspected.⁴

Chemotherapy gloves must be disposed of in hazardous waste containers.⁴

D.2.10 Eye Protection

Safety goggles worn when working with cleaning agents or other liquids that pose a splash risk to the eyes must provide closed ventilation (non-ventilated) or indirect ventilation (preferred). Safety goggles must not provide direct ventilation.²⁶

Safety glasses provide impact protection but do not provide the same level of splash or droplet protection as safety goggles and therefore must not be used when there is a risk of chemical splashes and/or sprays to the eyes.²⁷

When working with cleaning agents or other liquids that pose a splash risk to the face and eyes, a face shield must be worn in addition to safety goggles, not as a substitute.²⁸

A face shield and safety goggles or a full face mask respirator must be worn for splash protection:¹

- when working at or above eye level
- when decontaminating or cleaning the containment primary engineering control
- when cleaning up a hazardous drug spill
- when performing manipulations with splash risk to the face and eyes
- when unpacking shipments suspected of containing damaged drugs

Safety goggles, face shields, and reusable respirators must be decontaminated using a decontaminating agent daily after use.⁴

If glasses are worn for vision correction, a face shield and safety goggles or a full face mask respirator must be worn over them whenever there is a risk of splash to the eyes when using cleaning agents or other liquids.⁴

Removal of a medical mask to don a respirator, or removal of a respirator to don a medical mask (e.g., to decontaminate the C-PEC or after the C-PEC has been decontaminated) must not occur in the clean room.²

D.3 Hand Hygiene

Hand hygiene must be performed by all personnel prior to entry into the clean room to minimize microbial contamination of sterile products. After handling hazardous drugs and removing chemotherapy gloves, hand washing is performed⁴ to remove possible drug contamination.

D.3.1.1 Alcohol-Based Hand Rub

Alcohol-based hand rubs used to disinfect hands before compounding parenteral hazardous drugs must have a minimum alcohol concentration of 70%, and be used in conjunction with plain or an antimicrobial soap.²⁹

D.3.2 Hand Washing After Handling Hazardous Drugs

Personnel handling hazardous drugs must wash their hands with soap and water immediately after removal of chemotherapy gloves.⁴

D.3.2.1 Compounding personnel

Personnel compounding sterile hazardous drug preparations must perform hand hygiene before donning two pairs of sterile chemotherapy gloves.² Prior to performing hand hygiene, all jewellery including bracelets, rings and watches must be removed to prevent material from being trapped around or underneath them.² Hands must be dried with a clean, low lint towel.¹²

D.3.3 Nails and Nail Polish

Wearing of artificial nails or other nail applications is prohibited while working in the controlled area. Natural nails must be kept neat and trimmed, and must be free of nail polish.¹

D.4 Safety Stations

Eyewash stations and emergency showers must be easily accessible and clearly identified by signs which indicate their location and provide clear directions for their use.⁵

Personnel that are required to use emergency eyewash and shower facilities must be adequately trained in their location and proper use.⁵

For potential exposure to high risk materials: *WorkSafe BC Occupational Health and Safety (OHS) [Regulation Table 5-3: Provision and location of emergency washing equipment](#) states:*

“Eye Equipment: Tempered continuous flow eyewash facility with a minimum duration of 15 minutes (or more if required by the nature of the material)

Location: Within 5 seconds walking distance of the hazard area, but no further than 6 meters (20 feet).

Skin Equipment: Tempered, continuous flow emergency shower facility with a minimum duration of 15 minutes (or more if required by the nature of the material)

Location: Same location criteria as for high risk eyewash facility except that the shower may be located further than 6 meters, and

- (a) a supplementary emergency washing facility such as a non-tempered drench hose is located within 5 seconds walking distance of the hazard area but no further than 6 meters, and*
- (b) a tempered shower facility is available within the building to start emergency washing within 5 minutes of the contact”*

D.4.1.2 Hand Held Portable

Portable eyewash stations must be capable of delivering a minimum flush duration of 15 minutes.⁵

D.4.2 Emergency Showers

Emergency showers must not be used to flush the user's eyes because the high rate or pressure of water flow could possibly damage the eye.³⁰

D.4.3 Safety Stations Maintenance

Plumbed emergency eyewash and shower facilities must be full flow tested at least once per month, for a sufficient length of time to completely flush the branch of the water line supplying the eyewash.⁵

Hand held portable eyewash equipment must be inspected and maintained according to the manufacturer's instructions.³⁰

Section E - Supplies and Devices

E.1 Supplies

E.1.3 Sterile Alcohol Swabs

Single use, individually packaged sterile 70% isopropyl alcohol swabs must be used to disinfect a critical site prior to accessing.¹² Gauze pads or other particle-generating material moistened with alcohol must not be used to disinfect critical sites of containers prior to accessing.¹²

E.2 Devices

Devices used in the safe and accurate reconstitution and withdrawal of hazardous drug in a vial must support minimizing the production and release of HD aerosols and vapours, maintaining the sterility of hazardous drugs, and preventing HD leaks/spills.²

Staff must be trained to use the proper aseptic technique required with each device utilized in the safe preparation of hazardous drugs.¹³

The following criteria may be considered when deciding which devices are most suitable for the preparation of hazardous drugs.

- Device must be approved for use with hazardous drugs by the manufacturer³¹
- Venting devices used during preparation of parenteral hazardous drugs must have filters^{7,19}
- Luer-lock fittings must be used for all hazardous drug connections made during manipulation and dispensing³

E.2.1 Syringes

An appropriate size syringe must be selected so that no more than three-quarters (75%) of the syringe's maximum calibrated volume is filled with hazardous drug solution at any time during the compounding process.²

A syringe must not be used more than five times for a single compounding procedure (e.g., reconstitution).¹³

E.2.2 Syringe Tip Caps

Care must be taken to avoid touch-contaminating the end of the multi-function tip cap that will be luer-locked to either the syringe or the chemotherapy dispensing pin (critical site).¹²

E.2.3 Needles

All parts of a needle are critical sites. Needles must be manipulated by handling their paper over-wrap and/or needle caps. Paper-covered needles must be unwrapped by peeling apart the sides of the package just enough to expose the needle's luer-lock hub. Airflow to the hub must be maintained as the needle is un-wrapped and luer-locked onto a syringe. The needle cap must be left in place until the needle is ready to be used.¹⁵

Aluminum-free needles and devices must be used in the preparation and administration of CISplatin, CARBOplatin and oxaliplatin.³²

Safety Engineered Needles (SEN) must not be used in the preparation of hazardous drugs. There is a risk that the HD will spray droplets off of the needle point when the SEN cap is engaged.³¹

E.2.4 Needle Caps

Placing the open end of the needle cap directly on the work surface of the C-PEC must be avoided.¹²

E.2.7 Filters

Solutions prepared for parenteral administration must be filtered when there is a possibility that glass particles¹³ or particulate matter (e.g., core from a vial stopper) may be present and the solution is filterable.

E.2.7.1 Filter Needles

The same filter needle must not be used for both withdrawing and expelling solution.^{15,33}

E.2.7.2 Filter Discs

A filter disc used for hazardous drugs must be equipped with proximal and distal luer-locking connections.³

E.2.8 Filter Venting Devices

Negative pressure technique must not be used for hazardous drug reconstitution or withdrawal if filter venting devices⁷ or closed system drug transfer devices^{7,10} are available.

E.2.8.1 Chemotherapy Dispensing Pins

Chemotherapy dispensing pins or similar devices with spikes must not be used with vials of TAXOL® since they can cause the stopper to collapse resulting in loss of the sterile integrity³⁴ and the possible release of hazardous drug.

Note:

- Chemotherapy dispensing pins must be inspected for cracks prior to use. A cracked chemotherapy dispensing pin must be replaced prior to manipulation of HD solution³¹
- Chemotherapy dispensing pins must be disposed of in a HD sharps waste container if removed from a HD vial³¹
- A new chemotherapy dispensing pin must be used for each vial. Spraying of the solution or touch contamination can occur upon removal of the pin³⁵

E.2.8.2 Chemotherapy Vents

A new chemotherapy vent must be inserted prior to removal of a plugged chemotherapy vent.¹⁴

A hazardous drug vial stopper must be disinfected with sterile 70% alcohol prior to each puncture when multiple punctures are necessary.¹²

E.2.9 Syringe Fluid Dispensing Connectors/Syringe Tip Connectors

Both ends of the individually packaged fluid dispensing connector used with hazardous drugs must have luer-lock connections³ which allow transfer of solution from one syringe to another without leakage.

E.2.14 Closed System Drug Transfer Devices

Closed System Drug Transfer Devices must be used within the ISO Class 5 environment of a C-PEC during hazardous drug preparation.⁴ Protective clothing must be worn and best practice safety measures must be adhered to when using a Closed System Drug Transfer Device to prepare, administer and dispose of hazardous drugs.¹⁰

E.3 Containers

The length of time between opening an ampoule and transferring the solution into a closed-system (e.g., syringe) must be minimized.²

The neck of the ampoule must be wiped to disinfect using a sterile 70% alcohol swab before breaking and must not be touch-contaminated after being disinfected.¹²

Glass particles in solutions must be filtered prior to administration³⁶ unless the manufacturer indicates the solution cannot be filtered. Solution must not be withdrawn and injected using the same filtration equipment.²

All parts of an opened ampoule must be discarded into a sharps container.¹³

E.3.2 Vials

Removal of a flip top cap from a hazardous drug vial must be performed carefully inside the C-PEC to 'contain' and avoid spreading HD contamination to areas outside of the C-PEC.¹⁴

Hazardous drug vials must be wiped to disinfect (not sprayed) using a low-lint towel or gauze moistened with sterile 70% alcohol prior to placement inside the C-PEC.¹²

The date and time of puncture or the beyond use date and time must be written directly onto reconstituted and partial vials that will be saved for future use with ink that will not smudge or wipe off.²

E.3.4 Non-Di(2-ethylhexyl)phthalate (Non-DEHP) Bags

PACLitaxel, DOCEtaxel, temsirolimus, teniposide, etoposide, cabazitaxel, cycloSPORINE and ixabepilone must be prepared in non-DEHP containers and administered using non-DEHP tubing.³²

E.3.5 Empty Sterile Infusion Bags

Infusion bags used for hazardous drug solution waste must be disposed of as hazardous drug waste.⁴

E.4 Ambulatory Drug Delivery Infusion Devices

E.4.1.2 Infusor™ Flow Rates

The correct size of elastomeric Infusor™ with the correct infusion rate must be selected when preparing hazardous drug medication.³¹

To decrease the risk of accidental exposure to hazardous drug, the delivery tubing of the Infusor™ must be primed with hazardous drug-free solution.^{3,37}

E.4.1.4 Hazardous Drug Medication Infusion Device Labels

The intended infusion rate must be stated in millilitres per hour (mL/hour) on the medication label when hazardous drug is administered via an infusion device.³⁸

E.4.2 Computerized Ambulatory Drug Delivery (CADD®) Pump and Medication Cassette Reservoir

To decrease the risk of exposure to hazardous drug, the tubing of a CADD® medication cassette reservoir must be primed with hazardous drug-free solution.³

Section F - Aseptic and Protective Processes

F.1 Operational Standards for Sterile Hazardous Drug Preparation

Hazardous drugs shall be prepared only under conditions that protect the healthcare workers and other personnel in the controlled area.⁴

Operational standards must be adhered to when preparing sterile HD admixtures.²

F.1.1 Personnel Hygiene in the Controlled Area

- Eating, drinking, smoking, chewing gum **or candy**, or storing food in the controlled area is strictly prohibited^{1,3}
- Personnel with rashes, burns to the skin (including sunburn), weeping sores, cold sores, **other fresh wounds**, conjunctivitis, active respiratory infection **with coughing, sneezing or runny nose**, and wearing cosmetics are **prohibited from entering the controlled area**¹
- Before entering the controlled area, personnel must remove:
 - ✓ personal outer garments (e.g., bandana, coat, hat, jacket, scarf, sweater, vest, **boots, and outdoor shoes**) because they shed flakes and particles¹
 - ✓ jewellery, studs, and other accessories from fingers, wrists, forearms, face including nose, tongue and ears, and neck¹
 - ✓ all cosmetics, false eyelashes, perfume, hair products such as hairspray, henna tattoos and paper tattoos, as these products can generate particles that are possible sources of contamination¹
 - ✓ nail polish and other nail applications (nail extensions, synthetic nail-lengthening products), natural nails must be kept neat and trimmed¹

F.1.2 Personal Protective Equipment and Clean Room Garb in the Controlled Area

- **Street clothes must be replaced with fresh scrubs daily** by all personnel when the work assignment will take place in the controlled area¹
 - Personnel entering the controlled area must wear socks (that are long enough to reach higher than the bottom of the pant legs at all times, including when seated) and closed shoes¹
 - One pair of shoe covers must be donned in the gowning room to enter the controlled area,² a second pair of shoe covers must be donned if stepping from the dirty side of the anteroom to the clean side¹
- OR
- **Two pairs of disposable shoe covers (or a second pair if already wearing one pair of shoe covers) must be donned when stepping from the dirty side of the demarcation line in the anteroom to the clean side**¹

- Hair covers (covering hair and ears completely) and beard covers (as applicable), must be worn in the controlled area¹
- Hair covers and beard covers (as applicable) must be donned on the dirty side of the demarcation line in the anteroom¹
- Or
- Hair covers and beard covers (as applicable), must be donned in the gowning room²
- At a minimum, an AAMI rated Level 1 medical mask must be worn by personnel compounding hazardous sterile preparations in a containment primary engineering control^{1,39}
- When decontaminating interior surfaces of the C-PEC, when working in a clean room when a C-PEC is being decontaminated, or when cleaning up hazardous drug spills, additional PPE is required, including a NIOSH-approved elastomeric half face mask respirator with a face-shield and safety goggles or an elastomeric full face mask respirator. The respirator must be used with appropriate filter cartridges and be fit-tested for the operator¹
- A respirator must be donned on the dirty side of the demarcation line in the anteroom¹
- An isolation gown must be donned in the gowning room to enter the controlled area² when the controlled area comprises of rooms in addition to the anteroom and clean room
- A chemotherapy or isolation gown must be donned on the clean side of the demarcation line in the anteroom (depending on the work being performed)²
- A chemotherapy gown must be worn any time there is a risk of hazardous drug exposure (e.g., when working in the C-PEC, when cleaning up a HD spill)¹
- Hands must be cleansed and gloves donned in the gowning room²
- Hand hygiene must be performed and gloves donned on the clean side of the demarcation line in the anteroom to enter the clean room¹
- Gloves worn when touching surfaces that may be contaminated with hazardous drug must be tested with nine chemotherapy drugs as required in the ASTM Standard D6978-05. The reported breakthrough detection times must be used to determine if the gloves are appropriate and the length of time that each brand and type of chemotherapy glove may be worn while staff handles hazardous drugs^{4,25}
- Two pairs of disposable chemotherapy gloves must be worn at all times by all personnel when hazardous drug exposure is possible (e.g., when handling hazardous drug vials, when mixing hazardous drug in a C-PEC). Both pairs of chemotherapy gloves must be inspected for visible defects. Gloves that have visible defects must not be worn.¹
- Gloves must be powder-free because powder can contaminate the work area and can absorb and retain hazardous drug.⁴
- Gloves must be disinfected before being placed into the C-PEC by wiping with a low lint towel moistened with sterile 70% isopropyl alcohol. The gloves must be completely dry before performing aseptic compounding activities inside the C-PEC¹
- Both pairs of disposable chemotherapy gloves worn when handling hazardous drugs must be changed every 30 minutes (unless otherwise indicated by the manufacturer's documentation) or immediately if a tear, puncture or contamination is known or suspected.⁴
- Hands must be washed with soap and water every time gloves are removed.⁴
- Chemotherapy gowns worn when preparing hazardous drugs must be non-linting, impermeable, disposable chemotherapy gown with long sleeves and fitted cuffs, enclosed at the neck, closing at the back (no open front), and tied around the waist¹
- Lab coats and isolation gowns must not be worn in place of chemotherapy gowns when protection from HD exposure is required because they permit the permeation of hazardous drug and can hold spilled hazardous drug against the skin, thereby increasing exposure⁴

F.1.3 Containment Primary Engineering Control (C-PEC)

- The UV light inside the C-PEC may cause eye damage and must not be turned on when personnel are working in the clean room^{16,17}
- All interior surfaces of the C-PEC (except under the work surface) must be decontaminated, cleaned, and disinfected using appropriate agents followed by sterile 70% isopropyl alcohol¹² prior to commencing daily compounding. If the viewing window has been raised during cleaning and disinfecting, it must be lowered

to the manufacturers recommended operating level and the C-PEC must purge for at least 15 minutes after decontaminating before beginning compounding.⁸

- The viewing window must be kept at the manufacturer's recommended level during HD preparation^{16,17}
- Rapid arm movements that could disrupt the air curtain must be minimized^{16,17}
- The front air intake grill and the rear air exhaust route must not be blocked^{16,17}
- Manipulations must be performed at least six inches in from the front opening and side walls of the BSC²
- The work surface of the C-PEC must be decontaminated, cleaned, and disinfected between each preparation using appropriate agents followed by sterile 70% isopropyl alcohol²
- Following hazardous drug compounding, the C-PEC must purge for at least five minutes¹⁷ and then all interior surfaces (except under the work surface) must be decontaminated, cleaned, and disinfected using appropriate agents followed by sterile 70% isopropyl alcohol:¹²
 - ✓ after preparations within the C-PEC are completed for the day²
 - ✓ prior to compounding 'latex-free' preparations²
 - ✓ prior to compounding sterile HD preparations in a C-PEC once it has been used to compound non-sterile HD preparations²
 - ✓ prior to resuming compounding in a C-PEC that is turned off between aseptic processes for any reason (e.g., power interruption, maintenance)²

F.1.4 General Procedures

- Unnecessary items must not be taken into the C-PEC since airflow is disrupted in an overcrowded C-PEC¹⁶
- HD vials must be wiped with low-lint towels or gauze moistened with sterile 70% isopropyl alcohol to disinfect and physically remove HD contamination prior to placement inside the C-PEC¹²
- Prior to placement inside the C-PEC, the outer wrapping of unopened supplies (e.g., syringes) must be disinfected by wiping using a low lint towel moistened with sterile 70% isopropyl alcohol¹²
- Best practice standards for aseptic technique in vertical airflow must be adhered to when preparing sterile hazardous drug admixtures¹
- Compounding must occur in the direct compounding area (work zone) of the C-PEC such that critical sites are exposed to first air.¹ Supplies not immediately required for use must not be kept in the critical area of the C-PEC
- To decrease particle generation inside the C-PEC, paper coverings must be peeled away from needle hubs (critical sites) rather than pushing them through²
- Critical sites must be protected as soon as possible after being exposed and must not be touch contaminated¹²
- Infusion solution bag ports and vial stoppers must be disinfected using sterile 70% isopropyl alcohol prior to accessing¹²
- A new sterile alcohol swab must be used to disinfect each critical site¹²
- When reconstituting, the drug must be completely dissolved before withdrawing a dose⁸ or storing for future use
- Syringes must not be overfilled with hazardous drug. In most cases, syringes should not be more than three-quarters (75%) full, although some preparations require accurate volume measurements that necessitate the use of a smaller volume syringe²
- Each syringe barrel must be marked with a line indicating the volume of solution withdrawn and the name of the solution using a thin-tipped permanent marker. The line indicating the volume withdrawn into the syringe must be marked when the solution is in the syringe. Solution volumes that equal numbers marked on the syringe barrel by the manufacturer must also be marked with a line, not by circling the number at the calibrated volume³¹
- Negative pressure technique must not be used for hazardous drug reconstitution or withdrawal if filter venting devices² or closed system drug transfer devices¹⁰ are available
- A puncture-proof sharps container must be used for disposal of all sharp objects including needles, chemotherapy dispensing pins, and chemotherapy vents²

- All non-sharp waste generated during compounding of hazardous drugs must be placed inside a HD waste container (e.g., zip lock bag or sharps container) in the C-PEC for later removal and disposal⁴

F.1.5 Removing Products from the Containment Primary Engineering Control (C-PEC)

- Infusion solution bag ports that have been accessed must be wiped with an alcohol swab prior to removal from the C-PEC¹² to remove possible HD residue
- Infusion solution bags that have had hazardous drug added must be checked for leaks and particulate prior to removal from the C-PEC (if the injection port was used to add drug to the infusion solution bag)¹
- Outer chemotherapy gloves worn when compounding hazardous drugs must be removed, discarded within the C-PEC and replaced with a new pair of sterile chemotherapy gloves or wiped with a new towel moistened with a decontaminating agent prior to touching items for removal from the C-PEC²
- Surfaces of final preparation(s) may be contaminated with HD and must be decontaminated using a new towel moistened with a decontaminating agent prior to removal from the C-PEC²
- The final preparation must be labelled immediately after it is removed from the C-PEC with the patient-specific label and any required warning labels⁴⁰
- To remove a vial of HD that will be saved for reuse from the C-PEC:
 - ✓ the vial stopper must be wiped with a sterile 70% alcohol swab¹² to remove possible HD residue (if there is not a chemotherapy dispensing pin or CSTD inserted)
 - ✓ the puncture date and time or beyond use date and time must be written directly on the vial with a thin-tipped permanent marker²
 - ✓ the vial must be wiped with a new towel moistened with a decontaminating agent²
 - ✓ the vial must be placed inside a zip lock bag that is sealed inside the C-PEC or above the front grill upon removal from the C-PEC³³
- Containers used for HD waste (sharp and non-sharp) must be sealed and decontaminated using a new towel moistened with a decontaminating agent inside the C-PEC before removal from the cabinet¹⁰

F.1.6 Warning Labels

- All hazardous drugs and hazardous drug preparations must be easily identifiable by personnel involved in their handling³
- The container of hazardous drug must be appropriately labelled³ indicating the contents are hazardous in nature

F.1.7 Exiting the Clean Room

- PPE must be appropriately removed upon exiting the controlled area:^{3,4}
 - ✓ Outer chemotherapy gloves must be discarded into a hazardous waste container⁴ (inside or outside of the C-PEC) prior to exiting the clean room; outer gloves must NOT be worn outside the clean room once compounding hazardous drugs in the C-PEC has occurred⁶
 - ✓ The inner gloves must not be removed inside the clean room; inner gloves must be disposed of into a hazardous waste container¹
 - ✓ Both pairs of chemotherapy gloves worn when handling hazardous drugs must be changed every 30 minutes (unless otherwise indicated by the manufacturer's documentation) or immediately if a tear, puncture, or contaminates is known or suspected. Gloves must be disposed of into a hazardous waste container⁴
 - ✓ Removal and disposal of chemotherapy gowns must be done with care to avoid spreading HD contamination to other non-contaminated garments⁴
 - ✓ Chemotherapy gowns must be discarded into hazardous waste containers after:¹
 - no more than 3 hours of continuous compounding
 - contamination has occurred or is suspected
 - each removal
 - ✓ Chemotherapy gowns worn in the clean room during hazardous drug compounding must be removed and disposed of in the clean room, to prevent the spread of HD contamination from one area to another.³

- ✓ The outer pair of shoe covers must be removed with gloved hands upon exiting the clean room into the anteroom and must be discarded as hazardous waste⁴ **inside the clean room**
- ✓ The inner pair of shoe covers is removed upon stepping from the clean side of the demarcation line in the anteroom to the dirty side¹ and is disposed of into HD waste⁴
- Or
- ✓ The inner pair of shoe covers is removed in the gowning room² and is disposed of into a hazardous waste container⁴
- ✓ The medical mask or respirator and hair cover(s) must not be removed inside the clean room¹
- ✓ Disposable masks, respirators and hair cover(s) must be disposed of into a hazardous waste container¹
- ✓ **Disposable masks, respirators and beard covers must be changed after no more than 3.5 hours of continuous use¹**
- ✓ Reusable respirators, safety goggles, **and face shields** must be decontaminated daily after use¹
- ✓ Hands must be washed immediately with soap and water every time gloves are removed⁴
- ✓ An isolation gown must be donned over scrubs when exiting the anteroom¹³ into another room within the controlled area. If the anteroom exits directly into the general pharmacy, or to leave the controlled area, an isolation gown or lab coat must be donned over scrubs upon exiting¹³

F.2 Aseptic/Protective Routines

F.2.1 Critical Sites

Critical sites must be protected as much as possible and must not be touch-contaminated.¹² Protection of critical sites by precluding physical contact and airborne contamination must be given the highest priority in aseptic compounding practice.¹²

F.2.2 First Air

While working in the C-PEC, a path of first air must be maintained to critical sites at all times. It is vital to avoid reaching over or working directly above or in front of exposed or previously disinfected critical sites.¹

F.2.3 Disinfecting Critical Sites

The stopper on a vial or the port on an infusion solution bag must be disinfected using a sterile 70% alcohol swab just prior to penetration. At least 10 seconds must be allowed for the alcohol to dry (act) before manipulations begin.¹²

A new sterile swab must be used to disinfect each new surface.² The surface of sterile 70% alcohol swabs used to disinfect entry points on infusion solution bags and vials shall not contact any other object before contacting the surface of the entry point.¹²

Prior to removal from the C-PEC, the port of an infusion solution bag that has had drug added must be wiped with an alcohol swab¹² to remove possible HD residue.

F.2.4 Coring

Each vial and final product must be checked for particulate (e.g., coring) after each puncture of a vial stopper or infusion solution bag port.¹

F.2.5 Safely Capping Needles Used With Hazardous Drug

Needles are a critical site and therefore must be capped when not being used for injection or withdrawal.² Prior to manipulation of a hazardous drug-filled syringe, the needle must be capped to reduce aerosol release and prevent splashes from the needle tip.³³

For worker safety, two-handed recapping of a needle used for HD preparation is never an acceptable practice.²

F.3 Safe Handling Aseptic Techniques

F.3.1 Transfer of Hazardous Drug Solution from a Syringe

If too much hazardous drug solution has been drawn into a syringe, care must be taken to minimize aerosol and vapour production, and to contain hazardous drug solution while removing the excess volume.^{41,42}

Excess hazardous drug must NOT be ejected into the needle cap, sharps container, or any other open container⁴² as this could cause HD aerosolization, vaporization or contamination

F.3.2 Removal of Bubbles/Air from a Syringe

Bubbles and air must be removed carefully in a manner that prevents the release of HD solution and minimizes the production of HD aerosols in the C-PEC.^{41,42}

F.3.3 Attaching and Priming Solution / Secondary Administration Sets

Priming any intravenous administration set with hazardous drug solution in an uncontrolled environment must be avoided.³

To minimize exposure to HD, the administration tubing/line must be primed with HD-free solution whenever possible (e.g., unless contraindicated by the drug).³

Section G - Clean Up and Waste Disposal

G.1 Containment Primary Engineering Control (C-PEC) Waste Cleanup

The entire aseptic preparation area must be kept clean so that aseptically prepared products remain as free from potential microbial and hazardous drug contamination as possible.^{4,12}

G.2 Hazardous Waste Disposal

Hazardous waste containers must be available in all areas where hazardous drugs are received, stored, prepared and administered.³

All disposable items that may have come in contact with hazardous drugs during receipt, storage, preparation or administration must be treated as hazardous waste including PPE.³ Hazardous waste must be disposed of separately from general waste in hazardous waste containers with lids.² The hazardous waste container must be distinctly different from other types of waste containers.²

All disposable non-sharp HD waste must be disposed of in 4 mil thick plastic bags which are placed inside a rigid HD waste container or carton so that all waste is essentially 'double-bagged'.²

The warning label must identify the contents as hazardous so that individuals transporting the waste are alerted to the need for special handling.⁴

All sharps used for the preparation and administration of hazardous drug admixtures must be placed into a puncture-proof hazardous drug sharps container for disposal² without being crushed or clipped.^{13,41} Chemotherapy dispensing pins and chemotherapy vents removed from HD vials must also be disposed of in a hazardous drug sharps container.

The HD sharps container must be sealed when it is no more than three-quarters full or at the indicated maximum fill line.²

HD waste containers must not be overfilled and the contents must not be pushed down to make more room due to the risk of HD exposure.²

Two pairs of chemotherapy gloves must be worn while handling hazardous waste.¹⁰

While awaiting removal from the facility for disposal, hazardous waste must be stored in a secure area in securely sealed and properly labelled containers.¹³

Hazardous waste must be transported and disposed of according to Federal and Provincial regulations after leaving the facility.⁴

Section H - Safe Handling of Oral, Topical and Pre-Packaged Hazardous Drug Dosage Forms

All drugs listed on the facility's hazardous drug list must be handled according to the facility's hazardous drug safe handling guidelines.¹⁰ Oral, topical and pre-packaged hazardous drug dosage forms must be handled in a manner that prevents skin contact and minimizes the liberation of powdered or aerosolized HD into the air and cross contamination with other drugs.⁷

H.1 Oral Preparations

- Two pairs of chemotherapy gloves must be worn when handling hazardous drug tablets and capsules and when pouring HD oral solutions or suspensions⁴ in a designated area of the pharmacy dispensary^{4,3}
- All activities likely to result in particle generation, for example, weighing or mixing powder, crushing tablets/capsules, or filling capsules, must be performed in an externally vented, minimum Class I biological safety cabinet in a negative pressure room (with at least 12 APH) to minimize the risk of spreading HD contaminated particulate throughout the rest of the pharmacy⁴
- Counting of non-coated tablets or capsules that have visual evidence of HD powder residue on them or compounding HD oral solutions must be performed using containment strategies such as preparation inside an externally vented, minimum Class I biological safety cabinet⁴ to reduce the risk of HD exposure
- Dedicated 'chemotherapy' counting trays and spatulas must be used to count loose HD tablets and capsules^{2,4,3}
- "Chemotherapy" dedicated counting tray(s), spatula(s) and countertops should be decontaminated after each use using a decontaminating agent.² The towel used must be disposed of in HD waste⁴
- Hands must be washed with soap and water immediately after removing chemotherapy gloves⁴
- Gloves worn when handling hazardous drugs must be discarded in HD waste⁴
- Automated counting machines must not be used to count hazardous drug tablets and capsules⁴

H.2 Topical Preparations

- Two pairs of chemotherapy gloves must be worn when handling hazardous drug topical preparations that have been removed from the original packaging⁴
- Compounding hazardous topical products, especially activities likely to result in particle generation, must be performed in an externally vented minimum Class I biological safety cabinet⁴

All interior surfaces of a C-PEC (except under the work surface) used for both sterile and non-sterile HD preparations must be decontaminated, cleaned, and disinfected following non-sterile HD preparations using a decontaminating agent, a germicidal disinfectant detergent, and sterile 70% isopropyl alcohol. Once decontaminated, the C-PEC must purge for at least 15 minutes prior to compounding sterile HD products²

Section I - Hazardous Drug Spills

I.1 Hazardous Drug Spills

To minimize exposure of staff and patients to hazardous drugs, spills must be managed appropriately, according to established policies and procedures. Spill kits must be located in all areas where exposures may occur.³ These locations include hazardous drug preparation, dispensing, storage and receiving areas.

I.1.1 Recommended Spill Kit Contents

Disposable PPE and the filter cartridge⁴⁴ attached to the elastomeric respirator worn during a hazardous drug spill clean-up must be disposed of after use into a hazardous waste container. The respirator and other reusable PPE must be decontaminated using an appropriate decontaminating agent per the manufacturer's recommendation.¹

New employees must be advised of hazardous drug spill control procedures³ and be required to demonstrate competency in spill handling.¹⁹

Training and competency assessments should be documented.⁴

Section J - Accidental Exposure to Hazardous Drugs

J.1 Accidental Exposure to Hazardous Drugs

Healthcare workers must be made aware of how to manage accidental exposure to hazardous drugs.³

Any accidental hazardous drug exposure as a result of a spill, needle stick or other accident must be reported immediately to the professional practice leader/department manager⁴¹ and by calling the Provincial Workplace Health Call Centre reporting line at 1-866-922-9464. Appropriate documentation must be completed.²

J.1.2 Ingestion

Personnel must not take food, gum, drinks, cigarettes or personal medication into an area where hazardous drugs are handled³ (e.g., received, stored, prepared, administered and disposed).

Section K - Receipt, Storage, and Transportation of HD

K.1 Receipt and Unpacking

Personnel receiving and unpacking hazardous drugs must receive training to manage HD spills and be made aware of precautions and follow special handling procedures.⁴

Safe handling procedures must be followed to avoid breakage of hazardous drug containers, to minimize exposure to hazardous drugs, and to contain spills that occur when receiving and unpacking hazardous drugs within the pharmacy.²

K.1.1 Receipt

- Two pairs of chemotherapy gloves must be worn when packing and unpacking boxes containing hazardous drugs²
- The outside of cartons must be examined for possible damage prior to opening⁴
- Hazardous drugs requiring refrigeration must be unpacked and refrigerated immediately upon receipt^{8,15}

K.1.2 Receipt of a Damaged Shipment

Policies and procedures must be in place for handling damaged shipments of hazardous drugs.⁴

Damaged cartons, totes, and/or packages containing hazardous drugs that are received must NOT be opened and the receiver must don full PPE including a NIOSH-approved elastomeric half face mask respirator with a face-shield and safety goggles or a full face-piece respirator to handle the package¹ following site-specific procedures.

When cartons, totes, and/or packages are opened unknowingly with damaged contents inside, the receiver must don full PPE including a NIOSH-approved elastomeric half face mask respirator with a face-shield and safety goggles or a full face-piece respirator¹ and follow site-specific procedures.

HD spill kits with written procedures for use must be located in all areas where hazardous drugs are received.^{10,13}

K.2 Storage

- Containers, shelves and bins used for HD storage must be properly labelled with hazard warning labels identifying the drugs that require special handling³
- Barriers and other design features on bins and shelves must be present to contain accidental leakage and reduce the chance of drugs falling to the floor²
- Hazardous drugs must be stored in a manner that prevents spillage or breakage if the container falls⁴

- Hazardous drugs requiring manipulation other than counting or repackaging of final dosage forms must be stored separately from other inventory in a manner that prevents HD contamination and occupational exposure⁴
- Refrigerator temperatures must maintain a cold temperature range of 2°C to 8°C¹
- To prevent errors from occurring, medication that can be easily mistaken for another (sounds alike, looks alike, similar labelling) must be separated in all areas of the pharmacy²
- Access to areas where hazardous drugs are stored must be limited to authorized personnel²
- Hazardous drugs spill kits with written procedures for use must be readily available in areas where hazardous drugs are stored.³

K.3 Packaging and Transportation

Hazardous drugs must be packaged and transported in a manner which minimizes the risk of HD exposure due to a spill or breakage during transit.⁴

Pneumatic tubes or other mechanical transport systems that produce stress on the contents must not be used for hazardous drug transport.⁴

Module 2 – Pharmacy Medication Checks

Section A

A.1 Clinical Medication Order Check

The following must be reviewed and verified by a pharmacist for all parenteral, oral and topical oncology prescriptions prior to dispensing:⁴⁵

- pharmacy medication profile
- protocol code and cycle number (if appropriate)
- drug, dose (a maximum of a 5% variance is permitted in dose calculation unless a variance is prohibited by the treatment protocol), route, administration, timing and duration of each medication ordered; for cyclical therapies, no more than one cycle of medication will be dispensed at a time
- benefit status of the medication(s) and receipt of appropriate approval forms, if required
- patient-specific factors including allergies, alerts, and protocol required laboratory values; for new patients or patients beginning a new course of cancer treatment, baseline tests must have been conducted within four weeks of the start of therapy
- body surface area, calculated by the Mosteller equation, for the first treatment of each chemotherapy protocol only (based on actual body weight except for the designated high dose protocols described above, where ideal body weight is used) unless recalculated by the physician and documented in the patient's chart - if required for the cancer treatment being prescribed

A.2 Final Product and Computer Order Entry Check

The final product must be checked prior to dispensing:

- the volume/quantity calculation must be double checked, all components used must be appropriate and a visual inspection of the final product must be performed¹
- the drug volume(s) added to a final container must match the dose prescribed on the medication order⁷
 - reconstitution solution and volumes must be checked by any qualified person other than the individual who measured the solution^{8,46}

The computer order entry on the patient-specific label must be checked for accuracy.⁴⁵

The patient-specific label must be:

- affixed to the correct final product container;
- checked for completion; and
- checked that it accurately reflects what is written on the corresponding medication order.⁸

The final product must have the appropriate patient specific label, a HD warning label (if applicable) and any necessary auxiliary warning labels affixed to the container.^{7,46}

Section B

B.1 Documentation of Pharmacy Medication Checks

Standard operating procedures must be developed which include signed documentation that all the required pharmacy checks have been completed for each medication.⁷

- the pharmacist performing the clinical medication order review must indicate that the order is approved for preparation by documenting on the appropriate form prior to compounding⁴⁵
- the person performing the computer order entry (e.g., patient label) check must indicate the computer order entry is accurate as per the medication order by documenting on a permanent record⁷
- the person performing the final product check must indicate by documenting on a permanent record that they have performed a complete final product check⁷

Module 3 – Improving Patient Safety and Compliance

Section A

A.1 Intrathecal (IT) Doses

Preparation of IT doses must be performed accurately given the limitations of syringe technology while following the principles of aseptic technique for the safety of the patient and to minimize exposure of staff to the hazardous drugs.³¹

All intrathecal doses will have a label stating the patient name, date, generic drug name, dose and route in full (i.e. By INTRATHECAL Injection). The route will be highlighted by using all upper case letters and/or a coloured highlighting pen. This labelling will be attached to the syringe and to the outer zip-lock bag. Intrathecal doses will be provided in luer lock syringes (WorkSafe BC Health and Safety Regulation 6.54). Intrathecal syringes and labels will have an auxiliary label stating 'IT' attached to the syringe and the outer zip-lock bag. These will be bright in colour and clearly visible.

Intrathecal doses will be packaged in separate outer containers for transport to the centre specific treatment location (i.e. intrathecal doses will be delivered separately from all other chemotherapy doses).

For protocols in which intrathecal drugs and drugs administered by other parenteral routes are all to be given in one same treatment cycle, the non-intrathecal drugs will NOT be released from pharmacy until the nurse or physician confirms that the intrathecal drug administration is completed.

Note:

- All drug and diluent used in the preparation of IT doses MUST be preservative free³²
- All drug used MUST be labelled 'for intrathecal use' by the manufacturer³²
- Gloves MUST be changed immediately prior to preparing the intrathecal dose³¹

Section B

B.1 Vinca Alkaloids

All Vinca Alkaloids must be dispensed in a 50 mL minibag to be given over 5-15 minutes and labelled with an auxiliary label with the words 'WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES'⁴⁷

Section D

D.1 Outpatient Hazardous Drug Prescription Labelling

All hazardous drugs dispensed for outpatient use must be labelled according to the current labelling standards found in the Health Professions Act (HPA) Bylaws; College of Pharmacists of BC.

A recognized strategy to improve comprehension of the directions on outpatient prescription labels and to reduce inadvertent medication errors is to standardize drug labelling practices.⁴⁸

- All boxes/bottles must be individually labelled; multiple boxes must not be affixed together with a label on a single box⁴⁹
- Vials, blister cards, boxes, bottles and jars containing hazardous drugs must be clearly labelled with a “Chemotherapy” warning auxiliary label^{19,49}

Section E

E.1 Latex Allergies

Medication for patients allergic to latex must be prepared and administered in a latex-free environment. All surfaces of the biological safety cabinet must be decontaminated, cleaned and disinfected prior to compounding hazardous drug (HD) admixtures for latex-sensitive patients.⁵⁰

References

1. National Association of Pharmacy Regulatory Authorities, (NAPRA). Model Standards for Pharmacy Compounding of Hazardous Sterile Preparations. 2016. Available at: https://napra.ca/sites/default/files/2017-09/Mdl_Stnds_Pharmacy_Compounding_Hazardous_Sterile_Preparations_Nov2016_Revised_b.pdf. Accessed January 20, 2020
2. Canadian Society of Hospital Pharmacists. Compounding: Guidelines for Pharmacies. 1st ed. : CSHP; 2014
3. WorkSafe BC Publications. OHS Regulation Part 6 Substance Specific Requirements - Cytotoxic Drugs. 2006. Available at: <http://www2.worksafebc.com/publications/OHSRegulation/Part6.asp#top>. Accessed Oct. 9, 2007
4. United States Pharmacopeia. Hazardous Drugs - Handling in Healthcare Settings. USP Compounding Compendium. USP 39 - NF 34 ed. Rockville, MD: The United States Pharmacopeial Convention; 2016. p. 85-103
5. WorkSafe BC Publications. OHS Regulation Part 5 Chemical Agents and Biological Agents. Available at: <http://www2.worksafebc.com/publications/OHSRegulation/Part5.asp#SectionNumber:5.85>. Accessed July 21, 2010
6. WorkSafe BC Publications. OHS Regulation Part 8 Personal Protective Clothing and Equipment. Available at: <http://www2.worksafebc.com/publications/OHSRegulation/Part8.asp>. Accessed November 12, 2008
7. International Society of Oncology Pharmacy Practitioners Standards Committee. ISOPP Standards of Practice. Safe Handling of Cytotoxics. J.Oncol.Pharm.Pract. 2007;13 Suppl:1-81
8. CPhO. Standards of Practice for Oncology Pharmacy in Canada Version 2. 2009. Available at: <http://www.capho.org/standards-practice>. Accessed March 4, 2010
9. United States Pharmacopeia. Pharmaceutical Compounding - Sterile Preparations; 2nd Supplement. The United States Pharmacopeial Convention 2019:1-37
10. National Institute for Occupational Safety and Health, (NIOSH). NIOSH Alert Preventing Occupational Exposure to Antineoplastic and Other Hazardous Drugs in Health Care Settings. 2004;2004-165
11. BC Sterile Compounding Standards Working Group. Hazardous Drug Storage Interpretation Document. British Columbia 2021:1-11
12. United States Pharmacopeia. Pharmaceutical Compounding - Sterile Preparations. USP Compounding Compendium. USP 39 - NF 34 ed. Rockville, MD: USP Convention; 2016. p. 39-84
13. Power L, Jorgenson J. Safe Handling of Hazardous Drugs - Video Training Program. Bethesda, MD: ASHP; 2006
14. ACB Department of Pharmacy, Cross Cancer Institute, Tom Baker Cancer Centre editors. Alberta Cancer Board Pharmacy - Training Manual for the Preparation of Parenteral Cytotoxic Admixtures. 8th Edition ed. Alberta: Alberta Cancer Board; June2005
15. American Society of Health-System Pharmacists. Basics of Aseptic Compounding Technique - Video Training Program. Bethesda, MD: ASHP; 2006
16. Casey L, Wilson D. Primary Containment for Biohazards: Selection, Installation and Use of Biological Safety Cabinets 3rd Edition. 3rd Edition ed. : Department of Health & Human Services and National Institutes of Health; 2007. p. 291-329
17. Use of Biological Safety Cabinets. 2006. Available at: <http://www.uiowa.edu/~hpo/biosafety/BSC.htm>. Accessed April 16, 2008

18. WorkSafeBC Publications. OHS Regulation Part 30 Biological Safety Cabinets. 2008. Available at: <http://www2.worksafebc.com/publications/OHSRegulation/Part30.asp#SectionNumber:30.12>. Accessed April 7, 2010
19. ASHP Council on Professional Affairs. ASHP Guidelines on Handling Hazardous Drugs. American Journal of Health-System Pharmacy 2006;63(12):1172-1191
20. CriticalPoint. Decontamination, Cleaning and Disinfection in Hazardous Drug Sterile Compounding Environments. CriticalPoint 2017
21. National Association of Pharmacy Regulatory Authorities, (NAPRA). Model Standards for Pharmacy Compounding of Non-Hazardous Sterile Preparations. 2016. Available at: https://napra.ca/sites/default/files/2017-09/Mdl_Stnds_Pharmacy_Compounding_NonHazardous_Sterile_Preparations_Nov2016_Revised_b.pdf. Accessed January 20, 2020
22. British Columbia Occupational Health and Safety and Pharmacy Professionals. Hazardous Drug Exposure Control Program Draft. 2017. Available at: <http://2pod.phsa.ca/quality-safety/safety/exposure-prevention/drugs/Documents/Hazardous%20Drugs%20ECP.pdf>. Accessed January 20, 2020
23. Eric Kastango, Kate Douglass and Abby Roth. (2019). CriticalPoint Peer Network.
24. National Association of Pharmacy Regulatory Authorities, (NAPRA). Model Standards for Pharmacy Compounding of Non-Sterile Preparations. 2018. Available at: https://napra.ca/sites/default/files/documents/Mdl_Stnds_Pharmacy_Compounding_Nonsterile_Preparations_March2018_FINAL.pdf. Accessed January 20, 2020
25. American Society for Testing and Materials, (ASTM). Standard Practice for Assessment of Resistance of Medical Gloves to Permeation by Chemotherapy Drugs. 2005;6978-05
26. Albany County Fasteners. Indirect vs. Direct Ventilation Safety Goggles - When to Use Each. 2021. Available at: <https://www.albanycountyfasteners.com/blog/indirect-vs-direct-ventilation-safety-goggles/>. Accessed April, 2021
27. National Institute for Occupational Safety & Health (NIOSH). Eye Safety Tool Box Talk - Instructor's Guide. 2013. Available at: <https://www.cdc.gov/niosh/topics/eye/toolbox-eye.html>. Accessed April, 2021
28. International Safety Equipment Association. American National Standard Z87.1-2010 for Occupational and Educational Personal Eye and Face Protection Devices. 2010th ed. Arlington, Virginia: International Safety Equipment Association; April 13, 2010
29. British Columbia Provincial Hand Hygiene Working Group. Best Practices for Hand Hygiene in All Healthcare Settings and Programs. 2012;Appendix 1 - 923454
30. WorkSafe BC Publications. OHS Answers: Emergency Showers and Eyewash Stations. 2005. Available at: http://www.ccohs.ca/oshanswers/safety_haz/emer_showers.html. Accessed June 3, 2008
31. BCCA Division of Pharmacy. BCCA Best Practice Safe Handling Standards. 2008
32. BCCA Division of Pharmacy. BCCA Cancer Drug Manual. BCCA, 2007. Available at: <http://www.bccancer.bc.ca/HPI/DrugDatabase/default.htm>. Accessed June 4, 2008
33. Chow L, Moore M, BCCA Division of Pharmacy. Aseptic and Safe Preparation of Cytotoxic Drugs - Procedures and Checklists. Vancouver, BC: BC Cancer Agency; August 1995
34. Bristol-Myers Squibb. Taxol® (paclitaxel) Injection Package Insert. 2000. Available at: <http://www.fda.gov/cder/foi/label/2002/20262slr038l1bl.pdf>. Accessed June 4, 2008
35. Kendall-LTP. Vial Venting System. 2009. Available at: <http://www.kendall-ltp.com/Kendall-LTP/pageBuilder.aspx?webPageID=0&topicID=147871&xsl=xsl/productPagePrint.xsl>. Accessed Oct 29, 2009

36. Kempen PM, Sulkowski E, Sawyer RA. Glass ampules and associated hazards. *Crit.Care Med.* 1989;17(8):812-813
37. Baxter. Baxter Elastomeric Pumps - Clinician Guide Promoting Patient Mobility. Baxter, 2010. Available at: <http://www.capca.ca/wp-content/uploads/Baxter-Elastomeric-Pumps-Clinician-Guide1.pdf>. Accessed December, 2010
38. BCCA Division of Pharmacy. BCCA Systemic Therapy Update - Elastomeric Infusors. 2008;2(11)
39. PriMed. Mask Protection Standards & Medical Face Mask Information for Use. Available at: <https://www.primed.ca/resources/astm-mask-protection-standards/>. Accessed May, 2021
40. ASSTSAS Quebec. Prevention Guide Safe Handling of Hazardous Drugs. IRSST, 2008. Available at: <http://www.irsst.qc.ca/files/documents/PubIRSST/CG-002.pdf>
41. Canadian Society of Hospital Pharmacists, (CSHP). Guidelines for Handling and Disposal of Hazardous Pharmaceuticals (Including Cytotoxic Drugs). : CSHP; -4 2003. p. 31-42
42. American Society of Health-System Pharmacists. ASHP Technical Assistance Bulletin on Handling Cytotoxic and Hazardous Drugs. *Am.J.Hosp.Pharm.* 1990;47(5):1033-1049
43. Bartel SB. Safe practices and financial considerations in using oral chemotherapeutic agents. *Am.J.Health.Syst.Pharm.* 2007;64(9 Suppl 5):S8-S14
44. Andrea Griffiths, Occupational Hygienist, Prevention Advisor, MAC, PhD. Recommended Strategy Change Out Schedule for Canisters Used in Pharmacy for: Decontaminating/Cleaning of Biological Safety Cabinet and Spill Clean Up.
45. BCCA Provincial Systemic Program Committee. BCCA Systemic Policy Number III-10 - Systemic Therapy Treatment Delivery Process. BCCA, 2009. Available at: <http://www.bccancer.bc.ca/HPI/ChemotherapyProtocols/Policies.htm>. Accessed June 15
46. ASHP Council on Professional Affairs. ASHP guidelines on preventing medication errors with antineoplastic agents. *Am.J.Health.Syst.Pharm.* 2002;59(17):1648-1668
47. WHO. World Health Organization Information Exchange System Alert No. 115. 2007;115
48. BCCA Provincial Pharmacy Safety Officer/P4C. BCCA Pharmacy Directive Number VI-90-02 - Outpatient Prescription Labels: Guiding Principles for Patient Instructions. BCCA 2011
49. SHPA Committee of Specialty Practice in Cancer Services. SHPA Standards of Practice for the Provision of Oral Chemotherapy for the Treatment of Cancer. *Journal of Pharmacy Practice and Research* 2007;37(2):149-152
50. Canadian Society of Hospital Pharmacists, (CSHP). Guidelines for Preparing Medications for Natural Rubber Latex (NRL) Sensitive/Allergic Patients. : CSHP; -4 2003. p. 107-111