



Statistics by Cancer Type - Hodgkin Lymphoma

General Summary

In 2017, an estimated 120 British Columbians will be newly diagnosed with Hodgkin's lymphoma and 15 will die of it. One in 581 females and one in 474 males is expected to develop Hodgkin's lymphoma during their lifetime. One in 4776 females and one in 2892 males is expected to die of Hodgkin's lymphoma.

Number of New Hodgkin Lymphoma Cancer Cases Diagnosed in 2014

| Age at Diagnosis | Males | Females | Total |
|------------------|-----------|-----------|------------|
| 0-19 | 15 | 5 | 15 |
| 20-39 | 35 | 20 | 55 |
| 40-59 | 15 | 10 | 25 |
| 60-79 | 10 | 5 | 15 |
| 80+ | 5 | 5 | 10 |
| Total | 70 | 40 | 115 |

Due to rounding, totals may not equal the sum of individual values

Number of Hodgkin Lymphoma Cancer Deaths in 2014

| Age at Death | Males | Females | Total |
|--------------|-----------|----------|-----------|
| 0-19 | 0 | 0 | 0 |
| 20-39 | 0 | 0 | 5 |
| 40-59 | 0 | 5 | 5 |
| 60-79 | 10 | 0 | 10 |
| 80+ | 0 | 0 | 5 |
| Total | 10 | 5 | 20 |

Due to rounding, totals may not equal the sum of individual values

Incidence Rates (per 100,000 population) for Hodgkin Lymphoma Cancer Cases Diagnosed in 2014

| Age at Diagnosis | Males | Females | Total |
|------------------|------------|------------|------------|
| 0-19 | 2.2 | 0.9 | 1.6 |
| 20-39 | 5.1 | 3.1 | 4.1 |
| 40-59 | 2.3 | 1.3 | 1.8 |
| 60-79 | 2.5 | 0.9 | 1.7 |
| 80+ | 3.5 | 4.1 | 3.9 |
| Total | 3.1 | 1.8 | 2.4 |

Mortality Rates (per 100,000 population) for Hodgkin Lymphoma Cancer Deaths in 2014

| Age at Death | Males | Females | Total |
|--------------|------------|------------|------------|
| 0-19 | 0 | 0 | 0 |
| 20-39 | 0.3 | 0.5 | 0.4 |
| 40-59 | 0.3 | 0.3 | 0.3 |
| 60-79 | 1.9 | 0.2 | 1 |
| 80+ | 1.2 | 0.8 | 1 |
| Total | 0.6 | 0.3 | 0.4 |



Statistics by Cancer Type - Hodgkin Lymphoma

Estimated New Hodgkin Lymphoma Cancer Diagnoses in 2017 by Health Authority

| Health Authority | Age at Diagnosis | | | | | Total* |
|------------------|------------------|-----------|-----------|-----------|----------|------------|
| | 0-19 | 20-39 | 40-59 | 60-79 | 80+ | |
| Interior | 0 | 10 | 5 | 5 | 0 | 20 |
| Fraser | 5 | 15 | 10 | 10 | 0 | 40 |
| Coastal | 0 | 15 | 5 | 5 | 0 | 25 |
| Island | 0 | 10 | 0 | 5 | 0 | 20 |
| Northern | 0 | 0 | 0 | 0 | 0 | 5 |
| BC | 10 | 50 | 25 | 25 | 5 | 120 |

Due to rounding and unknown health authority of residence at time of diagnosis for some cases, totals may not equal the sum of individual values

Estimated Hodgkin Lymphoma Cancer Deaths in 2017 by Health Authority

| Health Authority | Age at Diagnosis | | | | | Total* |
|------------------|------------------|----------|----------|----------|----------|-----------|
| | 0-19 | 20-39 | 40-59 | 60-79 | 80+ | |
| Interior | 0 | 0 | 0 | 0 | 0 | 5 |
| Fraser | 0 | 0 | 0 | 0 | 0 | 5 |
| Coastal | 0 | 0 | 0 | 0 | 0 | 5 |
| Island | 0 | 0 | 0 | 0 | 0 | 5 |
| Northern | 0 | 0 | 0 | 0 | 0 | 0 |
| BC | 0 | 0 | 0 | 5 | 0 | 15 |

Due to rounding and unknown health authority of residence at time of death for some cases, totals may not equal the sum of individual values

1-, 3- and 5-year Observed Survival and Age-standardized Relative Survival for BC Adults (age 15-99) Diagnosed with Hodgkin Lymphoma Cancer

| | Observed Survival Proportion % | | | Relative Survival Proportion % | | |
|-------------------------|--------------------------------|-------------|-------------|--------------------------------|-------------|-------------|
| | 1 Year | 3 Year | 5 Year | 1 Year | 3 Year | 5 Year |
| Survival Proportion | 91.6 | 88.8 | 84.5 | 92.4 | 90.6 | 86.1 |
| 95% Confidence Interval | 88.6 - 93.9 | 85.3 - 91.4 | 80.6 - 87.6 | 89.6 - 94.5 | 87.2 - 93.1 | 81.6 - 89.5 |