

Statistics by Cancer Type - Hodgkin Lymphoma

General Summary

In 2019, an estimated 120 British Columbians will be newly diagnosed with Hodgkin's lymphoma and 15 will die of it. One in 544 females and one in 445 males is expected to develop Hodgkin's lymphoma during their lifetime. One in 4605 females and one in 3303 males is expected to die of Hodgkin's lymphoma.

Number of New Hodgkin Lymphoma Cancer Cases Diagnosed in 2016

Age at Diagnosis	Males	Females	Total
0-19	5	5	5
20-39	25	25	45
40-59	15	15	25
60-79	15	15	25
80+	0	0	0
Total	55	55	105

Due to rounding, totals may not equal the sum of individual values

Number of Hodgkin Lymphoma Cancer Deaths in 2016

Age at Death	Males	Females	Total
0-19	0	0	0
20-39	0	0	0
40-59	0	5	0
60-79	5	0	5
80+	0	0	0
Total	5	0	10

Due to rounding, totals may not equal the sum of individual values

Incidence Rates (per 100,000 population) for Hodgkin Lymphoma Cancer Cases Diagnosed in 2016

Age at Diagnosis	Males	Females	Total
0-19	0.6	1.1	0.8
20-39	3.4	3.7	3.6
40-59	2.3	1.8	2.0
60-79	2.8	2.5	2.6
80+	1.1	0.8	0.9
Total	2.3	2.3	2.3

Mortality Rates (per 100,000 population) for Hodgkin Lymphoma Cancer Deaths in 2016

Age at Death	Males	Females	Total
0-19	0.0	0.0	0.0
20-39	0.0	0.0	0.0
40-59	0.0	0.1	0.1
60-79	0.9	0.2	0.5
80+	0.0	1.6	0.9
Total	0.2	0.2	0.2

Statistics by Cancer Type - Hodgkin Lymphoma

Estimated New Hodgkin Lymphoma Cancer Diagnoses in 2019 by Health Authority

Health Authority	Age at Diagnosis					Total*
	0-19	20-39	40-59	60-79	80+	
Interior	5	10	5	10	5	20
Fraser	0	20	10	5	0	40
Coastal	5	20	5	10	0	30
Island	5	10	5	5	0	20
Northern	0	5	5	0	0	5
BC	15	55	25	30	5	120

*Due to rounding and unknown health authority of residence at time of diagnosis for some cases, totals may not equal the sum of individual values

Estimated Hodgkin Lymphoma Cancer Deaths in 2019 by Health Authority

Health Authority	Age at Diagnosis					Total*
	0-19	20-39	40-59	60-79	80+	
Interior	0	0	0	0	0	0
Fraser	0	0	0	5	0	5
Coastal	0	0	0	0	0	0
Island	0	0	0	5	0	0
Northern	0	0	0	0	0	0
BC	0	0	10	10	0	15

*Due to rounding and unknown health authority of residence at time of death for some cases, totals may not equal the sum of individual values

1-, 3- and 5-year Observed Survival and Age-standardized Relative Survival for BC Adults (age 15-99) Diagnosed with Hodgkin Lymphoma Cancer

	Observed Survival Proportion %			Relative Survival Proportion %		
	1 Year	3 Year	5 Year	1 Year	3 Year	5 Year
Survival Proportion	92.0	88.5	84.2	92.5	89.9	86.0
95% Confidence Interval	89.2 - 94.2	85.2 - 91.1	80.5 - 87.3	89.8 - 94.5	86.7 - 92.4	82.2 - 89.1