



Statistics by Cancer Type - Non-Hodgkin Lymphoma

General Summary

In 2017, an estimated 1200 British Columbians will be newly diagnosed with Non-Hodgkin lymphoma and 350 will die of it. One in 47 females and one in 39 males is expected to develop Non-Hodgkin's lymphoma during their lifetime. One in 126 females and one in 102 males is expected to die of Non-Hodgkin's lymphoma.

Number of New Non-Hodgkin Lymphoma Cancer Cases Diagnosed in 2014

Age at Diagnosis	Males	Females	Total
0-19	15	0	10
20-39	30	20	45
40-59	140	95	240
60-79	340	240	585
80+	135	135	270
Total	660	495	1155

Due to rounding, totals may not equal the sum of individual values

Number of Non-Hodgkin Lymphoma Cancer Deaths in 2014

Age at Death	Males	Females	Total
0-19	0	5	0
20-39	5	0	5
40-59	20	10	35
60-79	105	55	165
80+	60	60	120
Total	190	125	320

Due to rounding, totals may not equal the sum of individual values

Incidence Rates (per 100,000 population) for Non-Hodgkin Lymphoma Cancer Cases Diagnosed in 2014

Age at Diagnosis	Males	Females	Total
0-19	2.6	0.2	1.5
20-39	4.8	2.9	3.8
40-59	21.4	14.5	17.9
60-79	79.3	53.9	66.4
80+	157.5	110.8	130
Total	28.7	21.2	24.9

Mortality Rates (per 100,000 population) for Non-Hodgkin Lymphoma Cancer Deaths in 2014

Age at Death	Males	Females	Total
0-19	0.2	0	0.1
20-39	0.5	0.2	0.3
40-59	3.3	1.5	2.4
60-79	24.7	12.5	18.5
80+	69.4	50.1	58
Total	8.3	5.5	6.9



Statistics by Cancer Type - Non-Hodgkin Lymphoma

Estimated New Non-Hodgkin Lymphoma Cancer Diagnoses in 2017 by Health Authority

Health Authority	Age at Diagnosis					Total*
	0-19	20-39	40-59	60-79	80+	
Interior	0	10	45	140	55	240
Fraser	5	15	85	205	80	390
Coastal	0	10	60	135	60	270
Island	0	10	45	140	60	250
Northern	0	0	15	35	10	70
BC	10	45	250	660	260	1200

Due to rounding and unknown health authority of residence at time of diagnosis for some cases, totals may not equal the sum of individual values

Estimated Non-Hodgkin Lymphoma Cancer Deaths in 2017 by Health Authority

Health Authority	Age at Diagnosis					Total*
	0-19	20-39	40-59	60-79	80+	
Interior	0	0	0	35	35	70
Fraser	0	0	10	55	45	110
Coastal	0	0	5	35	30	70
Island	0	0	5	40	35	80
Northern	0	0	0	10	10	15
BC	0	0	25	165	155	350

Due to rounding and unknown health authority of residence at time of death for some cases, totals may not equal the sum of individual values

1-, 3- and 5-year Observed Survival and Age-standardized Relative Survival for BC Adults (age 15-99) Diagnosed with Non-Hodgkin Lymphoma Cancer

	Observed Survival Proportion %			Relative Survival Proportion %		
	1 Year	3 Year	5 Year	1 Year	3 Year	5 Year
Survival Proportion	79.2	68.9	61.6	82.0	74.9	70.4
95% Confidence Interval	77.4 - 80.9	66.8 - 70.8	59.4 - 63.7	80.3 - 83.6	72.8 - 76.9	68.1 - 72.7