

## Statistics by Cancer Type - Non-Hodgkin Lymphoma

### General Summary

In 2020, an estimated 1320 British Columbians will be newly diagnosed with Non-Hodgkin lymphoma. One in 46 females and one in 39 males is expected to develop Non-Hodgkin's lymphoma during their lifetime. One in 128 females and one in 94 males is expected to die of Non-Hodgkin's lymphoma.

### Number of New Non-Hodgkin Lymphoma Cancer Cases Diagnosed in 2017

Age at Diagnosis	Males	Females	Total
0-19	10	0	10
20-39	25	25	45
40-59	135	95	230
60-79	345	260	605
80+	130	125	255
<b>Total</b>	<b>645</b>	<b>505</b>	<b>1150</b>

Due to rounding, totals may not equal the sum of individual values

### Number of Non-Hodgkin Lymphoma Cancer Deaths in 2017

Age at Death	Males	Females	Total
0-19	5	0	5
20-39	5	0	5
40-59	15	10	30
60-79	105	75	180
80+	105	90	190
<b>Total</b>	<b>235</b>	<b>175</b>	<b>410</b>

Due to rounding, totals may not equal the sum of individual values

### Incidence Rates (per 100,000 population) for Non-Hodgkin Lymphoma Cancer Cases Diagnosed in 2017

Age at Diagnosis	Males	Females	Total
0-19	1.6	0.2	0.9
20-39	4.0	3.3	3.7
40-59	19.8	13.4	16.5
60-79	71.1	50.6	60.5
80+	141.1	100.5	117.5
<b>Total</b>	<b>26.4</b>	<b>20.3</b>	<b>23.3</b>

### Mortality Rates (per 100,000 population) for Non-Hodgkin Lymphoma Cancer Deaths in 2017

Age at Death	Males	Females	Total
0-19	0.2	0.0	0.1
20-39	0.7	0.2	0.4
40-59	2.7	1.8	2.2
60-79	22.1	14.5	18.2
80+	113.8	70.4	88.6
<b>Total</b>	<b>9.6</b>	<b>7.2</b>	<b>8.4</b>

## Statistics by Cancer Type - Non-Hodgkin Lymphoma

### Estimated New Non-Hodgkin Lymphoma Cancer Diagnoses in 2020 by Health Authority

Health Authority	Age at Diagnosis					Total*
	0-19	20-39	40-59	60-79	80+	
Interior	0	10	40	150	55	260
Fraser	0	20	90	230	95	425
Coastal	0	15	50	145	65	280
Island	0	10	40	165	65	280
Northern	0	5	15	45	10	75
<b>BC</b>	<b>10</b>	<b>60</b>	<b>230</b>	<b>735</b>	<b>295</b>	<b>1320</b>

\*Due to rounding and unknown health authority of residence at time of diagnosis for some cases, totals may not equal the sum of individual values

### 1-, 3- and 5-year Observed Survival and Age-standardized Relative Survival for BC Adults (age 15-99) Diagnosed with Non-Hodgkin Lymphoma Cancer

	Observed Survival Proportion %			Relative Survival Proportion %		
	1 Year	3 Year	5 Year	1 Year	3 Year	5 Year
Survival Proportion	79.5	70.0	63.6	82.7	76.3	72.6
95% Confidence Interval	77.7 - 81.1	68.0 - 71.9	61.5 - 65.6	81.1 - 84.3	74.2 - 78.3	70.2 - 74.9