# Quality Breast Cancer Care: Can we do better?

Rona Cheifetz SON Fall Update October 24, 2009

# Do we need to improve breast cancer care in BC?

# How can we improve breast cancer care in BC?

 Review some of literature 'guidelines' and specialist recommendations

National Library of Medicine, Agency for Health Care Research and Quality Oct 2004 (for example)

- Discuss whether goal is important
- Discuss barriers
- Discuss requirements to achieve goal

 If a palpable breast mass is detected at least one of the following procedures should be completed within 3 months: FNA, Mammo, US, Bx or F/U visit

- Important?
- Barriers?
- Requirements to achieve?

 Tissue diagnosis should be obtained at the initial specialty visit for a palpable breast mass

- Important?
- Barriers?
- Requirements to achieve?

 If an FNA cannot rule out malignancy a tissue diagnosis should be obtained within 6 weeks

- Important?
- Barriers?
- Requirements to achieve?

 Suspicious screening mammography should be evaluated within 3 weeks with further assessment

- Important?
- Barriers?
- Requirements to achieve?

 Women with suspicious imaging requiring surgery for diagnosis should be operated on within 14 days of the surgical decision

- Important?
- Barriers?
- Requirements to achieve?

 Core biopsy rather than surgical biopsy should be procedure of choice for initial breast cancer diagnosis

- Important?
- Barriers?
- Requirements to achieve?

 Urgent referrals with confirmed breast cancer should be seen within 5 working days

- Important?
- Barriers?
- Requirements to achieve?

 As appropriate, women with stage 1 or 2 breast cancer should be offered the choice of mastectomy or breast conservation

- Important?
- Barriers?
- Requirements to achieve?

 Therapeutic surgery should be available within 21 days of surgical consultation for breast cancer

- Important?
- Barriers?
- Requirements to achieve?

 Women have breast conserving surgery should not require more than 2 therapeutic operations

- Important?
- Barriers?
- Requirements to achieve?

 Women having breast conserving surgery, should have their biopsy cavities marked with clips to facilitate radiation planning

- Important?
- Barriers?
- Requirements to achieve?

 Operative reports should contain details of the margins (eg pectoralis fascia was taken, very thin anterior margin)

- Important?
- Barriers?
- Requirements to achieve?

 Women who are clinically node negative, should be offered sentinel node biopsy for axillary staging (unless contra-indicated)

- Important?
- Barriers?
- Requirements to achieve?

 All women with curable breast cancer should receive an opinion regarding adjuvant therapy

- Important?
- Barriers?
- Requirements to achieve?

 Early referral or discussion, especially for persons with LABC or with a definite indication for chemotherapy

- Important?
- Barriers?
- Requirements to achieve?

 Early referral for persons with medical issues and/or social issues that will impact their breast cancer care

- Important?
- Barriers?
- Requirements to achieve?