

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca/terms-of-use and according to acceptable standards of care.

PROTOCOL CODE: GUBEP

Page 1 of 2

DOCTOR'S ORDERS Htcm Wtkg BSA_	m²	
REMINDER: Please ensure drug allergies and previous bleomycin are documented on t	he Allergy & Alert Form	
	Cycle #:	
Date of Previous Cycle:		
No Treatment Delay for Day 1 Bloodwork.		
Dose modification for: Other Toxicity		
Proceed with treatment based on blood work from		
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm		
ondansetron 8 mg PO 30 to 60 minutes prior to treatment on Days 1 to 5		
dexamethasone 8 mg or 12 mg (select one) PO 30 to 60 minutes prior to treatment on Day 1 ; then		
dexamethasone 4 mg PO BID on Days 2 to 5		
aprepitant 125 mg PO 30 to 60 minutes prior to treatment on Day 1; then 80 mg PO	daily on Day 2 and 3	
hydrocortisone 100 mg IV prior to bleomycin on Day 1, Day 8, and Day 15		
hydrocortisone 100 mg IV prior to etoposide or etoposide phosphate (ETOPOPH	· · · · · · · · · · · · · · · · · · ·	
diphenhydrAMINE 50 mg IV prior to etoposide or etoposide phosphate (ETOPOP	,	
Have Hypersensitivity Reaction Tray and Protocol Available		
PRE-HYDRATION:		
1000 mL NS with 20 mEq potassium chloride and 2 g magnesium sulfate over 1 hour	prior to CISplatin	
CHEMOTHERAPY:		
CISplatin 20 mg/m²/day x BSA = mg		
☐ Dose Modification:% =mg/m² x BSA = mg		
IV in 100 mL NS over 30 minutes on Days 1 to 5		
etoposide 100 mg/m²/day x BSA x = mg		
☐ Dose Modification: % = mg/m² x BSA = mg		
IV in 250 to 1000 mL (non-DEHP bag) NS over 45 min to 1 hour 30 min on Days 1	to 5 (use non-DEHP	
tubing with 0.2 micron in-line filter).		
bleomycin = units (dose is 30 units)* IV in 50 mL NS over 10 minutes Day 1, Day 8, and Day 15		
*bleomycin dose will need to be filled in with suggested dosing. Any dose modifications can	result in inferior outcomes.	
If hypersensitivity to etoposide:		
Omit etoposide. Give etoposide phosphate (ETOPOPHOS)* 100 mg/m²/day x BSA = mg		
□ Dose Modification: % = mg/m² x BSA = mg		
IV in 500 mL NS over 45 min to 1 hour 30 min on Days 1 to 5 .		
*Approval from the Health Canada Special Access Program must be obtained for eacl	n natient	
POST-HYDRATION: 500 mL NS over 30 minutes to 1 hour post chemotherapy on Days 1 to 5		
STANDING ORDER FOR ETOPOSIDE OR ETOPOSIDE PHOSPHATE (ETOPOPHOS) TOXICITY:		
hydrocortisone 100 mg IV prn / diphenhydrAMINE 50 mg IV prn	IONICIT:	
DOCTOR'S SIGNATURE:	SIGNATURE:	
	UC:	



Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca/terms-of-use and according to acceptable standards of care.

PROTOCOL CODE: GUBEP

Page 2 of 2

DOCTOR'S ORDERS		
DATE:		
RETURN APPOINTMENT ORDERS		
☐ Return in three weeks for Doctor and Cycle Book chemo Days 1 to 5, Day 8 & 15☐ Last Cycle. Return in week(s).		
CBC & Diff, Platelets, Creatinine, LDH, AFP, beta hCG tumour marker, magnesium, sodium, potassium, random glucose prior to each cycle		
Creatinine within 24 hours prior to Day 8 and Day 15, if patient receiving bleomycin		
CBC & Diff, Platelets on Day 5, if ANC on Day 1 less than 1.0 x 109/L		
Creatinine on Day 5, if creatinine on Day 1 greater than upper limit of normal		
☐ Other tests:☐ Consults:		
DOCTOR'S SIGNATURE:	SIGNATURE:	
	UC:	