

PROTOCOL CODE: LUAJPMB6

(Page 1 of 2)

DOCTOR'S ORDERS		Ht _____ cm	Wt _____ kg	BSA _____	m^2						
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form											
DATE:	To be given:			Cycle #:							
Date of Previous Cycle:											
<input type="checkbox"/> Delay treatment _____ week(s)											
May proceed with dose as written if within 96 hours ALT less than or equal to 3 times the upper limit of normal, total bilirubin less than or equal to 1.5 times the upper limit of normal , creatinine less than or equal to 1.5 times the upper limit of normal and less than or equal to 1.5 times baseline .											
Proceed with treatment based on blood work from _____											
PREMEDIcATIONS: Patient to take own supply. RN/Pharmacist to confirm _____.											
For prior infusion reaction:											
<input type="checkbox"/> diphenhydRamine 50 mg PO 30 minutes prior to treatment <input type="checkbox"/> acetaminophen 325 to 975 mg PO 30 minutes prior to treatment <input type="checkbox"/> hydrocortisone 25 mg IV 30 minutes prior to treatment											
TREATMENT:											
pembrolizumab 4 mg/kg x _____ kg = _____ mg (max. 400 mg) every 6 weeks IV in 50 mL NS over 30 minutes using a 0.2 micron in-line filter											
Pharmacist to select dose band per last page of PPO. Complete table below (please print)											
<table border="1"> <thead> <tr> <th>Drug</th> <th>Dose Band (mg)</th> <th>Pharmacist Initial and Date</th> </tr> </thead> <tbody> <tr> <td>pembrolizumab</td> <td></td> <td></td> </tr> </tbody> </table>						Drug	Dose Band (mg)	Pharmacist Initial and Date	pembrolizumab		
Drug	Dose Band (mg)	Pharmacist Initial and Date									
pembrolizumab											
RETURN APPOINTMENT ORDERS											
<input type="checkbox"/> Return in six weeks for Doctor and Cycle _____											
<input type="checkbox"/> Last cycle. Return in _____ week(s)											
CBC & Diff, creatinine, alkaline phosphatase, ALT, total bilirubin, LDH, sodium, potassium, TSH prior to each treatment											
If clinically indicated: <input type="checkbox"/> ECG <input type="checkbox"/> chest X-ray <input type="checkbox"/> serum HCG or <input type="checkbox"/> urine HCG – required for woman of child bearing potential <input type="checkbox"/> Free T3 and free T4 <input type="checkbox"/> lipase <input type="checkbox"/> morning serum cortisol <input type="checkbox"/> random glucose <input type="checkbox"/> serum ACTH levels <input type="checkbox"/> testosterone <input type="checkbox"/> estradiol <input type="checkbox"/> FSH <input type="checkbox"/> LH											
<input type="checkbox"/> Weekly nursing assessment <input type="checkbox"/> Other consults: <input type="checkbox"/> See general orders sheet for additional requests.											
DOCTOR'S SIGNATURE:				SIGNATURE:							
				UC:							

PROTOCOL CODE: LUAJPMB6

(Page 2 of 2)

PEMBROLIZUMAB DOSE BANDING TABLE (4 mg/kg capped 400 mg)

Ordered Dose (mg)		Rounded dose (mg)
From:	To:	
Less than 137.5		Pharmacy prepares specific dose
137.5	162.49	150
162.5	187.49	175
187.5	221.49	200
221.5	242.49	225
242.5	264.49	250
265.5	284.49	275
285.5	332.49	300
333.5	374.49	350
374.5	400	400